



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 01:26 am BST

PDB ID : 2VIR
Title : INFLUENZA VIRUS HEMAGGLUTININ COMPLEXED WITH A NEUTRALIZING ANTIBODY
Authors : Bizebard, T.; Fleury, D.; Gigant, B.; Wharton, S.A.; Skehel, J.J.; Knossow, M.
Deposited on : 1997-12-22
Resolution : 3.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1251 (3.30-3.22)
Ramachandran outliers	138981	1229 (3.30-3.22)
Sidechain outliers	138945	1228 (3.30-3.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	210	
2	B	221	
3	C	282	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5339 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IMMUNOGLOBULIN (IGG1, LAMBDA).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	210	1589	994	266	323	6	0	0	0

- Molecule 2 is a protein called IMMUNOGLOBULIN (IGG1, LAMBDA).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	221	1672	1065	270	328	9	0	0	0

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	3	GLN	LYS	CONFLICT	GB 4096752
B	5	LYS	GLN	CONFLICT	GB 4096752
B	28	LEU	SER	CONFLICT	GB 4096752
B	30	ILE	THR	CONFLICT	GB 4096752
B	32	ASN	TYR	CONFLICT	GB 4096752
B	63	LEU	HIS	CONFLICT	GB 4096752
B	69	ILE	PHE	CONFLICT	GB 4096752
B	83	LYS	ASN	CONFLICT	GB 4096752
B	92	MET	LEU	CONFLICT	GB 4096752
B	98	ASP	-	INSERTION	GB 4096752
B	99	PHE	-	INSERTION	GB 4096752
B	100	TYR	-	INSERTION	GB 4096752
B	102	TYR	HIS	CONFLICT	GB 4096752
B	103	ASP	GLY	CONFLICT	GB 4096752
B	105	PHE	-	INSERTION	GB 4096752
B	106	TYR	-	INSERTION	GB 4096752
B	107	TYR	-	INSERTION	GB 4096752
B	108	ALA	-	INSERTION	GB 4096752
B	109	MET	-	INSERTION	GB 4096752
B	110	ASP	-	INSERTION	GB 4096752

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	117	SER	LEU	CONFLICT	GB 4096752
B	122	SER	ALA	CONFLICT	GB 4096752
B	135	PRO	SER	CONFLICT	GB 4096752

- Molecule 3 is a protein called HEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	267	2075	1304	360	400	11	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	38	ASP	ASN	CONFLICT	UNP P03437
C	81	ASP	ASN	MODIFIED RESIDUE	UNP P03437
C	165	ASP	ASN	MODIFIED RESIDUE	UNP P03437
C	285	ASP	ASN	MODIFIED RESIDUE	UNP P03437

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total 2	Zn 2	0	0
4	C	1	Total 1	Zn 1	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: IMMUNOGLOBULIN (IGG1, LAMBDA)



- Molecule 2: IMMUNOGLOBULIN (IGG1, LAMBDA)



- Molecule 3: HEMAGGLUTININ





4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	85.50Å 85.50Å 515.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	7.00 – 3.25	Depositor
% Data completeness (in resolution range)	(Not available) (7.00-3.25)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.84	Depositor
R, R_{free}	0.198 , 0.249	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5339	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/1627	0.87	2/2224 (0.1%)
2	B	0.59	0/1718	0.83	1/2351 (0.0%)
3	C	0.61	0/2127	0.84	1/2896 (0.0%)
All	All	0.60	0/5472	0.85	4/7471 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	45	LEU	CA-CB-CG	6.62	130.53	115.30
2	B	153	GLY	N-CA-C	5.72	127.40	113.10
3	C	108	LEU	CA-CB-CG	5.16	127.16	115.30
1	A	45	LEU	N-CA-C	-5.01	97.48	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	143	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1589	0	1532	70	0
2	B	1672	0	1635	75	0
3	C	2075	0	2011	108	0
4	A	2	0	0	0	0
4	C	1	0	0	0	0
All	All	5339	0	5178	247	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 247 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:134:ALA:HB2	2:B:219:ILE:HD11	1.40	1.03
1:A:147:VAL:HG12	1:A:200:HIS:HB2	1.47	0.97
3:C:283:THR:HG23	3:C:285:ASP:H	1.33	0.93
2:B:40:PRO:HG2	2:B:43:LYS:HB2	1.51	0.91
3:C:45:SER:HB3	3:C:296:ASN:OD1	1.81	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	208/210 (99%)	183 (88%)	19 (9%)	6 (3%)	4 25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	219/221 (99%)	189 (86%)	26 (12%)	4 (2%)	8	35
3	C	265/282 (94%)	232 (88%)	25 (9%)	8 (3%)	4	24
All	All	692/713 (97%)	604 (87%)	70 (10%)	18 (3%)	5	27

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	87	THR
2	B	173	HIS
2	B	212	SER
3	C	210	GLN
2	B	153	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/178 (100%)	158 (89%)	20 (11%)	6	23
2	B	192/192 (100%)	180 (94%)	12 (6%)	18	47
3	C	236/250 (94%)	208 (88%)	28 (12%)	5	20
All	All	606/620 (98%)	546 (90%)	60 (10%)	8	28

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	183	LEU
3	C	48	THR
3	C	264	LYS
2	B	218	LYS
3	C	57	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	32	ASN
2	B	142	ASN
3	C	210	GLN
2	B	3	GLN
3	C	189	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.