



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 8, 2023 – 04:11 am GMT

PDB ID : 2W6G
Title : Low resolution structures of bovine mitochondrial F1-ATPase during controlled dehydration: Hydration State 3.
Authors : Sanchez-Weatherby, J.; Felisaz, F.; Gobbo, A.; Huet, J.; Ravelli, R.B.G.; Bowler, M.W.; Cipriani, F.
Deposited on : 2008-12-18
Resolution : 6.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 6.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1049 (8.00-3.90)
Ramachandran outliers	138981	1016 (8.00-3.86)
Sidechain outliers	138945	1017 (8.00-3.82)
RSRZ outliers	127900	1015 (8.20-3.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	553	 47% 33% 7% • 12%
1	B	553	 47% 33% 6% • 13%
1	C	553	 51% 33% • • 11%
2	D	528	 55% 28% • • 12%
2	E	528	 45% 36% 6% • 12%
2	F	528	 57% 28% • 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	298	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment (3%), a green segment (34%), a yellow segment (12%), and a large grey segment (53%).</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 22795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP SYNTHASE SUBUNIT ALPHA HEART ISOFORM, MITOCHONDRIAL.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	487	Total 3715	C 2341	N 656	O 706	S 12	0	0	0
1	B	479	Total 3656	C 2303	N 647	O 694	S 12	0	0	0
1	C	492	Total 3748	C 2360	N 661	O 715	S 12	0	0	0

- Molecule 2 is a protein called ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	467	Total 3539	C 2243	N 601	O 684	S 11	0	0	0
2	E	466	Total 3530	C 2238	N 600	O 681	S 11	0	0	0
2	F	466	Total 3530	C 2238	N 600	O 681	S 11	0	0	0

- Molecule 3 is a protein called ATP SYNTHASE SUBUNIT GAMMA, MITOCHONDRIAL.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	G	140	Total 1077	C 671	N 195	O 205	S 6	0	0	0




• Molecule 1: ATP SYNTHASE SUBUNIT ALPHA HEART ISOFORM, MITOCHONDRIAL





• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL






• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL



• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	109.35Å 132.93Å 275.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	101.53 – 6.00 80.74 – 6.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (101.53-6.00) 65.8 (80.74-6.00)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 6.18Å)	Xtrriage
Refinement program	REFMAC 5.5.0038	Depositor
R, R_{free}	0.330 , (Not available) 0.329 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	156.4	Xtrriage
Anisotropy	0.372	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.36$, $\langle L^2 \rangle = 0.20$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.78	EDS
Total number of atoms	22795	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/3766	1.31	29/5080 (0.6%)
1	B	0.61	0/3704	1.36	34/4995 (0.7%)
1	C	0.62	0/3799	1.38	24/5126 (0.5%)
2	D	0.62	0/3596	1.36	23/4879 (0.5%)
2	E	0.60	0/3587	1.32	18/4867 (0.4%)
2	F	0.62	0/3587	1.36	27/4867 (0.6%)
3	G	0.33	0/1083	0.69	2/1448 (0.1%)
All	All	0.60	0/23122	1.33	157/31262 (0.5%)

There are no bond length outliers.

The worst 5 of 157 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	408	ARG	CD-NE-CZ	23.30	156.22	123.60
2	E	408	ARG	CD-NE-CZ	13.97	143.16	123.60
1	C	291	ARG	NE-CZ-NH2	-13.30	113.65	120.30
1	B	40	ARG	NE-CZ-NH1	13.08	126.84	120.30
1	B	279	ARG	NE-CZ-NH1	12.73	126.67	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3715	0	3813	184	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3656	0	3764	156	58
1	C	3748	0	3843	163	56
2	D	3539	0	3593	158	0
2	E	3530	0	3587	217	0
2	F	3530	0	3587	124	2
3	G	1077	0	1139	66	0
All	All	22795	0	23326	961	58

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 961 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:390:ILE:HD11	3:G:242:MET:CE	1.54	1.36
2:F:390:ILE:CD1	3:G:242:MET:CE	2.07	1.29
2:E:390:ILE:HG13	3:G:25:MET:SD	1.75	1.25
1:C:127:ARG:HH12	1:C:255:GLU:HB2	1.00	1.13
2:F:390:ILE:CD1	3:G:242:MET:HE3	1.77	1.09

The worst 5 of 58 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:314:ASP:O	1:C:22:SER:C[4_455]	0.42	1.78
1:B:314:ASP:CA	1:C:22:SER:N[4_455]	0.82	1.38
1:B:318:GLY:C	1:C:20:ASP:OD2[4_455]	0.87	1.33
1:B:314:ASP:O	1:C:23:VAL:N[4_455]	0.93	1.27
1:B:314:ASP:OD1	1:C:20:ASP:O[4_455]	0.94	1.26

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/553 (88%)	443 (91%)	35 (7%)	7 (1%)	11	46
1	B	475/553 (86%)	427 (90%)	41 (9%)	7 (2%)	10	45
1	C	490/553 (89%)	444 (91%)	38 (8%)	8 (2%)	9	44
2	D	465/528 (88%)	419 (90%)	43 (9%)	3 (1%)	25	66
2	E	464/528 (88%)	407 (88%)	48 (10%)	9 (2%)	8	38
2	F	464/528 (88%)	432 (93%)	30 (6%)	2 (0%)	34	72
3	G	134/298 (45%)	130 (97%)	3 (2%)	1 (1%)	22	62
All	All	2977/3541 (84%)	2702 (91%)	238 (8%)	37 (1%)	13	50

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	407	GLY
2	E	393	MET
1	A	57	SER
1	A	405	GLN
1	A	409	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/444 (88%)	351 (89%)	42 (11%)	6	24
1	B	388/444 (87%)	341 (88%)	47 (12%)	5	20
1	C	397/444 (89%)	369 (93%)	28 (7%)	14	39
2	D	377/417 (90%)	346 (92%)	31 (8%)	11	34
2	E	376/417 (90%)	344 (92%)	32 (8%)	10	33
2	F	376/417 (90%)	354 (94%)	22 (6%)	19	45
3	G	116/251 (46%)	110 (95%)	6 (5%)	23	48
All	All	2423/2834 (86%)	2215 (91%)	208 (9%)	10	33

5 of 208 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	502	THR
2	D	405	SER
2	F	405	SER
2	D	56	SER
2	D	232	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	328	HIS
2	F	443	GLN
2	E	51	GLN
3	G	225	GLN
2	F	221	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	487/553 (88%)	0.00	7 (1%) 75 66	20, 20, 20, 20	0
1	B	479/553 (86%)	-0.01	2 (0%) 92 87	20, 20, 20, 20	0
1	C	492/553 (88%)	-0.14	1 (0%) 95 93	20, 20, 20, 20	0
2	D	467/528 (88%)	-0.06	3 (0%) 89 84	20, 20, 20, 20	0
2	E	466/528 (88%)	-0.13	3 (0%) 89 84	20, 20, 20, 20	0
2	F	466/528 (88%)	-0.06	4 (0%) 84 77	20, 20, 20, 20	0
3	G	140/298 (46%)	0.48	9 (6%) 19 18	36, 59, 81, 89	0
All	All	2997/3541 (84%)	-0.04	29 (0%) 82 75	20, 20, 20, 89	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	67	LEU	4.0
3	G	202	ARG	3.5
1	A	234	ALA	3.4
1	B	489	ILE	3.2
3	G	69	ILE	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.