



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2025 – 06:49 am BST

PDB ID : 2WR1 / pdb_00002wr1
Title : structure of influenza H2 hemagglutinin with human receptor
Authors : Liu, J.; Stevens, D.J.; Haire, L.F.; Walker, P.A.; Coombs, P.J.; Russell, R.J.; Gamblin, S.J.; Skehel, J.J.
Deposited on : 2009-08-29
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6234 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 13094 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	4	0	0
			3882	2436	671	753	22			
1	B	490	Total	C	N	O	S	0	0	0
			3879	2434	670	753	22			
1	C	485	Total	C	N	O	S	0	0	0
			3844	2413	664	745	22			

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	3	Total	C	N	O	0	0	0
			46	25	2	19			
2	G	3	Total	C	N	O	0	0	0
			46	25	2	19			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-3)-beta-D-mannopyranose

-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	4	Total	C	N	O	0	0	1
			40	23	2	15			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	H	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	426	Total	O	0	0
			426	426		
7	B	405	Total	O	0	0
			405	405		
7	C	406	Total	O	0	0
			406	406		

MolProbity failed to run properly - this section is therefore empty.

3 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.56Å 141.46Å 199.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 30.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-2.10) 99.8 (30.00-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.10Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.196 , 0.230 0.194 , 0.226	Depositor DCC
R_{free} test set	5783 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	36.5	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13094	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

Of 16 monosaccharides modelled in this entry, 15 were used for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	D	1	2	15,15,15	1.10	1 (6%)	21,21,21	1.97	5 (23%)
2	GAL	D	2	2	11,11,12	0.54	0	15,15,17	1.67	4 (26%)
2	SIA	D	3	2	20,20,21	1.34	1 (5%)	24,28,31	1.29	3 (12%)
3	NAG	E	1	3	14,14,15	0.67	0	17,19,21	1.71	2 (11%)
3	NAG	E	2	3	14,14,15	0.75	0	17,19,21	1.26	3 (17%)
4	NAG	F	1	4	14,14,15	0.68	0	17,19,21	2.00	5 (29%)
4	NAG	F	2	4	14,14,15	0.52	0	17,19,21	0.86	0
4	BMA	F	3	4	11,11,12	0.75	0	15,15,17	0.62	0
2	NAG	G	1	2	15,15,15	1.55	3 (20%)	21,21,21	3.82	15 (71%)
2	GAL	G	2	2	11,11,12	0.77	0	15,15,17	3.10	8 (53%)
2	SIA	G	3	2	20,20,21	1.36	3 (15%)	24,28,31	1.52	5 (20%)
5	NAG	H	1	5	14,14,15	0.55	0	17,19,21	1.62	3 (17%)
5	NAG	H	2	5	14,14,15	0.62	0	17,19,21	0.98	1 (5%)
5	BMA	H	3	5	11,11,12	0.79	0	15,15,17	0.96	1 (6%)
5	MAN	H	4	5	11,11,12	0.82	1 (9%)	15,15,17	1.84	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	2	-	0/6/26/26	0/1/1/1
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1
2	SIA	D	3	2	-	2/18/34/38	0/1/1/1
3	NAG	E	1	3	-	4/6/23/26	0/1/1/1
3	NAG	E	2	3	-	4/6/23/26	0/1/1/1
4	NAG	F	1	4	-	5/6/23/26	0/1/1/1
4	NAG	F	2	4	-	4/6/23/26	0/1/1/1
4	BMA	F	3	4	-	1/2/19/22	0/1/1/1
2	NAG	G	1	2	-	0/6/26/26	0/1/1/1
2	GAL	G	2	2	-	0/2/19/22	0/1/1/1
2	SIA	G	3	2	-	2/18/34/38	0/1/1/1
5	NAG	H	1	5	-	2/6/23/26	0/1/1/1
5	NAG	H	2	5	-	2/6/23/26	0/1/1/1
5	BMA	H	3	5	-	0/2/19/22	0/1/1/1
5	MAN	H	4	5	-	2/2/19/22	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	SIA	C2-C1	4.69	1.56	1.52
2	G	1	NAG	C2-N2	4.20	1.52	1.45
2	G	3	SIA	C4-C5	3.08	1.55	1.53
2	G	3	SIA	O6-C6	-2.80	1.39	1.44
2	G	1	NAG	C8-C7	2.60	1.55	1.50
2	G	3	SIA	C2-C1	2.47	1.54	1.52
2	G	1	NAG	C1-C2	2.24	1.55	1.52
5	H	4	MAN	O2-C2	2.11	1.47	1.43
2	D	1	NAG	C8-C7	2.10	1.54	1.50

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1	NAG	C6-C5-C4	-9.22	91.40	113.00
2	G	2	GAL	C1-O5-C5	-7.10	102.57	112.19
2	G	1	NAG	O5-C1-C2	-6.96	102.52	109.52
2	G	1	NAG	C1-C2-N2	-6.51	103.19	110.73
2	D	1	NAG	O5-C5-C6	5.19	119.33	106.44
2	G	1	NAG	O5-C5-C6	4.80	118.37	106.44
2	G	2	GAL	C2-C3-C4	-4.56	103.00	110.89
2	G	3	SIA	C3-C4-C5	-4.55	105.96	111.46
5	H	1	NAG	C1-O5-C5	-4.48	106.12	112.19
2	G	2	GAL	O6-C6-C5	-4.25	96.72	111.29
3	E	1	NAG	O5-C1-C2	-4.24	104.59	111.29
5	H	4	MAN	O3-C3-C2	-4.11	102.11	109.99
2	G	1	NAG	O1-C1-C2	-4.11	100.68	109.22
2	G	1	NAG	C3-C4-C5	4.08	117.52	110.24
4	F	1	NAG	C4-C3-C2	3.93	116.78	111.02
2	D	1	NAG	C6-C5-C4	-3.91	103.85	113.00
2	G	1	NAG	C3-C2-N2	-3.77	103.51	110.62
2	D	1	NAG	C1-C2-N2	-3.76	106.38	110.73
2	G	2	GAL	O2-C2-C1	-3.71	101.57	109.15
4	F	1	NAG	C3-C4-C5	3.60	116.66	110.24
4	F	1	NAG	C8-C7-N2	3.57	122.14	116.10
3	E	1	NAG	C3-C4-C5	3.50	116.48	110.24
2	G	2	GAL	O5-C5-C6	-3.47	101.76	107.20
5	H	4	MAN	C1-C2-C3	3.47	113.93	109.67
2	G	1	NAG	C4-C3-C2	-3.44	105.30	110.34
2	G	2	GAL	C6-C5-C4	-3.11	105.72	113.00
2	D	1	NAG	O3-C3-C2	-3.07	103.46	109.66
2	G	1	NAG	O3-C3-C4	-3.06	103.28	110.35
4	F	1	NAG	O7-C7-C8	-3.04	116.42	122.06

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1	NAG	O4-C4-C3	3.03	117.35	110.35
2	G	1	NAG	O3-C3-C2	-2.99	103.63	109.66
2	D	3	SIA	O9-C9-C8	-2.97	104.60	111.07
3	E	2	NAG	C2-N2-C7	-2.82	118.88	122.90
3	E	2	NAG	C1-O5-C5	2.82	116.01	112.19
2	D	3	SIA	C3-C4-C5	-2.79	108.08	111.46
4	F	1	NAG	C2-N2-C7	2.70	126.75	122.90
5	H	4	MAN	C3-C4-C5	2.68	115.03	110.24
2	G	2	GAL	O2-C2-C3	-2.67	104.79	110.14
2	D	2	GAL	O3-C3-C4	-2.61	104.31	110.35
2	D	2	GAL	C1-O5-C5	-2.52	108.78	112.19
2	G	3	SIA	C4-C5-N5	-2.49	105.46	110.38
2	D	2	GAL	O3-C3-C2	-2.48	105.24	109.99
2	G	1	NAG	O7-C7-N2	2.46	126.47	121.95
3	E	2	NAG	C4-C3-C2	-2.33	107.60	111.02
5	H	4	MAN	C2-C3-C4	2.32	114.91	110.89
5	H	3	BMA	C1-C2-C3	-2.28	106.86	109.67
5	H	1	NAG	O4-C4-C3	-2.24	105.17	110.35
2	G	1	NAG	O5-C5-C4	-2.23	105.64	109.69
2	D	1	NAG	O5-C1-C2	-2.22	107.28	109.52
5	H	1	NAG	O5-C1-C2	-2.18	107.85	111.29
2	D	3	SIA	O1A-C1-C2	-2.17	117.45	122.57
2	G	3	SIA	O1B-C1-C2	2.16	119.19	113.03
2	G	3	SIA	O7-C7-C8	-2.12	103.70	108.81
2	G	2	GAL	C3-C4-C5	-2.11	106.47	110.24
2	G	3	SIA	O1A-C1-C2	-2.11	117.58	122.57
2	G	1	NAG	C1-C2-C3	-2.10	107.68	110.54
2	G	1	NAG	O7-C7-C8	-2.09	118.17	122.06
2	D	2	GAL	O2-C2-C1	-2.09	104.87	109.15
5	H	2	NAG	C8-C7-N2	2.07	119.61	116.10

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	1	NAG	C4-C5-C6-O6
3	E	1	NAG	O5-C5-C6-O6
4	F	1	NAG	O5-C5-C6-O6
5	H	4	MAN	C4-C5-C6-O6
5	H	4	MAN	O5-C5-C6-O6
2	D	3	SIA	C11-C10-N5-C5
2	D	3	SIA	O10-C10-N5-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	G	3	SIA	C11-C10-N5-C5
2	G	3	SIA	O10-C10-N5-C5
3	E	1	NAG	C8-C7-N2-C2
3	E	1	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2
3	E	2	NAG	O7-C7-N2-C2
4	F	1	NAG	C8-C7-N2-C2
4	F	1	NAG	O7-C7-N2-C2
4	F	2	NAG	C8-C7-N2-C2
4	F	2	NAG	O7-C7-N2-C2
5	H	1	NAG	C8-C7-N2-C2
5	H	1	NAG	O7-C7-N2-C2
5	H	2	NAG	C8-C7-N2-C2
5	H	2	NAG	O7-C7-N2-C2
4	F	1	NAG	C4-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6
4	F	1	NAG	C3-C2-N2-C7
4	F	2	NAG	C4-C5-C6-O6
4	F	2	NAG	O5-C5-C6-O6
3	E	2	NAG	O5-C5-C6-O6
4	F	3	BMA	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	4	MAN	0	2

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.


Oligosaccharide Chain D




Bond lengths



Bond angles






Torsions




Rings

Oligosaccharide Chain G







4.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	A	502	-	14,14,15	0.69	0	17,19,21	0.73	0
6	NAG	C	1500	-	14,14,15	0.65	0	17,19,21	1.10	1 (5%)
6	NAG	A	501	-	14,14,15	0.55	0	17,19,21	0.84	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	502	-	-	5/6/23/26	0/1/1/1
6	NAG	C	1500	-	-	2/6/23/26	0/1/1/1
6	NAG	A	501	-	-	5/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	1500	NAG	C6-C5-C4	-2.35	107.49	113.00
6	A	501	NAG	C1-O5-C5	2.21	115.19	112.19

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	501	NAG	C1-C2-N2-C7
6	A	501	NAG	C8-C7-N2-C2
6	A	501	NAG	O7-C7-N2-C2
6	A	502	NAG	C1-C2-N2-C7
6	A	502	NAG	C8-C7-N2-C2
6	A	502	NAG	O7-C7-N2-C2
6	A	501	NAG	O5-C5-C6-O6
6	A	502	NAG	O5-C5-C6-O6
6	A	502	NAG	C4-C5-C6-O6
6	C	1500	NAG	O5-C5-C6-O6
6	A	501	NAG	C4-C5-C6-O6
6	C	1500	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data ⓘ

5.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	490/509 (96%)	0.00	17 (3%)	47 49	21, 36, 65, 153	1 (0%)
1	B	490/509 (96%)	0.12	26 (5%)	33 35	24, 38, 66, 139	0
1	C	485/509 (95%)	0.08	17 (3%)	47 49	25, 39, 64, 145	0
All	All	1465/1527 (95%)	0.07	60 (4%)	42 44	21, 38, 66, 153	1 (0%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	453	MET	5.7
1	A	74	LEU	5.3
1	B	74	LEU	5.1
1	B	463	GLY	4.9
1	A	459	VAL	4.8
1	A	141	LEU	4.7
1	B	173	GLU	4.5
1	A	498	ALA	4.4
1	B	75	LEU	4.1
1	C	493	GLU	4.0
1	B	114	HIS	4.0
1	C	187	ASP	4.0
1	B	141	LEU	4.0
1	C	459	VAL	3.9
1	C	277	CYS	3.7
1	B	259	ILE	3.5
1	A	75	LEU	3.5
1	B	262	ARG	3.5
1	A	496	LYS	3.4
1	B	492	GLU	3.3
1	C	157	GLY	3.2
1	B	459	VAL	3.2
1	B	390	THR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	326	ILE	3.1
1	C	262	ARG	3.1
1	B	124	ASP	3.1
1	B	142	ASP	3.1
1	A	277	CYS	3.1
1	B	495	SER	3.1
1	B	464	ASN	3.0
1	C	74	LEU	3.0
1	B	76	SER	2.9
1	A	73	ARG	2.9
1	B	113	THR	2.8
1	B	197	VAL	2.8
1	C	141	LEU	2.7
1	C	276	ASN	2.7
1	B	277	CYS	2.6
1	A	324	PRO	2.6
1	A	72	ASP	2.6
1	B	58	ASP	2.6
1	C	348	ASP	2.5
1	A	262	ARG	2.5
1	C	495	SER	2.5
1	A	142	ASP	2.5
1	A	497	ALA	2.5
1	B	73	ARG	2.4
1	A	493	GLU	2.3
1	C	75	LEU	2.3
1	B	494	GLU	2.2
1	C	494	GLU	2.2
1	A	325	GLN	2.2
1	C	18	GLU	2.2
1	A	466	CYS	2.2
1	C	158	SER	2.1
1	B	393	GLU	2.1
1	B	498	ALA	2.1
1	C	462	LEU	2.1
1	C	390	THR	2.1
1	B	350	TRP	2.0

5.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	BMA	F	4	1/12	0.26	0.29	89,89,89,89	0
4	BMA	F	3	11/12	0.63	0.10	88,88,89,89	0
2	NAG	D	1	15/15	0.64	0.13	87,90,92,95	0
4	NAG	F	1	14/15	0.64	0.17	86,88,88,88	0
3	NAG	E	2	14/15	0.67	0.12	75,76,76,76	0
2	NAG	G	1	15/15	0.68	0.17	62,65,66,69	0
4	NAG	F	2	14/15	0.77	0.12	87,87,88,88	0
2	GAL	G	2	11/12	0.78	0.11	47,55,58,58	0
3	NAG	E	1	14/15	0.79	0.15	74,75,75,76	0
5	BMA	H	3	11/12	0.79	0.10	55,56,57,57	0
5	NAG	H	2	14/15	0.80	0.11	55,56,56,56	0
5	MAN	H	4	11/12	0.80	0.14	56,56,56,57	0
2	GAL	D	2	11/12	0.81	0.10	73,80,83,83	0
2	SIA	D	3	20/21	0.86	0.12	60,65,72,73	0
5	NAG	H	1	14/15	0.90	0.11	56,57,57,58	0
2	SIA	G	3	20/21	0.93	0.09	35,40,46,48	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NAG	A	501	14/15	0.75	0.14	92,94,95,95	0
6	NAG	A	502	14/15	0.82	0.13	59,61,62,62	0
6	NAG	C	1500	14/15	0.82	0.14	61,63,63,63	0

5.5 Other polymers [i](#)

There are no such residues in this entry.