



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 11, 2024 – 09:11 AM EST

PDB ID : 3BTP
Title : Crystal structure of Agrobacterium tumefaciens VirE2 in complex with its chaperone VirE1: a novel fold and implications for DNA binding
Authors : Dym, O.; Albeck, S.; Unger, T.; Elbaum, M.; Israel Structural Proteomics Center (ISPC)
Deposited on : 2007-12-30
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	556	 2% 51% 15% 31%
2	B	63	 33% 6% 5% 56%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NH4	B	602	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3466 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Single-strand DNA-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	386	3158	1980	570	595	13	0	1	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	171	LEU	ILE	conflict	UNP P08062

- Molecule 2 is a protein called Protein virE1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	28	227	144	37	44	2	0	0	0

- Molecule 3 is AMMONIUM ION (three-letter code: NH4) (formula: H₄N).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total N 1 1	0	0
3	B	1	Total N 1 1	0	0
3	B	1	Total N 1 1	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	61	Total O 61 61	0	0
5	B	10	Total O 10 10	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Single-strand DNA-binding protein



- Molecule 2: Protein virE1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.02Å 96.27Å 112.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.14 – 2.30 48.14 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.14-2.30) 99.5 (48.14-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.68 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.209 , 0.256 0.208 , 0.253	Depositor DCC
R_{free} test set	1288 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3466	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NH4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.17	9/3225 (0.3%)	1.14	19/4343 (0.4%)
2	B	1.28	0/233	1.07	1/316 (0.3%)
All	All	1.18	9/3458 (0.3%)	1.13	20/4659 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	324	GLU	CG-CD	7.83	1.63	1.51
1	A	324	GLU	CB-CG	6.75	1.65	1.52
1	A	165	ALA	CA-CB	6.73	1.66	1.52
1	A	116	GLU	CG-CD	6.12	1.61	1.51
1	A	258	SER	CB-OG	5.43	1.49	1.42
1	A	324	GLU	CD-OE2	5.31	1.31	1.25
1	A	463	GLU	CD-OE2	5.14	1.31	1.25
1	A	318	LYS	CD-CE	5.09	1.64	1.51
1	A	389	GLU	CG-CD	5.01	1.59	1.51

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	491	ARG	NE-CZ-NH2	-13.66	113.47	120.30
1	A	356	GLN	C-N-CD	-10.78	96.87	120.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	402	ASP	CB-CG-OD1	9.58	126.92	118.30
1	A	356	GLN	N-CA-C	8.81	134.80	111.00
1	A	402	ASP	CB-CG-OD2	-8.58	110.58	118.30
1	A	356	GLN	C-N-CA	8.50	157.71	122.00
1	A	491	ARG	NE-CZ-NH1	8.45	124.53	120.30
1	A	277	ARG	NE-CZ-NH1	-8.01	116.29	120.30
1	A	183	GLU	N-CA-C	7.75	131.92	111.00
1	A	113	ARG	NE-CZ-NH2	-7.30	116.65	120.30
1	A	179	LEU	CA-CB-CG	6.77	130.87	115.30
1	A	243	ARG	NE-CZ-NH1	-6.11	117.25	120.30
2	B	40	ASP	CB-CG-OD2	-5.84	113.05	118.30
1	A	201	ARG	NE-CZ-NH1	5.67	123.13	120.30
1	A	485	ARG	NE-CZ-NH1	5.63	123.11	120.30
1	A	277	ARG	NE-CZ-NH2	5.38	122.99	120.30
1	A	357	PRO	CA-N-CD	-5.37	103.99	111.50
1	A	345	GLN	N-CA-C	5.20	125.04	111.00
1	A	353	MET	CB-CG-SD	5.07	127.61	112.40
1	A	340	ARG	NE-CZ-NH2	-5.05	117.77	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	183	GLU	Peptide
1	A	356	GLN	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3158	0	3089	84	0
2	B	227	0	209	7	0
3	A	1	0	0	1	0
3	B	2	0	0	1	0
4	A	7	0	10	3	0
5	A	61	0	0	2	0
5	B	10	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3466	0	3308	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:GLN:H	1:A:364:THR:HG22	1.09	1.15
1:A:277:ARG:HH11	1:A:277:ARG:CG	1.61	1.12
1:A:325:ARG:HH11	1:A:325:ARG:CG	1.65	1.07
1:A:277:ARG:HG2	1:A:277:ARG:NH1	1.63	1.03
2:B:37:THR:HG22	2:B:40:ASP:H	1.19	1.01
1:A:325:ARG:HG3	1:A:325:ARG:NH1	1.68	1.00
1:A:325:ARG:HH11	1:A:325:ARG:HG3	0.85	1.00
1:A:356:GLN:N	1:A:364:THR:HG22	1.77	0.99
1:A:315:HIS:HD2	1:A:331:ARG:HH12	1.15	0.93
1:A:447:PHE:N	1:A:447:PHE:HD2	1.66	0.91
1:A:277:ARG:HH11	1:A:277:ARG:HG2	0.78	0.91
1:A:447:PHE:N	1:A:447:PHE:CD2	2.28	0.89
1:A:315:HIS:CD2	1:A:331:ARG:HH12	1.92	0.88
1:A:507:LYS:HE3	1:A:508:LEU:O	1.76	0.85
1:A:476:SER:HB2	1:A:502:GLU:OE1	1.77	0.84
1:A:137:ASP:OD1	1:A:138:HIS:HD2	1.63	0.81
1:A:184:PRO:O	1:A:186:GLU:N	2.15	0.80
2:B:37:THR:HG21	5:B:607:HOH:O	1.81	0.78
1:A:350:THR:HG23	1:A:368:HIS:HB3	1.66	0.77
1:A:121:ARG:HH12	1:A:131:THR:HG22	1.49	0.77
3:A:601:NH4:N	3:B:603:NH4:N	2.33	0.76
1:A:413:GLU:HG3	1:A:420:PHE:CD2	2.20	0.75
1:A:413:GLU:HG3	1:A:420:PHE:CE2	2.23	0.73
1:A:178:GLN:O	1:A:178:GLN:HG2	1.89	0.73
1:A:152:TYR:OH	1:A:203:HIS:HE1	1.73	0.71
1:A:179:LEU:C	1:A:182:LEU:HA	2.14	0.67
1:A:178:GLN:O	1:A:179:LEU:HB2	1.93	0.66
1:A:350:THR:CG2	1:A:368:HIS:HB3	2.25	0.66
2:B:31:HIS:CE1	2:B:32:GLN:NE2	2.65	0.65
1:A:325:ARG:CG	1:A:325:ARG:NH1	2.39	0.65
1:A:368:HIS:HD2	5:A:708:HOH:O	1.80	0.64
1:A:171:LEU:CD1	1:A:191:CYS:SG	2.86	0.64
1:A:199:TRP:O	1:A:203:HIS:HD2	1.82	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:LEU:HD13	1:A:191:CYS:SG	2.40	0.62
1:A:450:GLN:O	1:A:491:ARG:NH2	2.34	0.60
1:A:239:GLU:HG2	1:A:240:LYS:HG3	1.83	0.59
1:A:173:GLU:HG3	1:A:174:GLN:NE2	2.16	0.59
1:A:201:ARG:O	1:A:205:GLU:HG3	2.03	0.59
1:A:277:ARG:CD	1:A:332:PHE:O	2.51	0.59
1:A:356:GLN:HB3	1:A:357:PRO:HD2	1.84	0.59
1:A:315:HIS:HD2	1:A:331:ARG:NH1	1.95	0.59
1:A:397:GLN:HE22	1:A:495:TYR:HD2	1.53	0.57
1:A:325:ARG:HD2	1:A:325:ARG:N	2.19	0.57
1:A:413:GLU:OE1	1:A:434:GLY:O	2.23	0.56
1:A:486:ASN:OD1	1:A:490:GLN:HG2	2.05	0.56
1:A:137:ASP:OD1	1:A:138:HIS:CD2	2.53	0.56
1:A:138:HIS:HE1	1:A:144:GLU:OE1	1.89	0.55
1:A:469:PRO:HA	1:A:472:LEU:CD1	2.38	0.54
1:A:482:ILE:HD11	1:A:498:VAL:HG13	1.91	0.53
1:A:296:LYS:HB2	1:A:301:SER:HA	1.90	0.53
1:A:325:ARG:HD2	1:A:325:ARG:H	1.73	0.53
1:A:469:PRO:HA	1:A:472:LEU:HD11	1.89	0.53
1:A:175:LEU:HD23	1:A:179:LEU:HD22	1.91	0.53
1:A:350:THR:HG22	5:A:708:HOH:O	2.09	0.53
1:A:367:ARG:HE	1:A:460:ASN:ND2	2.07	0.52
1:A:277:ARG:HD3	1:A:332:PHE:O	2.10	0.51
2:B:37:THR:CG2	2:B:40:ASP:H	2.07	0.51
2:B:37:THR:HG22	2:B:40:ASP:N	2.05	0.51
1:A:114:THR:HB	1:A:134:CYS:HB3	1.92	0.51
1:A:171:LEU:HD11	1:A:191:CYS:SG	2.51	0.50
1:A:155:GLY:CA	4:A:701:PEG:H21	2.42	0.50
1:A:326:GLY:O	1:A:330:ILE:HG13	2.12	0.50
1:A:329:ASP:OD2	1:A:340:ARG:HD2	2.12	0.50
1:A:486:ASN:CG	1:A:490:GLN:HG2	2.32	0.49
1:A:173:GLU:HG3	1:A:174:GLN:HE21	1.77	0.49
1:A:292:GLU:O	1:A:296:LYS:HG2	2.12	0.49
1:A:121:ARG:NH1	1:A:131:THR:HG22	2.24	0.49
1:A:152:TYR:OH	1:A:203:HIS:CE1	2.61	0.49
1:A:486:ASN:ND2	1:A:490:GLN:HG2	2.27	0.49
1:A:182:LEU:C	1:A:183:GLU:HG3	2.34	0.48
1:A:397:GLN:NE2	1:A:495:TYR:HD2	2.12	0.48
1:A:178:GLN:O	1:A:178:GLN:CG	2.56	0.48
1:A:155:GLY:HA3	4:A:701:PEG:H21	1.97	0.47
4:A:701:PEG:H41	2:B:51:LEU:HD21	1.95	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:ASN:HD21	1:A:212:ASP:H	1.62	0.46
1:A:477:GLN:HG3	1:A:479:ASP:H	1.80	0.46
1:A:324:GLU:HB3	1:A:327:SER:OG	2.15	0.46
1:A:469:PRO:O	1:A:472:LEU:HD12	2.15	0.46
1:A:296:LYS:HE2	1:A:296:LYS:HB3	1.74	0.46
2:B:46:LEU:O	2:B:50:VAL:HB	2.15	0.45
1:A:277:ARG:CG	1:A:277:ARG:NH1	2.38	0.45
1:A:476:SER:O	1:A:476:SER:OG	2.34	0.44
1:A:179:LEU:O	1:A:182:LEU:HD12	2.18	0.44
1:A:486:ASN:HD21	1:A:490:GLN:CG	2.30	0.44
1:A:511:ASP:OD1	1:A:512:ALA:N	2.51	0.43
1:A:186:GLU:O	1:A:190:GLN:HG3	2.18	0.43
1:A:338:GLU:OE1	1:A:340:ARG:NH2	2.50	0.43
1:A:486:ASN:ND2	1:A:490:GLN:CG	2.82	0.42
1:A:504:LEU:HB2	1:A:506[B]:MET:HG3	2.01	0.42
1:A:277:ARG:HD3	1:A:335:PHE:HB2	1.99	0.42
1:A:225:ARG:H	1:A:225:ARG:HG3	1.60	0.41
1:A:277:ARG:NH1	1:A:281:ASP:OD2	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/556 (67%)	361 (96%)	9 (2%)	5 (1%)	12	12
2	B	26/63 (41%)	26 (100%)	0	0	100	100
All	All	401/619 (65%)	387 (96%)	9 (2%)	5 (1%)	12	14

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	185	ARG
1	A	356	GLN
1	A	183	GLU
1	A	184	PRO
1	A	173	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/483 (70%)	312 (93%)	25 (7%)	13	17
2	B	27/58 (47%)	25 (93%)	2 (7%)	13	17
All	All	364/541 (67%)	337 (93%)	27 (7%)	13	17

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	131	THR
1	A	142	LYS
1	A	178	GLN
1	A	182	LEU
1	A	190	GLN
1	A	197	GLU
1	A	222	LYS
1	A	232	GLU
1	A	240	LYS
1	A	275	SER
1	A	277	ARG
1	A	307	LEU
1	A	314	MET
1	A	325	ARG
1	A	350	THR
1	A	356	GLN
1	A	376	GLN
1	A	427	VAL
1	A	447	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	448	ASP
1	A	461	LEU
1	A	470	GLU
1	A	472	LEU
1	A	498	VAL
1	A	508	LEU
2	B	37	THR
2	B	50	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	HIS
1	A	146	ASN
1	A	174	GLN
1	A	190	GLN
1	A	200	ASN
1	A	203	HIS
1	A	270	ASN
1	A	315	HIS
1	A	368	HIS
1	A	377	ASN
1	A	404	ASN
1	A	460	ASN
2	B	31	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 3 are modelled with single atom - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PEG	A	701	-	6,6,6	0.68	0	5,5,5	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	701	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	701	PEG	C4-C3-O2-C2
4	A	701	PEG	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	701	PEG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/556 (69%)	0.04	10 (2%) 56 63	15, 30, 55, 68	0
2	B	28/63 (44%)	-0.05	0 100 100	17, 24, 46, 51	0
All	All	414/619 (66%)	0.04	10 (2%) 59 66	15, 30, 55, 68	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	476	SER	4.5
1	A	449	ARG	3.3
1	A	226	ASN	2.5
1	A	178	GLN	2.5
1	A	173	GLU	2.4
1	A	517	ALA	2.4
1	A	341	ALA	2.3
1	A	185	ARG	2.1
1	A	447	PHE	2.1
1	A	469	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NH4	B	602	1/1	0.78	0.43	33,33,33,33	0
3	NH4	B	603	1/1	0.88	0.18	16,16,16,16	0
3	NH4	A	601	1/1	0.92	0.24	27,27,27,27	0
4	PEG	A	701	7/7	0.92	0.19	30,34,36,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.