



# Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 11:00 pm BST

PDB ID : 3CXN  
Title : Structure of the Urease Accessory Protein UreF from Helicobacter pylori  
Authors : Lam, R.; Johns, K.; Romanov, V.; Dong, A.; Wu-Brown, J.; Guthrie, J.; Dharamsi, A.; Thambipillai, D.; Mansoury, K.; Edwards, A.M.; Pai, E.F.; Chirgadze, N.Y.  
Deposited on : 2008-04-24  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	
1	B	274	
1	C	274	

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 5446 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urease accessory protein ureF.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	207	Total 1688	C 1084	N 272	O 322	S 2	Se 8	0	9	0
1	B	209	Total 1733	C 1115	N 278	O 330	S 2	Se 8	0	14	0
1	C	204	Total 1618	C 1031	N 265	O 313	S 2	Se 7	0	2	0

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MSE	-	EXPRESSION TAG	UNP Q09065
A	-18	GLY	-	EXPRESSION TAG	UNP Q09065
A	-17	SER	-	EXPRESSION TAG	UNP Q09065
A	-16	SER	-	EXPRESSION TAG	UNP Q09065
A	-15	HIS	-	EXPRESSION TAG	UNP Q09065
A	-14	HIS	-	EXPRESSION TAG	UNP Q09065
A	-13	HIS	-	EXPRESSION TAG	UNP Q09065
A	-12	HIS	-	EXPRESSION TAG	UNP Q09065
A	-11	HIS	-	EXPRESSION TAG	UNP Q09065
A	-10	HIS	-	EXPRESSION TAG	UNP Q09065
A	-9	SER	-	EXPRESSION TAG	UNP Q09065
A	-8	SER	-	EXPRESSION TAG	UNP Q09065
A	-7	GLY	-	EXPRESSION TAG	UNP Q09065
A	-6	LEU	-	EXPRESSION TAG	UNP Q09065
A	-5	VAL	-	EXPRESSION TAG	UNP Q09065
A	-4	PRO	-	EXPRESSION TAG	UNP Q09065
A	-3	ARG	-	EXPRESSION TAG	UNP Q09065
A	-2	GLY	-	EXPRESSION TAG	UNP Q09065
A	-1	SER	-	EXPRESSION TAG	UNP Q09065
A	0	HIS	-	EXPRESSION TAG	UNP Q09065
A	1	MSE	-	EXPRESSION TAG	UNP Q09065
B	-19	MSE	-	EXPRESSION TAG	UNP Q09065
B	-18	GLY	-	EXPRESSION TAG	UNP Q09065

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	SER	-	EXPRESSION TAG	UNP Q09065
B	-16	SER	-	EXPRESSION TAG	UNP Q09065
B	-15	HIS	-	EXPRESSION TAG	UNP Q09065
B	-14	HIS	-	EXPRESSION TAG	UNP Q09065
B	-13	HIS	-	EXPRESSION TAG	UNP Q09065
B	-12	HIS	-	EXPRESSION TAG	UNP Q09065
B	-11	HIS	-	EXPRESSION TAG	UNP Q09065
B	-10	HIS	-	EXPRESSION TAG	UNP Q09065
B	-9	SER	-	EXPRESSION TAG	UNP Q09065
B	-8	SER	-	EXPRESSION TAG	UNP Q09065
B	-7	GLY	-	EXPRESSION TAG	UNP Q09065
B	-6	LEU	-	EXPRESSION TAG	UNP Q09065
B	-5	VAL	-	EXPRESSION TAG	UNP Q09065
B	-4	PRO	-	EXPRESSION TAG	UNP Q09065
B	-3	ARG	-	EXPRESSION TAG	UNP Q09065
B	-2	GLY	-	EXPRESSION TAG	UNP Q09065
B	-1	SER	-	EXPRESSION TAG	UNP Q09065
B	0	HIS	-	EXPRESSION TAG	UNP Q09065
B	1	MSE	-	EXPRESSION TAG	UNP Q09065
C	-19	MSE	-	EXPRESSION TAG	UNP Q09065
C	-18	GLY	-	EXPRESSION TAG	UNP Q09065
C	-17	SER	-	EXPRESSION TAG	UNP Q09065
C	-16	SER	-	EXPRESSION TAG	UNP Q09065
C	-15	HIS	-	EXPRESSION TAG	UNP Q09065
C	-14	HIS	-	EXPRESSION TAG	UNP Q09065
C	-13	HIS	-	EXPRESSION TAG	UNP Q09065
C	-12	HIS	-	EXPRESSION TAG	UNP Q09065
C	-11	HIS	-	EXPRESSION TAG	UNP Q09065
C	-10	HIS	-	EXPRESSION TAG	UNP Q09065
C	-9	SER	-	EXPRESSION TAG	UNP Q09065
C	-8	SER	-	EXPRESSION TAG	UNP Q09065
C	-7	GLY	-	EXPRESSION TAG	UNP Q09065
C	-6	LEU	-	EXPRESSION TAG	UNP Q09065
C	-5	VAL	-	EXPRESSION TAG	UNP Q09065
C	-4	PRO	-	EXPRESSION TAG	UNP Q09065
C	-3	ARG	-	EXPRESSION TAG	UNP Q09065
C	-2	GLY	-	EXPRESSION TAG	UNP Q09065
C	-1	SER	-	EXPRESSION TAG	UNP Q09065
C	0	HIS	-	EXPRESSION TAG	UNP Q09065
C	1	MSE	-	EXPRESSION TAG	UNP Q09065

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	118	Total	O	0	0
			118	118		
3	B	185	Total	O	0	0
			185	185		
3	C	98	Total	O	0	0
			98	98		



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.22Å 89.42Å 66.04Å 90.00° 94.03° 90.00°	Depositor
Resolution (Å)	26.77 – 1.55 26.77 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.3 (26.77-1.55) 99.3 (26.77-1.55)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.01 (at 1.55Å)	Xtrriage
Refinement program	REFMAC refmac_5.2.0019	Depositor
R, $R_{free}$	0.187 , 0.211 0.192 , 0.213	Depositor DCC
$R_{free}$ test set	5637 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtrriage
Anisotropy	0.109	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 50.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5446	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/1736	0.60	0/2330
1	B	0.54	0/1796	0.64	0/2408
1	C	0.45	0/1641	0.57	0/2203
All	All	0.50	0/5173	0.60	0/6941

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1688	0	1734	15	0
1	B	1733	0	1795	13	0
1	C	1618	0	1648	23	0
2	B	6	0	8	0	0
3	A	118	0	0	1	0
3	B	185	0	0	3	0
3	C	98	0	0	2	0
All	All	5446	0	5185	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.



All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:VAL:HG11	1:C:191:ILE:HD11	1.47	0.95
1:C:123:ALA:O	1:C:126:LYS:HE2	1.70	0.92
1:B:85[B]:GLU:HG2	1:B:181[B]:TYR:OH	1.71	0.90
1:B:85[B]:GLU:OE2	1:B:181[B]:TYR:OH	1.92	0.86
1:C:73:LEU:HD22	1:C:189:MSE:HE3	1.70	0.73
1:B:85[B]:GLU:CG	1:B:181[B]:TYR:OH	2.37	0.71
1:A:42:VAL:HG11	1:C:191:ILE:CD1	2.18	0.70
1:C:126:LYS:HE3	3:C:306:HOH:O	1.91	0.70
1:B:76:ASN:ND2	3:B:405:HOH:O	2.27	0.67
1:C:32:GLU:OE1	1:C:176:LYS:NZ	2.29	0.65
1:C:186:THR:HG21	1:C:215:PHE:CZ	2.33	0.64
1:C:82:LEU:HB2	1:C:218:LEU:HD11	1.83	0.61
1:B:85[B]:GLU:CD	1:B:181[B]:TYR:OH	2.39	0.60
1:B:112:MSE:HG2	1:B:121:ARG:HD2	1.84	0.60
1:A:32:GLU:OE2	1:A:176:LYS:NZ	2.35	0.59
1:C:173:GLU:OE2	1:C:175:LYS:HB3	2.03	0.58
1:A:54:LEU:HD13	1:A:72:TYR:CE2	2.41	0.55
1:A:126:LYS:O	1:A:129:ASN:ND2	2.40	0.54
1:B:218[A]:LEU:HG	3:B:333:HOH:O	2.08	0.53
1:C:179:ARG:HG2	1:C:219:ILE:HD13	1.92	0.51
1:A:51:SER:HA	1:A:192:ASN:ND2	2.26	0.51
1:C:73:LEU:CD2	1:C:189:MSE:HE3	2.41	0.50
1:A:213[B]:SER:HB3	1:A:214:PRO:HD3	1.94	0.50
1:A:54:LEU:HG	1:A:58:ILE:CD1	2.41	0.49
1:A:42:VAL:CG1	1:C:191:ILE:HD11	2.30	0.49
1:C:221:LYS:HE3	1:C:225:LEU:HD21	1.92	0.49
1:B:60:GLN:C	1:B:61:LYS:HG2	2.34	0.47
1:C:186:THR:HG21	1:C:215:PHE:CE1	2.49	0.47
1:A:179[B]:ARG:HG3	1:C:33:PHE:CE1	2.50	0.47
1:A:54:LEU:HD23	1:A:192:ASN:ND2	2.30	0.47
1:C:186:THR:HG21	1:C:215:PHE:HZ	1.75	0.46
1:B:103:LYS:HE3	3:B:380:HOH:O	2.16	0.44
1:A:90[A]:LYS:HG3	1:A:174:LEU:HD21	2.00	0.44
1:C:123:ALA:O	1:C:126:LYS:HG3	2.18	0.43
1:B:82:LEU:HB2	1:B:218[B]:LEU:HD11	2.00	0.43
1:A:153:LYS:HB3	1:A:153:LYS:HE3	1.88	0.42
1:C:118:MSE:CE	1:C:119:GLU:HG3	2.49	0.42
1:C:118:MSE:HE3	1:C:119:GLU:N	2.34	0.42
1:C:28:HIS:H	1:C:28:HIS:CD2	2.38	0.42
1:B:50:HIS:HE1	1:B:85[A]:GLU:OE1	2.03	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:LEU:HG	3:A:350:HOH:O	2.20	0.41
1:C:186:THR:HG23	1:C:189:MSE:HE2	2.02	0.41
1:C:213:SER:HB3	1:C:214:PRO:HD3	2.03	0.41
1:C:67:GLU:HG3	3:C:305:HOH:O	2.19	0.41
1:A:57:TYR:CD2	1:B:118[A]:MSE:HE1	2.55	0.41
1:B:100:ASP:OD2	1:B:103:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/274 (78%)	211 (99%)	3 (1%)	0	100	100
1	B	221/274 (81%)	218 (99%)	3 (1%)	0	100	100
1	C	202/274 (74%)	196 (97%)	6 (3%)	0	100	100
All	All	637/822 (78%)	625 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/234 (82%)	190 (98%)	3 (2%)	62	35

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	200/234 (86%)	197 (98%)	3 (2%)	65	37
1	C	183/234 (78%)	176 (96%)	7 (4%)	33	6
All	All	576/702 (82%)	563 (98%)	13 (2%)	52	21

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	MSE
1	A	129	ASN
1	A	166	PHE
1	B	118[A]	MSE
1	B	118[B]	MSE
1	B	166	PHE
1	C	72	TYR
1	C	76	ASN
1	C	80	GLN
1	C	118	MSE
1	C	121	ARG
1	C	126	LYS
1	C	166	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	ASN
1	A	188	ASN
1	A	192	ASN
1	B	76	ASN
1	C	25	ASN
1	C	28	HIS
1	C	80	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	B	255	-	5,5,5	0.35	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	255	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	255	GOL	C1-C2-C3-O3
2	B	255	GOL	O1-C1-C2-C3
2	B	255	GOL	O2-C2-C3-O3
2	B	255	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	201/274 (73%)	0.98	27 (13%) 3 2	17, 24, 50, 57	1 (0%)
1	B	203/274 (74%)	0.52	12 (5%) 22 26	15, 20, 35, 47	1 (0%)
1	C	198/274 (72%)	0.92	33 (16%) 1 1	20, 29, 43, 46	0
All	All	602/822 (73%)	0.81	72 (11%) 4 3	15, 25, 44, 57	2 (0%)

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	52	PHE	15.2
1	A	120	LEU	7.3
1	B	59	GLN	7.0
1	A	53	GLY	6.6
1	A	27	ALA	6.4
1	A	48	TYR	6.1
1	C	25	ASN	6.0
1	B	52	PHE	6.0
1	A	122	LEU	5.9
1	A	47	SER	5.8
1	B	25	ASN	5.4
1	A	119	GLU	5.3
1	A	121	ARG	5.2
1	C	47	SER	5.2
1	C	120	LEU	5.1
1	A	28	HIS	5.0
1	B	56	THR	5.0
1	C	233	ALA	4.9
1	A	51	SER	4.9
1	A	54	LEU	4.6
1	A	123	ALA	4.3
1	A	55	GLU	3.9
1	C	224	GLU	3.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	60	GLN	3.9
1	A	29	VAL	3.8
1	C	122	LEU	3.8
1	C	46	GLY	3.8
1	A	117	PRO	3.7
1	A	49	THR	3.7
1	C	77	LEU	3.6
1	C	119	GLU	3.6
1	C	232	THR	3.6
1	C	83	TYR	3.6
1	A	59	GLN	3.5
1	C	116	SER	3.5
1	A	58	ILE	3.5
1	A	125	GLN	3.5
1	B	55	GLU	3.4
1	B	58	ILE	3.3
1	B	53	GLY	3.3
1	C	78	SER	3.3
1	A	127	LEU	3.2
1	C	53	GLY	3.2
1	C	225	LEU	3.1
1	A	50	HIS	3.0
1	C	121	ARG	3.0
1	A	126	LYS	2.9
1	C	76	ASN	2.9
1	B	51	SER	2.7
1	C	55	GLU	2.7
1	B	47	SER	2.7
1	C	226	ASP	2.6
1	C	222	THR	2.6
1	C	231	CYS	2.6
1	C	229	HIS	2.6
1	B	233	ALA	2.5
1	C	195	LYS	2.4
1	C	28	HIS	2.3
1	C	98	GLN	2.2
1	C	57	TYR	2.2
1	C	198	PRO	2.2
1	C	113	LEU	2.2
1	C	27	ALA	2.2
1	C	135	LEU	2.1
1	C	80	GLN	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	148	ASN	2.1
1	A	139	ASN	2.1
1	A	56	THR	2.0
1	A	140	GLU	2.0
1	A	181[A]	TYR	2.0
1	B	181[A]	TYR	2.0
1	C	45	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	B	255	6/6	0.59	0.25	53,53,54,54	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.