



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 23, 2023 – 03:00 PM EDT

PDB ID : 3DLJ  
Title : Crystal structure of human carnosine dipeptidase 1  
Authors : Dong, A.; Dobrovetsky, E.; Seitova, A.; He, H.; Tempel, W.; Kozieradzki, I.; Arrowsmith, C.H.; Weigelt, J.; Bountra, C.; Edwards, A.M.; Bochkarev, A.; Cossar, D.; Structural Genomics Consortium (SGC)  
Deposited on : 2008-06-27  
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	
1	B	485	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UNX	A	2004	-	-	-	X
3	UNX	A	2005	-	-	-	X
3	UNX	A	2006	-	-	-	X
3	UNX	A	2007	-	-	-	X
3	UNX	A	2008	-	-	-	X
3	UNX	A	2009	-	-	-	X
3	UNX	A	2010	-	-	-	X
3	UNX	A	2011	-	-	-	X
3	UNX	A	2012	-	-	-	X
3	UNX	B	2003	-	-	-	X
3	UNX	B	2004	-	-	-	X
3	UNX	B	2005	-	-	-	X
3	UNX	B	2006	-	-	-	X
3	UNX	B	2007	-	-	-	X
3	UNX	B	2008	-	-	-	X

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-Ala-His dipeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	471	3600	2322	591	672	15	0	0	0
1	B	467	3583	2300	598	670	15	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q96KN2
A	-2	ALA	-	expression tag	UNP Q96KN2
A	-1	MET	-	expression tag	UNP Q96KN2
A	0	ASP	-	expression tag	UNP Q96KN2
B	-3	GLY	-	expression tag	UNP Q96KN2
B	-2	ALA	-	expression tag	UNP Q96KN2
B	-1	MET	-	expression tag	UNP Q96KN2
B	0	ASP	-	expression tag	UNP Q96KN2


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	9	Total	X	0	0
			9	9		
3	B	6	Total	X	0	0
			6	6		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O S	0	0
			5	4 1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	109	Total	O	0	0
			109	109		
5	B	104	Total	O	0	0
			104	104		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-Ala-His dipeptidase



- Molecule 1: Beta-Ala-His dipeptidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.11Å 75.61Å 103.72Å 90.00° 109.79° 90.00°	Depositor
Resolution (Å)	29.91 – 2.26 29.90 – 2.26	Depositor EDS
% Data completeness (in resolution range)	97.3 (29.91-2.26) 97.3 (29.90-2.26)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 2.26Å)	Xtrriage
Refinement program	REFMAC 5.3.0037	Depositor
R, $R_{free}$	0.200 , 0.254 0.236 , 0.282	Depositor DCC
$R_{free}$ test set	1051 reflections (2.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.5	Xtrriage
Anisotropy	0.136	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 56.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.019 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7420	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN, UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.61	0/3683	0.67	2/5010 (0.0%)
1	B	0.64	1/3663 (0.0%)	0.67	1/4984 (0.0%)
All	All	0.62	1/7346 (0.0%)	0.67	3/9994 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	480	LEU	C-O	15.59	1.52	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	ARG	NE-CZ-NH2	-6.33	117.14	120.30
1	A	299	ARG	NE-CZ-NH2	-5.03	117.78	120.30
1	B	441	LEU	CA-CB-CG	5.01	126.83	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3600	0	3474	11	0
1	B	3583	0	3440	30	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	9	0	0	0	0
3	B	6	0	0	0	0
4	A	5	0	0	0	0
5	A	109	0	0	1	0
5	B	104	0	0	0	0
All	All	7420	0	6914	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302:LYS:HG2	1:B:378:SER:HB3	1.72	0.71
1:B:48:LEU:HG	1:B:85:PRO:HG3	1.76	0.68
1:B:200:ILE:HB	1:B:441:LEU:HD22	1.74	0.68
1:B:35:SER:HB2	1:B:48:LEU:HD13	1.78	0.65
1:A:10:GLU:O	1:A:14:GLN:HG3	2.00	0.62
1:B:307:THR:HG23	1:B:310:GLU:H	1.71	0.56
1:B:411:VAL:HG21	1:B:469:LEU:HA	1.91	0.52
1:B:289:LEU:HD11	1:B:294:TYR:HB2	1.91	0.52
1:A:33:ILE:HG12	1:A:50:ARG:NH1	2.26	0.51
1:B:84:ILE:HG23	1:B:172:MET:HG3	1.93	0.50
1:A:226:GLU:OE2	1:A:342:ARG:CZ	2.61	0.47
1:B:91:GLU:HB3	1:B:166:LYS:HG2	1.95	0.47
1:B:408:ILE:HG13	1:B:469:LEU:HD11	1.97	0.47
1:B:104:TYR:HB3	1:B:200:ILE:HG12	1.97	0.46
1:B:138:THR:HG21	1:B:449:ASP:HB3	1.97	0.46
1:A:291:LEU:HD11	1:A:311:ILE:HG13	1.98	0.46
1:A:84:ILE:HG23	1:A:172:MET:HG3	1.97	0.45
1:B:234:PHE:HB2	1:B:339:ILE:HB	1.99	0.45
1:B:31:VAL:HG21	1:B:125:LEU:HD13	1.98	0.44
1:B:104:TYR:OH	1:B:170:GLU:OE2	2.24	0.44
1:B:353:PRO:HA	1:B:354:HIS:HA	1.74	0.44
1:A:161:LEU:HA	1:A:162:PRO:HD3	1.88	0.44
1:A:302:LYS:CG	1:B:378:SER:HB3	2.45	0.43
1:B:427:ILE:HG13	1:B:430:MET:HE2	2.01	0.42
1:B:212:ALA:HB1	1:B:419:ILE:HG12	2.01	0.42
1:A:356:ASN:ND2	5:A:2097:HOH:O	2.52	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:400:GLN:HG3	1:B:438:SER:OG	2.19	0.42
1:B:62:LEU:HD21	1:B:152:SER:HB3	2.01	0.42
1:B:118:TRP:HB3	1:B:120:THR:O	2.19	0.42
1:B:431:PHE:CE2	1:B:437:LYS:HB2	2.55	0.42
1:B:389:LEU:HD22	1:B:426:PRO:HB3	2.02	0.42
1:A:228:LYS:HD2	1:A:342:ARG:HD2	2.02	0.41
1:B:19:HIS:O	1:B:22:GLU:HG2	2.20	0.41
1:B:15:TYR:CE1	1:B:19:HIS:CE1	3.08	0.41
1:B:180:LEU:HG	1:B:430:MET:HE1	2.02	0.41
1:B:48:LEU:O	1:B:52:MET:HG2	2.20	0.41
1:B:34:GLU:HG2	1:B:123:TYR:CG	2.56	0.41
1:B:214:THR:HB	1:B:441:LEU:HB2	2.02	0.40
1:A:228:LYS:HB2	1:A:342:ARG:HG3	2.02	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/485 (96%)	452 (98%)	11 (2%)	0	100	100
1	B	459/485 (95%)	448 (98%)	11 (2%)	0	100	100
All	All	922/970 (95%)	900 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/418 (89%)	367 (99%)	4 (1%)	73	82
1	B	370/418 (88%)	366 (99%)	4 (1%)	73	82
All	All	741/836 (89%)	733 (99%)	8 (1%)	73	82

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	THR
1	A	278	GLU
1	A	279	GLU
1	A	317	ARG
1	B	113	ASP
1	B	114	ARG
1	B	295	ARG
1	B	317	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	19	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 4 are monoatomic and 15 are unknown - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	A	2003	-	4,4,4	0.18	0	6,6,6	0.87	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	471/485 (97%)	0.40	35 (7%) <b>14</b> <b>15</b>	28, 36, 50, 58	0
1	B	467/485 (96%)	0.61	44 (9%) <b>8</b> <b>9</b>	26, 36, 45, 56	0
All	All	938/970 (96%)	0.51	79 (8%) <b>11</b> <b>12</b>	26, 36, 48, 58	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	PRO	5.2
1	B	189	ASP	4.6
1	B	96	PRO	4.3
1	A	434	ILE	4.0
1	B	436	HIS	4.0
1	A	396	ILE	3.8
1	A	436	HIS	3.8
1	A	76	LEU	3.8
1	A	395	ASN	3.6
1	B	477	MET	3.6
1	A	189	ASP	3.5
1	B	478	ALA	3.4
1	A	8	LEU	3.2
1	B	119	LEU	3.2
1	A	79	GLY	3.2
1	B	323	ILE	3.2
1	B	143	PRO	3.2
1	A	143	PRO	3.1
1	B	296	ASN	3.1
1	A	201	SER	3.1
1	A	288	HIS	3.0
1	A	81	SER	2.9
1	A	80	GLN	2.9
1	A	394	ALA	2.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	204	LEU	2.8
1	B	286	ALA	2.8
1	A	5	PRO	2.8
1	B	106	HIS	2.8
1	B	99	GLY	2.7
1	A	480	LEU	2.7
1	B	97	THR	2.7
1	A	402	LEU	2.7
1	B	422	GLY	2.7
1	A	397	ASP	2.7
1	B	107	LEU	2.6
1	A	146	ALA	2.6
1	A	251	VAL	2.6
1	B	288	HIS	2.6
1	A	103	PHE	2.5
1	B	128	VAL	2.4
1	B	185	GLU	2.4
1	B	98	LYS	2.4
1	B	193	SER	2.4
1	A	144	VAL	2.4
1	B	395	ASN	2.4
1	B	407	ALA	2.3
1	B	129	ASP	2.3
1	B	406	ARG	2.3
1	B	397	ASP	2.3
1	B	461	TRP	2.3
1	B	434	ILE	2.3
1	A	250	LEU	2.3
1	A	7	ALA	2.2
1	B	200	ILE	2.2
1	A	399	THR	2.2
1	A	82	LEU	2.2
1	B	475	LEU	2.2
1	B	140	ASN	2.2
1	B	291	LEU	2.2
1	B	100	THR	2.2
1	A	433	GLU	2.2
1	B	120	THR	2.2
1	A	203	ASN	2.1
1	B	435	VAL	2.1
1	A	72	GLY	2.1
1	B	15	TYR	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	200	ILE	2.1
1	A	194	GLY	2.1
1	B	452	HIS	2.1
1	B	410	THR	2.1
1	B	105	GLY	2.1
1	A	142	GLY	2.1
1	B	402	LEU	2.1
1	B	93	GLY	2.1
1	B	414	THR	2.1
1	B	394	ALA	2.1
1	B	474	PHE	2.1
1	A	75	GLN	2.0
1	B	196	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	UNX	A	2010	1/1	-0.68	1.45	2,2,2,2	1
3	UNX	A	2008	1/1	-0.44	2.47	2,2,2,2	1
3	UNX	A	2006	1/1	-0.43	1.60	2,2,2,2	1
3	UNX	B	2008	1/1	-0.35	3.61	2,2,2,2	1
3	UNX	A	2009	1/1	-0.26	4.31	2,2,2,2	1
3	UNX	B	2007	1/1	-0.14	1.26	2,2,2,2	1
3	UNX	B	2004	1/1	-0.14	2.71	2,2,2,2	1
3	UNX	A	2011	1/1	-0.06	3.28	2,2,2,2	1
3	UNX	B	2006	1/1	0.04	3.27	2,2,2,2	1
3	UNX	A	2005	1/1	0.06	2.83	2,2,2,2	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	UNX	A	2012	1/1	0.07	2.85	2,2,2,2	1
3	UNX	A	2007	1/1	0.30	2.85	2,2,2,2	1
3	UNX	A	2004	1/1	0.41	1.84	2,2,2,2	1
3	UNX	B	2003	1/1	0.50	4.00	2,2,2,2	1
3	UNX	B	2005	1/1	0.52	3.36	2,2,2,2	1
2	ZN	B	2001	1/1	0.97	0.11	26,26,26,26	0
4	SO4	A	2003	5/5	0.97	0.13	34,36,38,40	0
2	ZN	A	2001	1/1	0.98	0.09	34,34,34,34	0
2	ZN	B	2002	1/1	0.98	0.14	29,29,29,29	0
2	ZN	A	2002	1/1	0.99	0.10	30,30,30,30	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.