



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 02:19 pm BST

PDB ID : 3E9D
Title : Structure of full-length TIGAR from *Danio rerio*
Authors : Li, H.; Jogl, G.
Deposited on : 2008-08-21
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	 8% 88% 6% 6%
1	B	265	 8% 86% 9%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4546 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Zgc:56074.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	1945	1218	358	358	11	0	0	0
1	B	240	1881	1180	348	342	11	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	258	LEU	-	EXPRESSION TAG	UNP Q7ZVE3
A	259	GLU	-	EXPRESSION TAG	UNP Q7ZVE3
A	260	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
A	261	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
A	262	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
A	263	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
A	264	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
A	265	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
B	258	LEU	-	EXPRESSION TAG	UNP Q7ZVE3
B	259	GLU	-	EXPRESSION TAG	UNP Q7ZVE3
B	260	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
B	261	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
B	262	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
B	263	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
B	264	HIS	-	EXPRESSION TAG	UNP Q7ZVE3
B	265	HIS	-	EXPRESSION TAG	UNP Q7ZVE3

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total K 1 1	0	0
3	A	1	Total K 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	365	Total O 365 365	0	0
4	B	333	Total O 333 333	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	127.66Å 115.54Å 61.31Å 90.00° 104.66° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 29.63 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.9 (30.00-2.00) 96.9 (29.63-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.5.0044	Depositor
R, R_{free}	0.171 , 0.213 0.172 , 0.215	Depositor DCC
R_{free} test set	2835 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtrriage
Anisotropy	0.095	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4546	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/1982	0.61	2/2677 (0.1%)
1	B	0.56	0/1916	0.57	0/2584
All	All	0.56	0/3898	0.59	2/5261 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	222	ARG	NE-CZ-NH1	-8.26	116.17	120.30
1	A	222	ARG	NE-CZ-NH2	5.53	123.06	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1945	0	1944	20	0
1	B	1881	0	1882	10	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	365	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	333	0	0	1	0
All	All	4546	0	3826	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:ARG:HG3	1:A:222:ARG:HH11	1.08	1.10
1:A:157:GLU:HG2	1:B:237:THR:CG2	1.88	1.03
1:A:222:ARG:CG	1:A:222:ARG:HH11	1.78	0.94
1:A:7:THR:OG1	1:A:222:ARG:NH1	2.02	0.93
1:A:23:GLN:OE1	1:A:29:THR:OG1	1.92	0.88
1:A:157:GLU:HG2	1:B:237:THR:HG22	1.54	0.88
1:A:157:GLU:HA	1:B:238:ARG:HH21	1.37	0.87
1:A:222:ARG:NH1	1:A:222:ARG:HG3	1.93	0.78
1:A:134:MET:CE	4:A:613:HOH:O	2.30	0.78
1:B:238:ARG:HD3	4:B:476:HOH:O	1.92	0.69
1:A:134:MET:HE1	4:A:613:HOH:O	1.92	0.62
1:A:222:ARG:CG	1:A:222:ARG:NH1	2.51	0.61
1:A:22:LEU:HD11	1:A:103:LEU:HD22	1.83	0.60
1:A:157:GLU:HG2	1:B:237:THR:HG21	1.80	0.59
1:B:229:ARG:HE	1:B:232:SER:HA	1.69	0.58
1:A:27:ILE:HD12	1:A:96:GLU:HG2	1.86	0.56
1:A:70:LEU:HD12	1:A:81:MET:HE2	1.88	0.55
1:A:157:GLU:HA	1:B:238:ARG:NH2	2.16	0.55
1:B:22:LEU:HD11	1:B:103:LEU:HD22	1.91	0.52
1:A:27:ILE:HG22	1:A:29:THR:HG23	1.96	0.48
1:A:55:PHE:HZ	1:A:143:MET:HE1	1.79	0.47
1:A:98:ARG:HB2	4:A:376:HOH:O	2.18	0.43
1:A:55:PHE:CZ	1:A:143:MET:HE1	2.54	0.42
1:B:10:ARG:O	1:B:218:THR:HA	2.19	0.42
1:B:1:MET:HG3	1:B:228:HIS:CE1	2.55	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/265 (94%)	244 (98%)	4 (2%)	0	100	100
1	B	236/265 (89%)	233 (99%)	3 (1%)	0	100	100
All	All	484/530 (91%)	477 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/225 (93%)	206 (98%)	4 (2%)	57	61
1	B	202/225 (90%)	200 (99%)	2 (1%)	76	81
All	All	412/450 (92%)	406 (98%)	6 (2%)	65	69

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	LYS
1	A	98	ARG
1	A	115	ARG
1	A	222	ARG
1	B	57	SER
1	B	237	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	B	141	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	266	-	4,4,4	0.86	0	6,6,6	0.38	0
2	PO4	B	267	-	4,4,4	0.80	0	6,6,6	0.51	0
2	PO4	A	267	-	4,4,4	0.53	0	6,6,6	0.52	0
2	PO4	B	266	-	4,4,4	0.79	0	6,6,6	0.78	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/265 (94%)	0.04	20 (8%) 12 11	16, 22, 47, 56	0
1	B	240/265 (90%)	0.12	22 (9%) 9 8	16, 22, 49, 57	0
All	All	490/530 (92%)	0.08	42 (8%) 10 9	16, 22, 49, 57	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	232	SER	7.6
1	B	231	GLU	6.2
1	B	114	CYS	5.5
1	A	111	GLY	5.3
1	A	110	ALA	5.1
1	A	114	CYS	4.9
1	A	155	PRO	4.6
1	B	111	GLY	4.6
1	B	230	GLU	4.4
1	A	108	ASN	4.3
1	B	1	MET	4.3
1	B	115	ARG	4.0
1	B	110	ALA	4.0
1	A	109	ALA	3.8
1	A	115	ARG	3.8
1	B	108	ASN	3.7
1	B	116	ASP	3.7
1	B	113	SER	3.5
1	A	158	ALA	3.5
1	B	233	VAL	3.5
1	A	157	GLU	3.4
1	B	229	ARG	3.4
1	A	116	ASP	3.3
1	A	112	GLN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	101	GLU	3.1
1	B	78	ALA	3.0
1	B	109	ALA	2.9
1	B	8	ILE	2.8
1	A	8	ILE	2.7
1	A	78	ALA	2.7
1	A	113	SER	2.6
1	A	232	SER	2.6
1	A	101	GLU	2.5
1	A	6	LEU	2.3
1	B	235	ARG	2.2
1	B	251	GLU	2.2
1	A	153	SER	2.1
1	B	112	GLN	2.1
1	A	159	ASP	2.1
1	A	231	GLU	2.1
1	B	187	ILE	2.0
1	B	228	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	B	267	5/5	0.95	0.10	46,46,48,48	0
2	PO4	A	267	5/5	0.95	0.12	46,46,47,49	0
2	PO4	A	266	5/5	0.99	0.10	24,25,27,28	0
2	PO4	B	266	5/5	1.00	0.09	25,25,28,28	0
3	K	A	268	1/1	1.00	0.04	24,24,24,24	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	K	B	268	1/1	1.00	0.04	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.