



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2024 – 12:54 AM EST

PDB ID : 3FID
Title : LpxR from Salmonella typhimurium
Authors : Rutten, L.; Gros, P.
Deposited on : 2008-12-11
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	
1	B	296	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5189 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative outer membrane protein (LpxR).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	296	Total 2294	C 1455	N 393	O 444	S 2	0	0	0
1	B	296	Total 2294	C 1455	N 393	O 444	S 2	0	0	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total 5	Zn 5	0	0
2	B	4	Total 4	Zn 4	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is PENTAETHYLENE GLYCOL MONODECYL ETHER (three-letter code: CXE) (formula: $C_{20}H_{42}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	12	1		
4	A	1	Total	C	O	0	0
			20	16	4		
4	A	1	Total	C	O	0	0
			15	13	2		
4	A	1	Total	C	O	0	0
			26	20	6		
4	B	1	Total	C	O	0	0
			13	12	1		
4	B	1	Total	C	O	6	0
			26	20	6		
4	B	1	Total	C	O	0	0
			15	13	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	B	1	26	20	6	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	202	202	202	0	0
5	B	212	212	212	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative outer membrane protein (LpxR)

Chain A:  95% 5%



- Molecule 1: Putative outer membrane protein (LpxR)

Chain B:  93% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	106.93Å 127.65Å 60.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.74 – 1.90 48.76 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.74-1.90) 100.0 (48.76-1.90)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.191 , 0.230 0.191 , 0.229	Depositor DCC
R_{free} test set	3357 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	16.7	Xtrriage
Anisotropy	0.445	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 63.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5189	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0450e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CXE, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/2346	0.78	5/3180 (0.2%)
1	B	0.71	0/2346	0.80	6/3180 (0.2%)
All	All	0.71	0/4692	0.79	11/6360 (0.2%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	187	LEU	CA-CB-CG	7.08	131.58	115.30
1	A	256	ARG	NE-CZ-NH2	6.52	123.56	120.30
1	B	77	ARG	NE-CZ-NH2	-6.49	117.06	120.30
1	B	256	ARG	NE-CZ-NH1	-6.45	117.07	120.30
1	B	256	ARG	NE-CZ-NH2	6.41	123.50	120.30
1	A	270	LEU	CA-CB-CG	5.89	128.86	115.30
1	B	100	ARG	NE-CZ-NH2	-5.79	117.41	120.30
1	A	256	ARG	NE-CZ-NH1	-5.58	117.51	120.30
1	A	100	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	3	LEU	CA-CB-CG	5.21	127.28	115.30
1	B	204	ARG	NE-CZ-NH1	5.07	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2294	0	2213	8	0
1	B	2294	0	2213	13	0
2	A	5	0	0	0	0
2	B	4	0	0	0	0
3	A	12	0	15	2	0
3	B	12	0	15	1	0
4	A	74	0	123	0	0
4	B	80	0	132	3	0
5	A	202	0	0	3	1
5	B	212	0	0	7	1
All	All	5189	0	4711	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:30:ARG:HD2	1:B:283:ASP:HB2	1.58	0.86
1:A:5:ILE:HD13	1:A:39:LEU:HD13	1.61	0.80
3:B:302:GOL:H2	5:B:553:HOH:O	1.85	0.75
3:A:303:GOL:H31	5:A:331:HOH:O	1.87	0.75
1:A:94:THR:HG21	1:A:96:TRP:CE2	2.22	0.74
1:B:131:PRO:O	1:B:134:GLN:HG2	1.88	0.74
1:B:44:ASP:HB3	5:B:349:HOH:O	1.94	0.68
1:B:283:ASP:OD1	5:B:367:HOH:O	2.10	0.67
3:A:303:GOL:H2	5:A:376:HOH:O	1.95	0.67
1:B:148:MET:CE	1:B:202:LEU:HB3	2.28	0.64
1:B:68:ARG:NH1	5:B:461:HOH:O	2.39	0.56
1:B:144:THR:HG22	5:B:383:HOH:O	2.07	0.54
1:B:228:VAL:O	1:B:250:LYS:HE2	2.10	0.51
1:A:44:ASP:HB3	5:A:481:HOH:O	2.11	0.50
1:B:151:LEU:HG	4:B:306:CXE:H101	1.93	0.50
1:A:104:ASP:HB2	1:A:144:THR:OG1	2.16	0.45
4:B:304:CXE:H162	4:B:304:CXE:H131	1.71	0.45
1:A:109:GLY:HA3	1:A:139:ASN:HA	2.00	0.44
1:A:94:THR:HG21	1:A:96:TRP:NE1	2.34	0.43
1:B:191:LYS:HE3	5:B:495:HOH:O	2.18	0.43
1:A:263:TYR:O	1:A:264:SER:C	2.56	0.43
1:A:94:THR:HG23	1:A:96:TRP:H	1.83	0.43
1:B:148:MET:HE3	1:B:202:LEU:HB3	2.00	0.42
1:B:256:ARG:NH1	5:B:517:HOH:O	2.39	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:PRO:HB2	4:B:306:CXE:H122	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:521:HOH:O	5:B:361:HOH:O[4_556]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/296 (99%)	290 (99%)	4 (1%)	0	100	100
1	B	294/296 (99%)	290 (99%)	4 (1%)	0	100	100
All	All	588/592 (99%)	580 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/237 (100%)	235 (99%)	2 (1%)	81	82
1	B	237/237 (100%)	235 (99%)	2 (1%)	81	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	474/474 (100%)	470 (99%)	4 (1%)	81	82

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	270	LEU
1	A	291	ASP
1	B	187	LEU
1	B	291	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 9 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	CXE	B	306	-	25,25,25	0.41	0	24,24,24	0.31	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CXE	A	305	-	19,19,25	0.42	0	18,18,24	0.39	0
4	CXE	B	303	-	12,12,25	0.35	0	11,11,24	0.56	0
3	GOL	A	303	-	5,5,5	0.32	0	5,5,5	0.55	0
3	GOL	A	302	-	5,5,5	0.29	0	5,5,5	1.09	0
4	CXE	A	306	-	14,14,25	0.35	0	13,13,24	0.52	0
4	CXE	B	304	-	25,25,25	4.07	1 (4%)	24,24,24	1.09	2 (8%)
3	GOL	B	301	-	5,5,5	0.47	0	5,5,5	0.64	0
3	GOL	B	302	-	5,5,5	0.33	0	5,5,5	0.51	0
4	CXE	A	304	-	12,12,25	0.28	0	11,11,24	0.63	0
4	CXE	A	307	-	25,25,25	0.49	0	24,24,24	0.41	0
4	CXE	B	305	-	14,14,25	0.32	0	13,13,24	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CXE	B	306	-	-	14/23/23/23	-
4	CXE	A	305	-	-	8/17/17/23	-
4	CXE	B	303	-	-	0/10/10/23	-
3	GOL	A	303	-	-	2/4/4/4	-
3	GOL	A	302	-	-	4/4/4/4	-
4	CXE	A	306	-	-	7/12/12/23	-
4	CXE	B	304	-	-	18/23/23/23	-
3	GOL	B	301	-	-	0/4/4/4	-
3	GOL	B	302	-	-	1/4/4/4	-
4	CXE	A	304	-	-	2/10/10/23	-
4	CXE	A	307	-	-	11/23/23/23	-
4	CXE	B	305	-	-	8/12/12/23	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	304	CXE	O20-C21	20.20	2.29	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	304	CXE	O20-C21-C22	3.76	127.33	110.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	304	CXE	O23-C22-C21	2.18	120.22	110.39

There are no chirality outliers.

All (75) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	GOL	C1-C2-C3-O3
3	A	303	GOL	C1-C2-C3-O3
3	A	303	GOL	O2-C2-C3-O3
4	B	306	CXE	O20-C21-C22-O23
4	A	307	CXE	O11-C12-C13-O14
4	A	307	CXE	O17-C18-C19-O20
4	A	307	CXE	O14-C15-C16-O17
4	A	305	CXE	O17-C18-C19-O20
4	B	304	CXE	O23-C24-C25-O26
4	B	306	CXE	O23-C24-C25-O26
4	B	304	CXE	C16-C15-O14-C13
4	B	305	CXE	O11-C10-C9-C8
4	B	304	CXE	C5-C6-C7-C8
4	B	305	CXE	C6-C7-C8-C9
4	B	304	CXE	C2-C3-C4-C5
4	B	306	CXE	O11-C12-C13-O14
4	B	306	CXE	C4-C5-C6-C7
4	B	306	CXE	C6-C7-C8-C9
4	B	306	CXE	O14-C15-C16-O17
4	A	306	CXE	C7-C8-C9-C10
4	B	305	CXE	C7-C8-C9-C10
4	A	306	CXE	C2-C3-C4-C5
4	A	306	CXE	C3-C4-C5-C6
4	B	306	CXE	C5-C6-C7-C8
4	A	307	CXE	C7-C8-C9-C10
4	B	304	CXE	O17-C18-C19-O20
4	A	306	CXE	C1-C2-C3-C4
4	B	304	CXE	C4-C5-C6-C7
3	A	302	GOL	O1-C1-C2-O2
3	A	302	GOL	O2-C2-C3-O3
4	B	304	CXE	C1-C2-C3-C4
4	B	305	CXE	C1-C2-C3-C4
4	A	305	CXE	C3-C4-C5-C6
4	B	304	CXE	C3-C4-C5-C6
4	A	306	CXE	O11-C10-C9-C8
4	B	306	CXE	C7-C8-C9-C10

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	B	304	CXE	C6-C7-C8-C9
4	A	306	CXE	C6-C7-C8-C9
4	B	305	CXE	C3-C4-C5-C6
4	B	305	CXE	C12-C13-O14-C15
4	A	305	CXE	C4-C5-C6-C7
4	B	304	CXE	C21-C22-O23-C24
4	A	307	CXE	C13-C12-O11-C10
4	B	306	CXE	C12-C13-O14-C15
4	B	306	CXE	C21-C22-O23-C24
4	A	307	CXE	C22-C21-O20-C19
4	A	305	CXE	C15-C16-O17-C18
4	B	304	CXE	C25-C24-O23-C22
4	A	307	CXE	C25-C24-O23-C22
4	B	304	CXE	C22-C21-O20-C19
4	A	305	CXE	C1-C2-C3-C4
4	B	306	CXE	C18-C19-O20-C21
4	B	304	CXE	C15-C16-O17-C18
4	B	304	CXE	C18-C19-O20-C21
4	A	307	CXE	C1-C2-C3-C4
4	A	307	CXE	C15-C16-O17-C18
4	A	306	CXE	C4-C5-C6-C7
4	A	304	CXE	C6-C7-C8-C9
4	B	306	CXE	O11-C10-C9-C8
4	B	306	CXE	C1-C2-C3-C4
3	B	302	GOL	C1-C2-C3-O3
4	B	306	CXE	C13-C12-O11-C10
4	B	304	CXE	C12-C13-O14-C15
4	B	305	CXE	C4-C5-C6-C7
4	B	304	CXE	O11-C12-C13-O14
4	A	305	CXE	C12-C13-O14-C15
3	A	302	GOL	O1-C1-C2-C3
4	A	305	CXE	O14-C15-C16-O17
4	A	304	CXE	O11-C10-C9-C8
4	B	304	CXE	O14-C15-C16-O17
4	B	304	CXE	O20-C21-C22-O23
4	A	305	CXE	O11-C12-C13-O14
4	A	307	CXE	C9-C10-O11-C12
4	A	307	CXE	O20-C21-C22-O23
4	B	305	CXE	C13-C12-O11-C10


There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	306	CXE	2	0
3	A	303	GOL	2	0
4	B	304	CXE	1	0
3	B	302	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/296 (100%)	-0.27	2 (0%) 87 88	9, 15, 26, 36	0
1	B	296/296 (100%)	-0.30	3 (1%) 82 84	8, 15, 27, 37	0
All	All	592/592 (100%)	-0.28	5 (0%) 86 87	8, 15, 27, 37	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	211	GLN	3.5
1	B	211	GLN	2.9
1	B	96	TRP	2.6
1	B	128	GLU	2.3
1	A	89	TRP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CXE	B	303	13/26	0.80	0.14	26,28,37,39	0
4	CXE	B	304	26/26	0.80	0.17	20,32,49,50	6
4	CXE	A	307	26/26	0.81	0.21	37,49,53,54	0
4	CXE	B	306	26/26	0.84	0.21	36,50,55,56	0
3	GOL	A	303	6/6	0.85	0.19	35,37,40,42	0
4	CXE	A	305	20/26	0.85	0.15	30,35,51,51	0
4	CXE	A	304	13/26	0.88	0.13	24,28,37,38	0
3	GOL	B	302	6/6	0.93	0.15	22,31,32,35	0
4	CXE	B	305	15/26	0.93	0.15	26,28,40,41	0
4	CXE	A	306	15/26	0.93	0.13	23,27,36,38	0
3	GOL	B	301	6/6	0.95	0.14	19,23,24,24	0
3	GOL	A	302	6/6	0.95	0.14	22,24,25,26	0
2	ZN	A	298	1/1	0.98	0.04	31,31,31,31	0
2	ZN	A	297	1/1	0.99	0.03	35,35,35,35	0
2	ZN	B	297	1/1	0.99	0.04	34,34,34,34	0
2	ZN	B	298	1/1	0.99	0.04	32,32,32,32	0
2	ZN	B	300	1/1	0.99	0.10	22,22,22,22	1
2	ZN	A	299	1/1	1.00	0.08	15,15,15,15	0
2	ZN	A	300	1/1	1.00	0.07	13,13,13,13	0
2	ZN	B	299	1/1	1.00	0.09	13,13,13,13	0
2	ZN	A	301	1/1	1.00	0.13	27,27,27,27	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers [i](#)

There are no such residues in this entry.