



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 25, 2023 – 03:14 AM EDT

PDB ID : 3H1C
Title : Crystal structure of Polynucleotide Phosphorylase (PNPase) core bound to RNase E and Tungstate
Authors : Nurmohamed, S.
Deposited on : 2009-04-11
Resolution : 3.57 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






















Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.66-3.50)
Clashscore	141614	1181 (3.66-3.50)
Ramachandran outliers	138981	1143 (3.66-3.50)
Sidechain outliers	138945	1143 (3.66-3.50)
RSRZ outliers	127900	1012 (3.66-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	 77% 22% ..
1	B	549	 76% 23% ..
1	C	549	 75% 23% ..
1	G	549	 77% 21% ..
1	I	549	 77% 21% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	549	 80% 18% ..
1	M	549	 82% 16% ..
1	O	549	 79% 19% ..
1	R	549	 81% 18% .
1	T	549	 78% 19% ..
1	V	549	 76% 22% ..
1	X	549	 78% 20% ..
2	D	41	 39% 12% 49%
2	E	41	 37% 15% 49% 2%
2	F	41	 39% 12% 49% 15%
2	H	41	 46% 5% 49% 7%
2	J	41	 44% 7% 49% 2%
2	L	41	 39% 12% 49%
2	N	41	 51% 49%
2	P	41	 49% 49% 2%
2	S	41	 32% 20% 49% 7%
2	U	41	 44% 7% 49% 5%
2	W	41	 37% 10% 49% 12%
2	Y	41	 37% 15% 49% 2%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	WO4	M	551	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 51109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyribonucleotide nucleotidyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	544	Total 4075	C 2559	N 704	O 792	S 20	0	0	0
1	B	544	Total 4107	C 2575	N 717	O 797	S 18	0	0	0
1	C	544	Total 4121	C 2583	N 720	O 798	S 20	0	0	0
1	G	543	Total 4095	C 2571	N 715	O 789	S 20	0	0	0
1	I	543	Total 4109	C 2576	N 716	O 797	S 20	0	0	0
1	K	544	Total 4127	C 2588	N 721	O 798	S 20	0	0	0
1	M	544	Total 4116	C 2581	N 718	O 797	S 20	0	0	0
1	O	544	Total 4124	C 2584	N 723	O 798	S 19	0	0	0
1	R	544	Total 4134	C 2590	N 724	O 800	S 20	0	0	0
1	T	543	Total 4116	C 2582	N 720	O 795	S 19	0	0	0
1	V	544	Total 4118	C 2580	N 719	O 799	S 20	0	0	0
1	X	544	Total 4113	C 2581	N 717	O 795	S 20	0	0	0

- Molecule 2 is a protein called Ribonuclease E.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	21	Total 138	C 82	N 31	O 25	0	0	0
2	E	21	Total 138	C 82	N 31	O 25	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	H	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	J	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	L	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	N	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	P	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	S	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	U	21	Total	C	N	O	0	0	0
			138	82	31	25			
2	W	21	Total	C	N	O	0	0	0
			131	78	29	24			
2	Y	21	Total	C	N	O	0	0	0
			138	82	31	25			

- Molecule 3 is TUNGSTATE(VI)ION (three-letter code: WO4) (formula: O₄W).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O W	0	0
			5	4 1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 5	O 4	W 1	0	0
3	B	1	Total 5	O 4	W 1	0	0
3	B	1	Total 5	O 4	W 1	0	0
3	C	1	Total 5	O 4	W 1	0	0
3	G	1	Total 5	O 4	W 1	0	0
3	I	1	Total 5	O 4	W 1	0	0
3	K	1	Total 5	O 4	W 1	0	0
3	K	1	Total 5	O 4	W 1	0	0
3	M	1	Total 5	O 4	W 1	0	0
3	M	1	Total 5	O 4	W 1	0	0
3	O	1	Total 5	O 4	W 1	0	0
3	O	1	Total 5	O 4	W 1	0	0
3	R	1	Total 5	O 4	W 1	0	0
3	R	1	Total 5	O 4	W 1	0	0
3	T	1	Total 5	O 4	W 1	0	0
3	T	1	Total 5	O 4	W 1	0	0
3	V	1	Total 5	O 4	W 1	0	0
3	V	1	Total 5	O 4	W 1	0	0
3	X	1	Total 5	O 4	W 1	0	0
3	X	1	Total 5	O 4	W 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polyrribonucleotide nucleotidyltransferase




- Molecule 1: Polyrribonucleotide nucleotidyltransferase





- Molecule 1: Polyrribonucleotide nucleotidyltransferase









● Molecule 1: Polyribonucleotide nucleotidyltransferase





● Molecule 1: Polyribonucleotide nucleotidyltransferase




● Molecule 1: Polyribonucleotide nucleotidyltransferase

Chain K:  80% 18%


- Molecule 1: Polyribonucleotide nucleotidyltransferase


Chain M:  82% 16%

- Molecule 1: Polyribonucleotide nucleotidyltransferase



Chain O:  79% 19%

- Molecule 1: Polyribonucleotide nucleotidyltransferase



Chain R:  81% 18%




• Molecule 1: Polyribonucleotide nucleotidyltransferase




• Molecule 1: Polyribonucleotide nucleotidyltransferase




• Molecule 1: Polyribonucleotide nucleotidyltransferase




- Molecule 2: Ribonuclease E




- Molecule 2: Ribonuclease E




- Molecule 2: Ribonuclease E




- Molecule 2: Ribonuclease E





- Molecule 2: Ribonuclease E







● Molecule 2: Ribonuclease E




● Molecule 2: Ribonuclease E





● Molecule 2: Ribonuclease E





● Molecule 2: Ribonuclease E





● Molecule 2: Ribonuclease E



● Molecule 2: Ribonuclease E



● Molecule 2: Ribonuclease E



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	167.74Å 262.89Å 264.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 3.57 20.15 – 3.57	Depositor EDS
% Data completeness (in resolution range)	94.2 (25.00-3.57) 94.2 (20.15-3.57)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 3.62Å)	Xtrriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.270 , 0.304 0.267 , 0.291	Depositor DCC
R_{free} test set	6514 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	79.4	Xtrriage
Anisotropy	0.028	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 26.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.038 for -h,l,k	Xtrriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	51109	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: WO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/4138	0.47	0/5621
1	B	0.30	0/4170	0.46	0/5660
1	C	0.37	0/4184	0.47	1/5675 (0.0%)
1	G	0.41	0/4158	0.51	2/5643 (0.0%)
1	I	0.38	0/4172	0.50	2/5660 (0.0%)
1	K	0.37	0/4190	0.54	4/5681 (0.1%)
1	M	0.33	0/4179	0.58	9/5668 (0.2%)
1	O	0.34	0/4187	0.47	2/5679 (0.0%)
1	R	0.30	0/4197	0.46	0/5690
1	T	0.32	0/4179	0.48	1/5667 (0.0%)
1	V	0.36	0/4180	0.47	1/5671 (0.0%)
1	X	0.30	0/4175	0.48	2/5661 (0.0%)
2	D	0.34	0/143	0.47	0/196
2	E	0.31	0/143	0.38	0/196
2	F	0.32	0/143	0.49	0/196
2	H	0.32	0/143	0.39	0/196
2	J	0.33	0/143	0.48	0/196
2	L	0.36	0/143	0.47	0/196
2	N	0.33	0/143	0.38	0/196
2	P	0.33	0/143	0.39	0/196
2	S	0.32	0/143	0.38	0/196
2	U	0.32	0/143	0.40	0/196
2	W	0.49	0/135	0.68	0/185
2	Y	0.32	0/143	0.41	0/196
All	All	0.35	0/51817	0.49	24/70317 (0.0%)

There are no bond length outliers.

The worst 5 of 24 bond angle outliers are listed below:

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	438	SER	CB-CA-C	13.85	136.42	110.10
1	M	434	SER	CB-CA-C	-13.49	84.47	110.10
1	K	37	ASP	CB-CA-C	-13.04	84.33	110.40
1	M	435	ASN	N-CA-CB	-10.55	91.60	110.60
1	M	394	GLY	N-CA-C	8.54	134.46	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4075	0	4044	154	0
1	B	4107	0	4096	134	0
1	C	4121	0	4126	186	0
1	G	4095	0	4087	168	0
1	I	4109	0	4108	138	0
1	K	4127	0	4141	148	0
1	M	4116	0	4116	98	0
1	O	4124	0	4130	119	0
1	R	4134	0	4150	101	0
1	T	4116	0	4127	160	0
1	V	4118	0	4115	173	0
1	X	4113	0	4117	176	0
2	D	138	0	125	5	0
2	E	138	0	125	8	0
2	F	138	0	125	16	0
2	H	138	0	125	2	0
2	J	138	0	125	2	0
2	L	138	0	125	8	0
2	N	138	0	125	0	0
2	P	138	0	125	1	0
2	S	138	0	125	9	0
2	U	138	0	125	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	W	131	0	115	21	0
2	Y	138	0	125	5	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
3	C	5	0	0	0	0
3	G	5	0	0	1	0
3	I	5	0	0	0	0
3	K	10	0	0	0	0
3	M	10	0	0	0	0
3	O	10	0	0	0	0
3	R	10	0	0	0	0
3	T	10	0	0	0	0
3	V	10	0	0	0	0
3	X	10	0	0	1	0
All	All	51109	0	50847	1659	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 1659 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:310:LEU:HD21	1:I:469:VAL:CG1	1.47	1.44
1:G:26:GLN:NE2	1:K:387:VAL:HA	1.36	1.40
1:R:51:LYS:HB3	1:R:54:GLN:CG	1.52	1.36
1:O:239:ASN:ND2	1:O:242:LEU:HB2	1.40	1.35
1:V:342:LEU:HD11	1:V:349:GLN:NE2	1.40	1.32

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	542/549 (99%)	496 (92%)	40 (7%)	6 (1%)	14	53
1	B	542/549 (99%)	508 (94%)	30 (6%)	4 (1%)	22	62
1	C	542/549 (99%)	497 (92%)	39 (7%)	6 (1%)	14	53
1	G	541/549 (98%)	500 (92%)	38 (7%)	3 (1%)	25	64
1	I	541/549 (98%)	502 (93%)	36 (7%)	3 (1%)	25	64
1	K	542/549 (99%)	502 (93%)	34 (6%)	6 (1%)	14	53
1	M	542/549 (99%)	501 (92%)	37 (7%)	4 (1%)	22	62
1	O	542/549 (99%)	507 (94%)	32 (6%)	3 (1%)	25	64
1	R	542/549 (99%)	505 (93%)	33 (6%)	4 (1%)	22	62
1	T	541/549 (98%)	499 (92%)	36 (7%)	6 (1%)	14	53
1	V	542/549 (99%)	506 (93%)	33 (6%)	3 (1%)	25	64
1	X	542/549 (99%)	510 (94%)	29 (5%)	3 (1%)	25	64
2	D	19/41 (46%)	19 (100%)	0	0	100	100
2	E	19/41 (46%)	19 (100%)	0	0	100	100
2	F	19/41 (46%)	17 (90%)	2 (10%)	0	100	100
2	H	19/41 (46%)	19 (100%)	0	0	100	100
2	J	19/41 (46%)	18 (95%)	1 (5%)	0	100	100
2	L	19/41 (46%)	19 (100%)	0	0	100	100
2	N	19/41 (46%)	17 (90%)	2 (10%)	0	100	100
2	P	19/41 (46%)	18 (95%)	1 (5%)	0	100	100
2	S	19/41 (46%)	19 (100%)	0	0	100	100
2	U	19/41 (46%)	18 (95%)	1 (5%)	0	100	100
2	W	19/41 (46%)	13 (68%)	4 (21%)	2 (10%)	0	7
2	Y	19/41 (46%)	19 (100%)	0	0	100	100
All	All	6729/7080 (95%)	6248 (93%)	428 (6%)	53 (1%)	19	59

5 of 53 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	38	THR
1	G	187	LEU
1	G	189	VAL
1	O	37	ASP
1	O	38	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/446 (96%)	422 (99%)	4 (1%)	78	90
1	B	432/446 (97%)	430 (100%)	2 (0%)	88	95
1	C	436/446 (98%)	429 (98%)	7 (2%)	62	83
1	G	429/446 (96%)	426 (99%)	3 (1%)	84	93
1	I	435/446 (98%)	432 (99%)	3 (1%)	84	93
1	K	437/446 (98%)	433 (99%)	4 (1%)	78	90
1	M	434/446 (97%)	429 (99%)	5 (1%)	71	87
1	O	436/446 (98%)	430 (99%)	6 (1%)	67	85
1	R	439/446 (98%)	437 (100%)	2 (0%)	88	95
1	T	435/446 (98%)	432 (99%)	3 (1%)	84	93
1	V	435/446 (98%)	434 (100%)	1 (0%)	93	98
1	X	433/446 (97%)	430 (99%)	3 (1%)	84	93
2	D	11/28 (39%)	11 (100%)	0	100	100
2	E	11/28 (39%)	11 (100%)	0	100	100
2	F	11/28 (39%)	11 (100%)	0	100	100
2	H	11/28 (39%)	11 (100%)	0	100	100
2	J	11/28 (39%)	11 (100%)	0	100	100
2	L	11/28 (39%)	11 (100%)	0	100	100
2	N	11/28 (39%)	11 (100%)	0	100	100
2	P	11/28 (39%)	11 (100%)	0	100	100
2	S	11/28 (39%)	11 (100%)	0	100	100
2	U	11/28 (39%)	11 (100%)	0	100	100
2	W	9/28 (32%)	8 (89%)	1 (11%)	6	31
2	Y	11/28 (39%)	11 (100%)	0	100	100
All	All	5337/5688 (94%)	5293 (99%)	44 (1%)	81	92

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	O	124	VAL
1	R	358	THR
1	O	222	LEU
1	O	358	THR
1	T	435	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 95 such sidechains are listed below:

Mol	Chain	Res	Type
1	O	349	GLN
2	S	1058	GLN
1	O	525	GLN
1	R	213	GLN
1	T	160	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	WO4	B	551	-	2,4,4	12.57	2 (100%)	-		
3	WO4	I	550	-	2,4,4	12.56	2 (100%)	-		
3	WO4	G	550	-	2,4,4	12.56	2 (100%)	-		
3	WO4	X	550	-	2,4,4	12.57	2 (100%)	-		
3	WO4	M	551	-	2,4,4	12.57	2 (100%)	-		
3	WO4	V	550	-	2,4,4	12.58	2 (100%)	-		
3	WO4	A	550	-	2,4,4	12.58	2 (100%)	-		
3	WO4	V	551	-	2,4,4	12.57	2 (100%)	-		
3	WO4	T	551	-	2,4,4	12.59	2 (100%)	-		
3	WO4	O	550	-	2,4,4	12.59	2 (100%)	-		
3	WO4	O	551	-	2,4,4	12.58	2 (100%)	-		
3	WO4	A	551	-	2,4,4	12.58	2 (100%)	-		
3	WO4	R	550	-	2,4,4	12.57	2 (100%)	-		
3	WO4	M	550	-	2,4,4	11.54	2 (100%)	-		
3	WO4	R	551	-	2,4,4	12.58	2 (100%)	-		
3	WO4	K	550	-	2,4,4	12.55	2 (100%)	-		
3	WO4	C	550	-	2,4,4	12.56	2 (100%)	-		
3	WO4	K	551	-	2,4,4	12.57	2 (100%)	-		
3	WO4	T	550	-	2,4,4	12.57	2 (100%)	-		
3	WO4	X	551	-	2,4,4	12.59	2 (100%)	-		
3	WO4	B	550	-	2,4,4	12.60	2 (100%)	-		

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	550	WO4	W-O2	12.61	2.04	1.74
3	X	551	WO4	W-O2	12.60	2.04	1.74
3	A	550	WO4	W-O2	12.60	2.04	1.74
3	T	551	WO4	W-O2	12.60	2.04	1.74
3	O	550	WO4	W-O2	12.60	2.04	1.74

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	550	WO4	1	0
3	X	550	WO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	544/549 (99%)	-0.28	2 (0%) 92 86	15, 50, 83, 103	1 (0%)
1	B	544/549 (99%)	-0.36	0 100 100	9, 39, 76, 101	1 (0%)
1	C	544/549 (99%)	-0.30	1 (0%) 95 91	14, 48, 83, 106	1 (0%)
1	G	543/549 (98%)	-0.29	3 (0%) 89 80	20, 51, 83, 111	1 (0%)
1	I	543/549 (98%)	-0.33	4 (0%) 87 77	14, 40, 76, 98	1 (0%)
1	K	544/549 (99%)	-0.34	2 (0%) 92 86	13, 43, 82, 115	1 (0%)
1	M	544/549 (99%)	-0.21	5 (0%) 84 71	20, 54, 89, 110	1 (0%)
1	O	544/549 (99%)	-0.33	1 (0%) 95 91	8, 41, 79, 97	1 (0%)
1	R	544/549 (99%)	-0.27	2 (0%) 92 86	13, 46, 83, 101	1 (0%)
1	T	543/549 (98%)	-0.28	2 (0%) 92 86	20, 53, 84, 108	1 (0%)
1	V	544/549 (99%)	-0.21	5 (0%) 84 71	19, 62, 92, 105	1 (0%)
1	X	544/549 (99%)	-0.22	2 (0%) 92 86	20, 52, 85, 100	1 (0%)
2	D	21/41 (51%)	0.54	0 100 100	44, 80, 95, 96	0
2	E	21/41 (51%)	0.50	1 (4%) 30 18	34, 69, 98, 104	0
2	F	21/41 (51%)	1.18	6 (28%) 0 0	70, 92, 111, 117	0
2	H	21/41 (51%)	0.64	3 (14%) 2 1	62, 95, 107, 109	0
2	J	21/41 (51%)	0.32	1 (4%) 30 18	43, 70, 85, 91	0
2	L	21/41 (51%)	0.30	0 100 100	49, 78, 103, 107	0
2	N	21/41 (51%)	0.32	0 100 100	64, 84, 97, 97	0
2	P	21/41 (51%)	0.26	1 (4%) 30 18	33, 73, 93, 98	0
2	S	21/41 (51%)	0.80	3 (14%) 2 1	64, 96, 117, 120	0
2	U	21/41 (51%)	0.62	2 (9%) 8 4	49, 91, 107, 111	0
2	W	21/41 (51%)	1.42	5 (23%) 0 0	78, 111, 130, 134	0
2	Y	21/41 (51%)	0.38	1 (4%) 30 18	62, 90, 101, 103	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	6777/7080 (95%)	-0.25	52 (0%) 86 73	8, 49, 88, 134	12 (0%)

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	W	1044	HIS	4.8
1	C	542	ASN	3.9
2	U	1039	GLY	3.5
2	H	1045	THR	3.4
2	W	1039	GLY	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	WO4	M	551	5/5	0.73	0.52	131,133,133,137	5
3	WO4	M	550	5/5	0.78	0.39	112,113,118,118	5
3	WO4	V	550	5/5	0.80	0.35	107,108,112,113	5
3	WO4	T	551	5/5	0.81	0.35	113,114,117,119	5
3	WO4	K	550	5/5	0.83	0.34	78,79,82,88	5
3	WO4	G	550	5/5	0.86	0.30	79,79,88,93	5
3	WO4	A	551	5/5	0.87	0.36	103,105,108,113	5
3	WO4	C	550	5/5	0.88	0.28	80,84,89,90	5
3	WO4	K	551	5/5	0.89	0.33	96,97,101,108	5
3	WO4	B	550	5/5	0.90	0.23	72,73,77,82	5
3	WO4	T	550	5/5	0.90	0.37	103,104,113,115	5
3	WO4	O	550	5/5	0.91	0.24	100,100,105,109	5
3	WO4	V	551	5/5	0.91	0.33	102,102,105,109	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	WO4	B	551	5/5	0.92	0.32	88,89,93,97	5
3	WO4	R	551	5/5	0.92	0.34	86,88,95,98	5
3	WO4	X	550	5/5	0.92	0.24	94,99,99,100	5
3	WO4	A	550	5/5	0.93	0.23	84,86,88,93	5
3	WO4	O	551	5/5	0.93	0.25	71,75,80,87	5
3	WO4	R	550	5/5	0.93	0.26	106,107,111,116	5
3	WO4	X	551	5/5	0.94	0.43	88,90,91,93	5
3	WO4	I	550	5/5	0.95	0.23	89,90,91,91	5

6.5 Other polymers [i](#)

There are no such residues in this entry.