

wwPDB X-ray Structure Validation Summary Report (i)

Aug 28, 2023 – 05:16 AM EDT

with

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)		
EDS	:	2.35
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	383	70%	21%	•	6%
1	В	383	72%	18%	•	7%
2	Е	182	2% 8 0%	15%	, D	• • •
2	F	182	4%	21%		••



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Δ	360	Total	С	Ν	0	\mathbf{S}	0	0	0
	A	300	2946	1861	528	542	15	0	0	
1	В	357	Total	С	Ν	0	S	0 0	0	0
	D	557	2921	1847	523	536	15	0	0	0

• Molecule 1 is a protein called Rho guanine nucleotide exchange factor 11.

Chain	Residue	Modelled	Actual	Comment	Reference
A	709	GLY	-	expression tag	UNP O15085
А	1086	HIS	-	expression tag	UNP O15085
A	1087	HIS	-	expression tag	UNP O15085
A	1088	HIS	-	expression tag	UNP O15085
А	1089	HIS	-	expression tag	UNP O15085
A	1090	HIS	-	expression tag	UNP O15085
А	1091	HIS	-	expression tag	UNP O15085
В	709	GLY	-	expression tag	UNP O15085
В	1086	HIS	-	expression tag	UNP O15085
В	1087	HIS	-	expression tag	UNP O15085
В	1088	HIS	-	expression tag	UNP O15085
В	1089	HIS	-	expression tag	UNP O15085
В	1090	HIS	-	expression tag	UNP O15085
В	1091	HIS	-	expression tag	UNP O15085

There are 14 discrepancies between the modelled and reference sequences:

• Molecule 2 is a protein called Transforming protein RhoA.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace			
2	E	179	Total	С	N O S O	0	Ο				
2	Ľ	115	1419	897	240	272	10	0	0	0	
9	Б	179	Total	C N O	Ο	\mathbf{S}	0	0	0		
	Ľ	119	1419	897	240	272	10		U	0	

There are 2 discrepancies between the modelled and reference sequences:




Chain	Residue	Modelled	Actual	Comment	Reference
Е	0	GLY	-	expression tag	UNP P61586
F	0	GLY	-	expression tag	UNP P61586

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Е	1	Total Mg 1 1	0	0
3	F	1	Total Mg 1 1	0	0

• Molecule 4 is 5'-GUANOSINE-DIPHOSPHATE-MONOTHIOPHOSPHATE (three-letter code: GSP) (formula: $C_{10}H_{16}N_5O_{13}P_3S$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
4	E	1	Total	С	Ν	Ο	Р	S	0	0
4	Ľ	1	32	10	5	13	3	1	0	0
4	Б	1	Total	С	Ν	Ο	Р	S	0	0
4	Ľ	1	32	10	5	13	3	1	0	0

• Molecule 5 is water.

[Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	5	А	36	Total O 36 36	0	0
	5	В	18	Total O 18 18	0	0

Continued on next page...




Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	Е	12	Total O 12 12	0	0
5	F	9	Total O 9 9	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Rho guanine nucleotide exchange factor 11

• Molecule 2: Transforming protein RhoA





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	98.17Å 111.84Å 138.39Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.45 - 2.70	Depositor
Resolution (A)	33.45 - 2.70	EDS
% Data completeness	90.8 (33.45-2.70)	Depositor
(in resolution range)	90.8(33.45-2.70)	EDS
R _{merge}	0.09	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.30 (at 2.68 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.234 , 0.283	Depositor
II, II, <i>free</i>	0.242 , 0.282	DCC
R_{free} test set	1947 reflections (5.05%)	wwPDB-VP
Wilson B-factor $(Å^2)$	52.8	Xtriage
Anisotropy	0.941	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.29 , 38.5	EDS
L-test for twinning ²	$ \langle L \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8846	wwPDB-VP
Average B, all atoms $(Å^2)$	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GSP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		Bond lengths		ond angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.74	1/2991~(0.0%)	0.76	0/4026
1	В	0.66	0/2966	0.71	0/3992
2	Е	0.63	0/1447	0.65	1/1956~(0.1%)
2	F	0.59	0/1447	0.68	0/1956
All	All	0.67	1/8851~(0.0%)	0.71	1/11930~(0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
1	А	1007	CYS	CB-SG	-5.30	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Ε	8	LEU	CA-CB-CG	5.21	127.30	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2946	0	3037	60	0
1	В	2921	0	3014	50	0

Continued on next page...



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Е	1419	0	1408	27	0
2	F	1419	0	1408	29	0
3	Е	1	0	0	0	0
3	F	1	0	0	0	0
4	Ε	32	0	12	2	0
4	F	32	0	12	1	0
5	А	36	0	0	1	0
5	В	18	0	0	0	0
5	Е	12	0	0	2	0
5	F	9	0	0	2	0
All	All	8846	0	8891	158	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 158 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:719:THR:O	1:B:720:VAL:HG22	1.42	1.16
1:A:731:ARG:NH2	1:A:897:GLU:HG2	1.67	1.08
2:F:8:LEU:C	2:F:8:LEU:HD12	1.78	1.02
1:A:972:LEU:HD22	1:A:986:VAL:HG11	1.45	0.95
1:B:719:THR:O	1:B:720:VAL:CG2	2.16	0.93

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	356/383~(93%)	341 (96%)	14 (4%)	1 (0%)	41 66
1	В	353/383~(92%)	330 (94%)	19 (5%)	4 (1%)	14 34

Continued on next page...



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
2	Ε	177/182~(97%)	168~(95%)	8 (4%)	1 (1%)	25	50
2	F	177/182~(97%)	169~(96%)	7 (4%)	1 (1%)	25	50
All	All	1063/1130~(94%)	1008 (95%)	48 (4%)	7(1%)	22	46

Continued from previous page...

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	949	SER
2	F	109	ASN
1	В	960	ASP
1	В	726	ALA
1	В	1051	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	328/347~(94%)	304~(93%)	24 (7%)	14 33
1	В	325/347~(94%)	294 (90%)	31 (10%)	8 20
2	Ε	156/157~(99%)	146 (94%)	10 (6%)	17 39
2	F	156/157~(99%)	142 (91%)	14 (9%)	9 22
All	All	965/1008~(96%)	886~(92%)	79 (8%)	11 26

5 of 79 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	Ε	68	ARG
2	F	129	ARG
2	Е	87	ASP
2	F	8	LEU
2	F	145	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:



Mol	Chain	Res	Type
1	В	926	GLN
2	Е	180	GLN
2	F	149	ASN
2	F	94	ASN
1	А	880	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Dec	Link	Bo	ond leng	ths	B	ond ang	les
	туре		nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2							
4	GSP	Е	538	3	26,34,34	1.76	3 (11%)	27,54,54	1.61	6 (22%)							
4	GSP	F	538	3	26,34,34	1.94	3 (11%)	27,54,54	1.41	5 (18%)							

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GSP	Е	538	3	-	2/17/38/38	0/3/3/3
4	GSP	F	538	3	-	1/17/38/38	0/3/3/3

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
4	F	538	GSP	PG-S1G	-8.05	1.73	1.90
4	Е	538	GSP	PG-S1G	-6.72	1.76	1.90
4	F	538	GSP	C5-C6	-3.92	1.39	1.47
4	Е	538	GSP	C5-C6	-3.86	1.39	1.47
4	Е	538	GSP	C2-N3	2.26	1.38	1.33

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	Ε	538	GSP	C5-C6-N1	3.49	120.11	113.95
4	Ε	538	GSP	C2-N1-C6	-2.96	119.64	125.10
4	Е	538	GSP	C8-N7-C5	2.92	108.54	102.99
4	F	538	GSP	C5-C6-N1	2.78	118.87	113.95
4	F	538	GSP	C8-N7-C5	2.74	108.21	102.99

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	Ε	538	GSP	PA-O3A-PB-O2B
4	F	538	GSP	PA-O3A-PB-O2B
4	Е	538	GSP	PA-O3A-PB-O1B

There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Е	538	GSP	2	0
4	F	538	GSP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	360/383~(93%)	-0.09	0 100 100	40, 58, 84, 95	0
1	В	357/383~(93%)	0.02	4 (1%) 80 82	46, 65, 95, 110	0
2	Ε	179/182~(98%)	0.34	4 (2%) 62 63	41, 70, 142, 171	0
2	F	179/182~(98%)	0.45	7 (3%) 39 38	45, 86, 171, 207	0
All	All	1075/1130~(95%)	0.11	15 (1%) 75 77	40, 66, 113, 207	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	947	ARG	4.0
2	F	86	ILE	3.9
2	F	89	PRO	3.6
1	В	979	ASP	3.1
2	F	134	MET	3.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q < 0.9
3	MG	Е	550	1/1	0.94	0.24	$59,\!59,\!59,\!59$	0
4	GSP	F	538	32/32	0.94	0.14	$60,\!83,\!98,\!105$	0
4	GSP	Е	538	32/32	0.97	0.11	55,73,88,90	0
3	MG	F	550	1/1	0.98	0.16	57,57,57,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

