



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 08:22 am BST

PDB ID : 3LH5
Title : Crystal Structure of the SH3-Guanylate kinase core domain of ZO-1
Authors : Lye, M.F.; Lavie, A.
Deposited on : 2010-01-21
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	251	 77% 18% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tight junction protein ZO-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	248	2026	1285	369	368	4	0	0	0


- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	73	Total	O	0	0
			73	73		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tight junction protein ZO-1

Chain A:  77% 18% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	125.80 Å 125.80 Å 35.60 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	27.24 – 2.60 27.24 – 2.41	Depositor EDS
% Data completeness (in resolution range)	99.6 (27.24-2.60) 98.4 (27.24-2.41)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 2.42 Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.205 , 0.285 0.243 , 0.317	Depositor DCC
R_{free} test set	1245 reflections (9.94%)	wwPDB-VP
Wilson B-factor (Å ²)	59.1	Xtrriage
Anisotropy	0.131	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.044 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2099	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/2070	0.66	1/2795 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	740	ARG	NE-CZ-NH2	-6.95	116.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2026	0	2025	37	0
2	A	73	0	0	1	0
All	All	2099	0	2025	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (\AA)	Clash overlap (\AA)
1:A:635:ARG:HH11	1:A:635:ARG:HG2	1.17	1.06

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:779:ASN:HD22	1:A:781:GLY:H	1.23	0.87
1:A:635:ARG:CG	1:A:635:ARG:HH11	1.90	0.83
1:A:748:ARG:CD	1:A:748:ARG:H	1.93	0.80
1:A:635:ARG:NH1	1:A:635:ARG:HG2	1.92	0.79
1:A:748:ARG:H	1:A:748:ARG:HD2	1.49	0.75
1:A:779:ASN:ND2	1:A:781:GLY:H	1.87	0.72
1:A:640:GLU:HG2	1:A:797:LEU:CD2	2.29	0.62
1:A:748:ARG:N	1:A:748:ARG:HD2	2.12	0.60
1:A:586:VAL:HG12	2:A:67:HOH:O	2.02	0.59
1:A:748:ARG:HH11	1:A:749:LYS:HG3	1.68	0.59
1:A:709:THR:HG22	1:A:712:ALA:H	1.69	0.58
1:A:709:THR:CG2	1:A:712:ALA:H	2.17	0.57
1:A:760:LYS:HD3	1:A:760:LYS:C	2.25	0.57
1:A:650:PHE:HB3	1:A:761:LEU:HD13	1.88	0.55
1:A:729:ASN:HB3	1:A:774:ASN:HD22	1.75	0.52
1:A:788:GLU:OE1	1:A:792:GLN:NE2	2.41	0.52
1:A:779:ASN:HD22	1:A:779:ASN:C	2.12	0.52
1:A:757:ARG:NH1	1:A:760:LYS:HD2	2.29	0.47
1:A:560:ILE:HG12	1:A:571:ARG:HG3	1.97	0.46
1:A:779:ASN:ND2	1:A:779:ASN:C	2.68	0.46
1:A:729:ASN:HD22	1:A:774:ASN:ND2	2.14	0.46
1:A:748:ARG:HH12	1:A:749:LYS:HZ1	1.63	0.45
1:A:635:ARG:CG	1:A:635:ARG:NH1	2.60	0.45
1:A:673:LYS:HD2	1:A:674:SER:H	1.82	0.45
1:A:640:GLU:HG2	1:A:797:LEU:HD22	1.97	0.43
1:A:527:TYR:HB3	1:A:537:PHE:CE1	2.53	0.43
1:A:673:LYS:HD3	1:A:673:LYS:HA	1.55	0.42
1:A:666:PRO:O	1:A:702:LYS:HE3	2.19	0.42
1:A:688:ILE:HD11	1:A:708:VAL:HA	2.02	0.42
1:A:655:ASP:OD2	1:A:740:ARG:NH2	2.53	0.42
1:A:788:GLU:O	1:A:792:GLN:HG2	2.20	0.41
1:A:783:TYR:CE2	1:A:787:LYS:HE2	2.55	0.41
1:A:731:ASP:OD1	1:A:731:ASP:N	2.52	0.41
1:A:576:ASN:ND2	1:A:579:ARG:H	2.19	0.41
1:A:675:GLU:HG2	1:A:688:ILE:O	2.20	0.41
1:A:762:ARG:O	1:A:766:HIS:HB3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	244/251 (97%)	236 (97%)	7 (3%)	1 (0%)	34 57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	626	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	217/220 (99%)	195 (90%)	22 (10%)	7 14

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	528	GLU
1	A	530	GLU
1	A	546	VAL
1	A	558	LEU
1	A	565	ASN
1	A	567	LYS
1	A	576	ASN
1	A	586	VAL
1	A	587	GLN
1	A	588	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	635	ARG
1	A	636	VAL
1	A	637	VAL
1	A	673	LYS
1	A	682	ASP
1	A	709	THR
1	A	714	ASP
1	A	733	LYS
1	A	748	ARG
1	A	775	LEU
1	A	779	ASN
1	A	791	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	576	ASN
1	A	578	ASN
1	A	700	GLN
1	A	711	ASN
1	A	774	ASN
1	A	779	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.