



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 24, 2023 – 01:13 PM EST

PDB ID : 3LWF  
Title : Crystal structure of Putative transcriptional regulator (NP\_470886.1) from LISTERIA INNOCUA at 2.06 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2010-02-23  
Resolution : 2.06 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	
1	B	159	
1	C	159	
1	D	159	

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 5000 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative transcriptional regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	144	1142	715	200	223	4	0	4	0
1	B	144	1150	722	200	224	4	0	5	0
1	C	144	1134	711	197	222	4	0	3	0
1	D	144	1132	712	192	224	4	0	5	0

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	expression tag	UNP Q92BJ8
A	-17	GLY	-	expression tag	UNP Q92BJ8
A	-16	SER	-	expression tag	UNP Q92BJ8
A	-15	ASP	-	expression tag	UNP Q92BJ8
A	-14	LYS	-	expression tag	UNP Q92BJ8
A	-13	ILE	-	expression tag	UNP Q92BJ8
A	-12	HIS	-	expression tag	UNP Q92BJ8
A	-11	HIS	-	expression tag	UNP Q92BJ8
A	-10	HIS	-	expression tag	UNP Q92BJ8
A	-9	HIS	-	expression tag	UNP Q92BJ8
A	-8	HIS	-	expression tag	UNP Q92BJ8
A	-7	HIS	-	expression tag	UNP Q92BJ8
A	-6	GLU	-	expression tag	UNP Q92BJ8
A	-5	ASN	-	expression tag	UNP Q92BJ8
A	-4	LEU	-	expression tag	UNP Q92BJ8
A	-3	TYR	-	expression tag	UNP Q92BJ8
A	-2	PHE	-	expression tag	UNP Q92BJ8
A	-1	GLN	-	expression tag	UNP Q92BJ8
A	0	GLY	-	expression tag	UNP Q92BJ8
B	-18	MSE	-	expression tag	UNP Q92BJ8
B	-17	GLY	-	expression tag	UNP Q92BJ8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	expression tag	UNP Q92BJ8
B	-15	ASP	-	expression tag	UNP Q92BJ8
B	-14	LYS	-	expression tag	UNP Q92BJ8
B	-13	ILE	-	expression tag	UNP Q92BJ8
B	-12	HIS	-	expression tag	UNP Q92BJ8
B	-11	HIS	-	expression tag	UNP Q92BJ8
B	-10	HIS	-	expression tag	UNP Q92BJ8
B	-9	HIS	-	expression tag	UNP Q92BJ8
B	-8	HIS	-	expression tag	UNP Q92BJ8
B	-7	HIS	-	expression tag	UNP Q92BJ8
B	-6	GLU	-	expression tag	UNP Q92BJ8
B	-5	ASN	-	expression tag	UNP Q92BJ8
B	-4	LEU	-	expression tag	UNP Q92BJ8
B	-3	TYR	-	expression tag	UNP Q92BJ8
B	-2	PHE	-	expression tag	UNP Q92BJ8
B	-1	GLN	-	expression tag	UNP Q92BJ8
B	0	GLY	-	expression tag	UNP Q92BJ8
C	-18	MSE	-	expression tag	UNP Q92BJ8
C	-17	GLY	-	expression tag	UNP Q92BJ8
C	-16	SER	-	expression tag	UNP Q92BJ8
C	-15	ASP	-	expression tag	UNP Q92BJ8
C	-14	LYS	-	expression tag	UNP Q92BJ8
C	-13	ILE	-	expression tag	UNP Q92BJ8
C	-12	HIS	-	expression tag	UNP Q92BJ8
C	-11	HIS	-	expression tag	UNP Q92BJ8
C	-10	HIS	-	expression tag	UNP Q92BJ8
C	-9	HIS	-	expression tag	UNP Q92BJ8
C	-8	HIS	-	expression tag	UNP Q92BJ8
C	-7	HIS	-	expression tag	UNP Q92BJ8
C	-6	GLU	-	expression tag	UNP Q92BJ8
C	-5	ASN	-	expression tag	UNP Q92BJ8
C	-4	LEU	-	expression tag	UNP Q92BJ8
C	-3	TYR	-	expression tag	UNP Q92BJ8
C	-2	PHE	-	expression tag	UNP Q92BJ8
C	-1	GLN	-	expression tag	UNP Q92BJ8
C	0	GLY	-	expression tag	UNP Q92BJ8
D	-18	MSE	-	expression tag	UNP Q92BJ8
D	-17	GLY	-	expression tag	UNP Q92BJ8
D	-16	SER	-	expression tag	UNP Q92BJ8
D	-15	ASP	-	expression tag	UNP Q92BJ8
D	-14	LYS	-	expression tag	UNP Q92BJ8
D	-13	ILE	-	expression tag	UNP Q92BJ8

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-12	HIS	-	expression tag	UNP Q92BJ8
D	-11	HIS	-	expression tag	UNP Q92BJ8
D	-10	HIS	-	expression tag	UNP Q92BJ8
D	-9	HIS	-	expression tag	UNP Q92BJ8
D	-8	HIS	-	expression tag	UNP Q92BJ8
D	-7	HIS	-	expression tag	UNP Q92BJ8
D	-6	GLU	-	expression tag	UNP Q92BJ8
D	-5	ASN	-	expression tag	UNP Q92BJ8
D	-4	LEU	-	expression tag	UNP Q92BJ8
D	-3	TYR	-	expression tag	UNP Q92BJ8
D	-2	PHE	-	expression tag	UNP Q92BJ8
D	-1	GLN	-	expression tag	UNP Q92BJ8
D	0	GLY	-	expression tag	UNP Q92BJ8

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 10 8 2	0	1
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	1
			10	8	2		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	2	Total	Cl	0	0
			2	2		
3	C	3	Total	Cl	0	0
			3	3		
3	D	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	84	Total O 84 84	0	1
4	B	82	Total O 82 82	0	1
4	C	68	Total O 68 68	0	0
4	D	96	Total O 96 96	0	2


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative transcriptional regulator




- Molecule 1: Putative transcriptional regulator



- Molecule 1: Putative transcriptional regulator



- Molecule 1: Putative transcriptional regulator





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	108.96Å 115.78Å 75.93Å 90.00° 120.70° 90.00°	Depositor
Resolution (Å)	72.83 – 2.06 41.17 – 2.06	Depositor EDS
% Data completeness (in resolution range)	99.8 (72.83-2.06) 99.8 (41.17-2.06)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 2.06Å)	Xtrriage
Refinement program	REFMAC 5.5.0102, PHENIX	Depositor
R, $R_{free}$	0.180 , 0.218 0.185 , 0.223	Depositor DCC
$R_{free}$ test set	2529 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.8	Xtrriage
Anisotropy	0.559	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 55.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5000	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	0/1165	0.76	0/1564
1	B	0.70	0/1176	0.74	0/1577
1	C	0.67	0/1154	0.73	0/1549
1	D	0.74	0/1158	0.77	0/1555
All	All	0.70	0/4653	0.75	0/6245

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	69	GLY	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1142	0	1158	10	0
1	B	1150	0	1170	12	0
1	C	1134	0	1147	5	0
1	D	1132	0	1142	6	0
2	A	45	0	0	1	0
2	B	20	0	0	1	0
2	C	25	0	0	0	0
2	D	15	0	0	0	0
3	A	1	0	0	1	0
3	B	2	0	0	1	0
3	C	3	0	0	1	0
3	D	1	0	0	0	0
4	A	84	0	0	0	0
4	B	82	0	0	3	0
4	C	68	0	0	1	0
4	D	96	0	0	0	1
All	All	5000	0	4617	29	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:93:MSE:HE3	1:D:95:ASP:OD1	1.79	0.81
1:B:50[A]:ARG:NH2	2:B:152:SO4:O3	2.27	0.67
1:C:16[A]:GLU:HG2	4:C:699:HOH:O	1.94	0.67
1:A:1:MSE:HE3	1:A:3:ILE:HD11	1.79	0.65
1:B:93:MSE:HE3	1:B:95:ASP:OD1	1.98	0.64

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:780:HOH:O	4:D:811:HOH:O[2_456]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	146/159 (92%)	144 (99%)	2 (1%)	0	100	100
1	B	147/159 (92%)	146 (99%)	1 (1%)	0	100	100
1	C	145/159 (91%)	143 (99%)	2 (1%)	0	100	100
1	D	147/159 (92%)	145 (99%)	2 (1%)	0	100	100
All	All	585/636 (92%)	578 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/131 (95%)	124 (100%)	0	100	100
1	B	124/131 (95%)	123 (99%)	1 (1%)	81	81
1	C	122/131 (93%)	121 (99%)	1 (1%)	81	81
1	D	122/131 (93%)	122 (100%)	0	100	100
All	All	492/524 (94%)	490 (100%)	2 (0%)	91	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	100	GLN
1	C	100	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	40	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 7 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	C	156[A]	-	4,4,4	0.16	0	6,6,6	0.68	0
2	SO4	A	146	-	4,4,4	0.17	0	6,6,6	0.30	0
2	SO4	A	142[A]	-	4,4,4	0.15	0	6,6,6	0.60	0
2	SO4	B	150	-	4,4,4	0.39	0	6,6,6	0.67	0
2	SO4	C	157	-	4,4,4	0.22	0	6,6,6	0.37	0
2	SO4	C	156[B]	-	4,4,4	0.23	0	6,6,6	0.51	0
2	SO4	A	142[B]	-	4,4,4	0.22	0	6,6,6	0.78	0
2	SO4	A	145	-	4,4,4	0.18	0	6,6,6	0.41	0
2	SO4	A	143	-	4,4,4	0.23	0	6,6,6	0.45	0
2	SO4	B	151	-	4,4,4	0.21	0	6,6,6	0.33	0
2	SO4	D	163	-	4,4,4	0.18	0	6,6,6	0.12	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	147	-	4,4,4	0.21	0	6,6,6	0.30	0
2	SO4	A	148	-	4,4,4	0.35	0	6,6,6	0.53	0
2	SO4	D	165	-	4,4,4	0.12	0	6,6,6	0.38	0
2	SO4	A	141	-	4,4,4	0.13	0	6,6,6	0.17	0
2	SO4	C	159	-	4,4,4	0.12	0	6,6,6	0.19	0
2	SO4	B	152	-	4,4,4	0.15	0	6,6,6	0.26	0
2	SO4	C	158	-	4,4,4	0.16	0	6,6,6	0.17	0
2	SO4	D	164	-	4,4,4	0.15	0	6,6,6	0.64	0
2	SO4	A	144	-	4,4,4	0.21	0	6,6,6	0.33	0
2	SO4	B	153	-	4,4,4	0.09	0	6,6,6	0.30	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	143	SO4	1	0
2	B	152	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	140/159 (88%)	-0.34	2 (1%) 75 76	19, 25, 46, 75	0
1	B	140/159 (88%)	-0.20	4 (2%) 51 54	18, 26, 46, 73	0
1	C	140/159 (88%)	-0.27	4 (2%) 51 54	19, 26, 45, 70	0
1	D	140/159 (88%)	-0.25	6 (4%) 35 36	18, 25, 45, 70	0
All	All	560/636 (88%)	-0.26	16 (2%) 51 54	18, 25, 46, 75	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	-2	PHE	6.6
1	D	-2	PHE	6.4
1	B	-3	TYR	6.4
1	A	-3	TYR	6.2
1	A	-2	PHE	6.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	C	158	5/5	0.70	0.27	58,62,67,68	5
2	SO4	B	153	5/5	0.72	0.28	48,55,59,61	5
2	SO4	A	148	5/5	0.79	0.25	31,39,47,54	5
3	CL	C	162	1/1	0.82	0.10	70,70,70,70	0
3	CL	B	155	1/1	0.83	0.10	72,72,72,72	0
2	SO4	A	147	5/5	0.83	0.26	47,49,56,57	5
2	SO4	A	143	5/5	0.84	0.18	62,64,70,74	5
2	SO4	A	146	5/5	0.84	0.26	55,55,58,58	5
2	SO4	D	164	5/5	0.85	0.25	36,38,43,50	5
2	SO4	D	165	5/5	0.86	0.22	42,43,50,53	5
2	SO4	D	163	5/5	0.86	0.23	43,47,49,54	5
2	SO4	C	159	5/5	0.86	0.35	50,51,54,56	5
2	SO4	A	145	5/5	0.89	0.17	30,45,53,54	5
2	SO4	A	142[A]	5/5	0.93	0.13	22,27,36,39	5
2	SO4	A	144	5/5	0.93	0.13	22,40,43,49	5
3	CL	A	149	1/1	0.93	0.10	63,63,63,63	0
3	CL	B	154	1/1	0.93	0.07	58,58,58,58	0
2	SO4	A	142[B]	5/5	0.93	0.13	20,22,33,33	5
3	CL	C	160	1/1	0.93	0.12	61,61,61,61	0
2	SO4	B	151	5/5	0.93	0.15	26,36,42,50	5
2	SO4	B	152	5/5	0.94	0.18	51,57,58,59	5
2	SO4	C	156[A]	5/5	0.95	0.17	22,22,27,29	5
2	SO4	C	156[B]	5/5	0.95	0.17	21,24,37,41	5
2	SO4	B	150	5/5	0.95	0.10	40,40,55,56	0
3	CL	D	166	1/1	0.96	0.05	59,59,59,59	0
2	SO4	C	157	5/5	0.97	0.13	32,34,40,41	5
3	CL	C	161	1/1	0.97	0.11	50,50,50,50	0
2	SO4	A	141	5/5	0.98	0.08	52,55,57,58	5

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.