



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2024 – 09:09 PM EST

PDB ID : 3N4Q  
Title : Human cytomegalovirus terminase nuclease domain, Mn soaked  
Authors : Nadal, M.; Mas, P.J.; Blanco, A.G.; Arnan, C.; Sola, M.; Hart, D.J.; Coll, M.  
Deposited on : 2010-05-22  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	
1	B	279	
1	C	279	
1	D	279	

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7056 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TERMINASE SUBUNIT UL89 PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	216	Total 1736	C 1113	N 292	O 325	S 6	0	0	0
1	B	221	Total 1775	C 1137	N 301	O 331	S 6	0	0	0
1	C	219	Total 1755	C 1126	N 293	O 330	S 6	0	0	0
1	D	214	Total 1722	C 1104	N 290	O 322	S 6	0	0	0

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	396	MET	-	expression tag	UNP P16732
A	397	GLY	-	expression tag	UNP P16732
A	398	HIS	-	expression tag	UNP P16732
A	399	HIS	-	expression tag	UNP P16732
A	400	HIS	-	expression tag	UNP P16732
A	401	HIS	-	expression tag	UNP P16732
A	402	HIS	-	expression tag	UNP P16732
A	403	HIS	-	expression tag	UNP P16732
A	404	ASP	-	expression tag	UNP P16732
A	405	TYR	-	expression tag	UNP P16732
A	406	ASP	-	expression tag	UNP P16732
A	407	ILE	-	expression tag	UNP P16732
A	408	PRO	-	expression tag	UNP P16732
A	409	THR	-	expression tag	UNP P16732
A	410	THR	-	expression tag	UNP P16732
A	411	GLU	-	expression tag	UNP P16732
A	412	ASN	-	expression tag	UNP P16732
A	413	LEU	-	expression tag	UNP P16732
A	414	TYR	-	expression tag	UNP P16732
A	415	PHE	-	expression tag	UNP P16732
A	416	GLN	-	expression tag	UNP P16732

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	417	GLY	-	expression tag	UNP P16732
B	396	MET	-	expression tag	UNP P16732
B	397	GLY	-	expression tag	UNP P16732
B	398	HIS	-	expression tag	UNP P16732
B	399	HIS	-	expression tag	UNP P16732
B	400	HIS	-	expression tag	UNP P16732
B	401	HIS	-	expression tag	UNP P16732
B	402	HIS	-	expression tag	UNP P16732
B	403	HIS	-	expression tag	UNP P16732
B	404	ASP	-	expression tag	UNP P16732
B	405	TYR	-	expression tag	UNP P16732
B	406	ASP	-	expression tag	UNP P16732
B	407	ILE	-	expression tag	UNP P16732
B	408	PRO	-	expression tag	UNP P16732
B	409	THR	-	expression tag	UNP P16732
B	410	THR	-	expression tag	UNP P16732
B	411	GLU	-	expression tag	UNP P16732
B	412	ASN	-	expression tag	UNP P16732
B	413	LEU	-	expression tag	UNP P16732
B	414	TYR	-	expression tag	UNP P16732
B	415	PHE	-	expression tag	UNP P16732
B	416	GLN	-	expression tag	UNP P16732
B	417	GLY	-	expression tag	UNP P16732
C	396	MET	-	expression tag	UNP P16732
C	397	GLY	-	expression tag	UNP P16732
C	398	HIS	-	expression tag	UNP P16732
C	399	HIS	-	expression tag	UNP P16732
C	400	HIS	-	expression tag	UNP P16732
C	401	HIS	-	expression tag	UNP P16732
C	402	HIS	-	expression tag	UNP P16732
C	403	HIS	-	expression tag	UNP P16732
C	404	ASP	-	expression tag	UNP P16732
C	405	TYR	-	expression tag	UNP P16732
C	406	ASP	-	expression tag	UNP P16732
C	407	ILE	-	expression tag	UNP P16732
C	408	PRO	-	expression tag	UNP P16732
C	409	THR	-	expression tag	UNP P16732
C	410	THR	-	expression tag	UNP P16732
C	411	GLU	-	expression tag	UNP P16732
C	412	ASN	-	expression tag	UNP P16732
C	413	LEU	-	expression tag	UNP P16732
C	414	TYR	-	expression tag	UNP P16732

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	415	PHE	-	expression tag	UNP P16732
C	416	GLN	-	expression tag	UNP P16732
C	417	GLY	-	expression tag	UNP P16732
D	396	MET	-	expression tag	UNP P16732
D	397	GLY	-	expression tag	UNP P16732
D	398	HIS	-	expression tag	UNP P16732
D	399	HIS	-	expression tag	UNP P16732
D	400	HIS	-	expression tag	UNP P16732
D	401	HIS	-	expression tag	UNP P16732
D	402	HIS	-	expression tag	UNP P16732
D	403	HIS	-	expression tag	UNP P16732
D	404	ASP	-	expression tag	UNP P16732
D	405	TYR	-	expression tag	UNP P16732
D	406	ASP	-	expression tag	UNP P16732
D	407	ILE	-	expression tag	UNP P16732
D	408	PRO	-	expression tag	UNP P16732
D	409	THR	-	expression tag	UNP P16732
D	410	THR	-	expression tag	UNP P16732
D	411	GLU	-	expression tag	UNP P16732
D	412	ASN	-	expression tag	UNP P16732
D	413	LEU	-	expression tag	UNP P16732
D	414	TYR	-	expression tag	UNP P16732
D	415	PHE	-	expression tag	UNP P16732
D	416	GLN	-	expression tag	UNP P16732
D	417	GLY	-	expression tag	UNP P16732

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mn 2 2	0	0
2	B	2	Total Mn 2 2	0	0
2	C	2	Total Mn 2 2	0	0
2	D	2	Total Mn 2 2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	11	Total O 11 11	0	0
4	B	17	Total O 17 17	0	0
4	C	15	Total O 15 15	0	0
4	D	13	Total O 13 13	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: TERMINASE SUBUNIT UL89 PROTEIN




#### • Molecule 1: TERMINASE SUBUNIT UL89 PROTEIN





#### • Molecule 1: TERMINASE SUBUNIT UL89 PROTEIN





● Molecule 1: TERMINASE SUBUNIT UL89 PROTEIN





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.70Å 88.10Å 190.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.05 – 3.20 44.05 – 3.19	Depositor EDS
% Data completeness (in resolution range)	89.6 (44.05-3.20) 89.6 (44.05-3.19)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.90 (at 3.19Å)	Xtrriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.207 , 0.283 0.200 , 0.271	Depositor DCC
$R_{free}$ test set	1109 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	75.1	Xtrriage
Anisotropy	0.820	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 68.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.046 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7056	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	84.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.61	0/1770	0.75	1/2399 (0.0%)
1	B	0.58	1/1809 (0.1%)	0.74	2/2452 (0.1%)
1	C	0.58	0/1789	0.74	3/2427 (0.1%)
1	D	0.55	0/1756	0.69	0/2380
All	All	0.58	1/7124 (0.0%)	0.73	6/9658 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	622	GLU	CG-CD	5.10	1.59	1.51

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	488	LEU	CA-CB-CG	7.49	132.52	115.30
1	C	458	LEU	CA-CB-CG	6.16	129.46	115.30
1	B	605	LEU	CA-CB-CG	5.69	128.38	115.30
1	C	569	GLN	N-CA-C	-5.46	96.27	111.00
1	B	578	LEU	CA-CB-CG	5.35	127.60	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1736	0	1707	64	0
1	B	1775	0	1751	67	0
1	C	1755	0	1732	70	0
1	D	1722	0	1691	59	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	11	0	0	0	0
4	B	17	0	0	2	0
4	C	15	0	0	1	0
4	D	13	0	0	0	0
All	All	7056	0	6881	254	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 254 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:664:ASP:HB2	4:B:18:HOH:O	1.55	1.03
1:D:464:PRO:HA	1:D:475:THR:HB	1.44	1.00
1:D:611:LYS:HB3	1:D:612:LEU:HD22	1.45	0.99
1:A:456:LYS:HD3	1:A:525:TYR:O	1.73	0.90
1:B:568:ASP:CG	1:B:569:GLN:H	1.77	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/279 (75%)	188 (90%)	15 (7%)	5 (2%)	6	34
1	B	213/279 (76%)	182 (85%)	29 (14%)	2 (1%)	17	56
1	C	213/279 (76%)	179 (84%)	30 (14%)	4 (2%)	8	39
1	D	206/279 (74%)	178 (86%)	26 (13%)	2 (1%)	15	54
All	All	840/1116 (75%)	727 (86%)	100 (12%)	13 (2%)	10	44

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	612	LEU
1	C	569	GLN
1	D	504	SER
1	A	537	THR
1	A	569	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/242 (78%)	154 (82%)	34 (18%)	1	8
1	B	192/242 (79%)	160 (83%)	32 (17%)	2	10
1	C	190/242 (78%)	154 (81%)	36 (19%)	1	8
1	D	186/242 (77%)	152 (82%)	34 (18%)	1	8
All	All	756/968 (78%)	620 (82%)	136 (18%)	1	9

5 of 136 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	521	SER
1	D	539	GLN
1	D	621	LEU
1	B	539	GLN
1	B	537	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	625	GLN
1	D	536	ASN
1	D	514	HIS
1	D	538	ASN
1	B	539	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	216/279 (77%)	-0.34	4 (1%) 66 53	48, 77, 119, 164	6 (2%)
1	B	221/279 (79%)	-0.41	0 100 100	53, 76, 110, 154	2 (0%)
1	C	219/279 (78%)	-0.34	2 (0%) 84 75	54, 82, 128, 180	4 (1%)
1	D	214/279 (76%)	-0.24	2 (0%) 84 75	57, 86, 122, 148	12 (5%)
All	All	870/1116 (77%)	-0.33	8 (0%) 84 75	48, 80, 123, 180	24 (2%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	612	LEU	3.2
1	D	613	SER	3.0
1	D	495	PHE	2.9
1	A	497	LEU	2.9
1	A	503	SER	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	B	2	1/1	0.88	0.10	40,40,40,40	0
3	MG	D	3	1/1	0.92	0.05	38,38,38,38	0
3	MG	C	4	1/1	0.93	0.17	65,65,65,65	0
3	MG	A	675	1/1	0.95	0.08	29,29,29,29	0
2	MN	A	1	1/1	0.96	0.12	58,58,58,58	0
2	MN	A	2	1/1	0.96	0.14	55,55,55,55	0
2	MN	B	4	1/1	0.96	0.07	52,52,52,52	0
2	MN	C	5	1/1	0.96	0.07	58,58,58,58	0
2	MN	D	7	1/1	0.97	0.09	60,60,60,60	0
2	MN	B	3	1/1	0.98	0.05	55,55,55,55	0
2	MN	D	8	1/1	0.98	0.18	64,64,64,64	0
2	MN	C	6	1/1	0.98	0.06	58,58,58,58	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.