



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 07:47 pm BST

PDB ID : 3NCY
Title : X-ray crystal structure of an arginine agmatine antiporter (AdiC) in complex with a Fab fragment
Authors : Fang, Y.; Jayaram, H.; Shane, T.; Komalkova-Partensky, L.; Wu, F.; Williams, C.; Xiong, Y.; Miller, C.
Deposited on : 2010-06-06
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	
1	B	445	
1	C	445	
1	D	445	
2	P	219	
2	Q	219	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	S	211	
3	W	211	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 18693 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AdiC.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	422	Total 3052	C 2018	N 489	O 524	S 21	0	0	0
1	B	420	Total 3039	C 2009	N 489	O 520	S 21	5	0	0
1	C	420	Total 3034	C 2006	N 487	O 520	S 21	0	0	0
1	D	419	Total 3031	C 2005	N 486	O 519	S 21	17	0	0

- Molecule 2 is a protein called Fab Heavy chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	Q	218	Total 1640	C 1044	N 264	O 325	S 7	16	0	0
2	P	219	Total 1647	C 1049	N 265	O 326	S 7	15	0	0

- Molecule 3 is a protein called Fab Light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	W	211	Total 1625	C 1011	N 271	O 333	S 10	40	0	0
3	S	211	Total 1625	C 1011	N 271	O 333	S 10	20	0	0




• Molecule 1: AdiC




• Molecule 1: AdiC







• Molecule 2: Fab Heavy chain




• Molecule 2: Fab Heavy chain





• Molecule 3: Fab Light chain





- Molecule 3: Fab Light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	79.66Å 104.15Å 154.03Å 81.96° 75.93° 73.73°	Depositor
Resolution (Å)	33.23 – 3.20 45.88 – 3.20	Depositor EDS
% Data completeness (in resolution range)	96.4 (33.23-3.20) 96.4 (45.88-3.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 3.19Å)	Xtrriage
Refinement program	PHENIX 1.6.1_357, REFMAC	Depositor
R, R_{free}	0.282 , 0.312 0.275 , 0.302	Depositor DCC
R_{free} test set	3612 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtrriage
Anisotropy	0.186	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.20 , 84.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.79	EDS
Total number of atoms	18693	wwPDB-VP
Average B, all atoms (Å ²)	115.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/3125	0.42	0/4281
1	B	0.22	0/3110	0.43	0/4257
1	C	0.22	0/3107	0.43	0/4256
1	D	0.22	0/3103	0.42	0/4250
2	P	0.22	0/1694	0.46	1/2319 (0.0%)
2	Q	0.23	0/1686	0.46	0/2307
3	S	0.23	0/1665	0.42	0/2260
3	W	0.23	0/1665	0.43	0/2260
All	All	0.22	0/19155	0.43	1/26190 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	134	VAL	N-CA-C	-5.71	95.58	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3052	0	3120	201	0
1	B	3039	0	3112	191	0
1	C	3034	0	3098	190	0
1	D	3031	0	3102	199	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	P	1647	0	1601	128	0
2	Q	1640	0	1594	135	0
3	S	1625	0	1550	83	0
3	W	1625	0	1550	85	0
All	All	18693	0	18727	1142	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 1142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:346:ALA:HA	1:A:347:ALA:CB	1.58	1.24
1:A:346:ALA:CA	1:A:347:ALA:HB3	1.76	1.12
1:A:23:ILE:HG12	1:A:206:GLY:HA3	1.33	1.11
1:D:356:VAL:HG11	1:D:409:GLU:HB3	1.30	1.10
1:B:23:ILE:HG12	1:B:206:GLY:HA3	1.32	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	416/445 (94%)	328 (79%)	59 (14%)	29 (7%)	1 8
1	B	414/445 (93%)	328 (79%)	57 (14%)	29 (7%)	1 8
1	C	414/445 (93%)	322 (78%)	68 (16%)	24 (6%)	1 13
1	D	413/445 (93%)	319 (77%)	71 (17%)	23 (6%)	2 14
2	P	217/219 (99%)	179 (82%)	27 (12%)	11 (5%)	2 15
2	Q	216/219 (99%)	180 (83%)	25 (12%)	11 (5%)	2 15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	S	209/211 (99%)	174 (83%)	28 (13%)	7 (3%)	4	25
3	W	209/211 (99%)	173 (83%)	29 (14%)	7 (3%)	4	25
All	All	2508/2640 (95%)	2003 (80%)	364 (14%)	141 (6%)	2	14

5 of 141 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	116	PHE
1	A	118	ILE
1	A	166	ALA
1	A	270	LEU
1	A	274	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/343 (90%)	280 (91%)	28 (9%)	9	34
1	B	306/343 (89%)	277 (90%)	29 (10%)	8	32
1	C	305/343 (89%)	286 (94%)	19 (6%)	18	53
1	D	306/343 (89%)	286 (94%)	20 (6%)	17	51
2	P	187/187 (100%)	166 (89%)	21 (11%)	6	25
2	Q	186/187 (100%)	162 (87%)	24 (13%)	4	19
3	S	185/185 (100%)	169 (91%)	16 (9%)	10	38
3	W	185/185 (100%)	168 (91%)	17 (9%)	9	33
All	All	1968/2116 (93%)	1794 (91%)	174 (9%)	10	36

5 of 174 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	35	LEU
2	Q	6	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	S	79	ILE
1	D	114	TYR
1	D	316	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	299	GLN
1	C	434	ASN
3	W	94	GLN
1	B	377	HIS
3	W	30	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	422/445 (94%)	0.75	59 (13%) 2 1	56, 127, 231, 339	0
1	B	419/445 (94%)	0.45	37 (8%) 10 5	62, 126, 220, 332	0
1	C	420/445 (94%)	0.39	31 (7%) 14 8	37, 105, 203, 283	0
1	D	416/445 (93%)	0.59	43 (10%) 6 4	33, 105, 197, 283	0
2	P	218/219 (99%)	0.50	15 (6%) 16 9	48, 104, 168, 282	2 (0%)
2	Q	216/219 (98%)	0.74	27 (12%) 3 2	44, 108, 171, 278	2 (0%)
3	S	209/211 (99%)	0.39	9 (4%) 35 22	30, 80, 169, 258	0
3	W	207/211 (98%)	0.62	28 (13%) 3 2	23, 81, 166, 266	0
All	All	2527/2640 (95%)	0.55	249 (9%) 7 4	23, 109, 203, 339	4 (0%)

The worst 5 of 249 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	Q	135	CYS	22.3
1	D	250	ILE	13.0
2	Q	137	ASP	11.5
1	B	264	ASP	8.5
1	A	268	MET	7.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.