

wwPDB X-ray Structure Validation Summary Report (i)

Sep 18, 2023 – 02:33 PM EDT

PDB ID : 3PCD

Title: PROTOCATECHUATE 3,4-DIOXYGENASE Y447H MUTANT

Authors: Orville, A.M.; Lipscomb, J.D.; Ohlendorf, D.H.

Deposited on : 1997-11-24

Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain	
1	A	200	80% 15%	•
1	В	200	78% 18%	•
1	С	200	77% 18%	•
1	D	200	78% 18%	
1	E	200	72% 23%	5%
1	F	200	72% 22%	6%
2	M	238		
			73% 21%	
2	N	238	78% 17%	• •



Continued from previous page...

Mol	Chain	Length	Quality of chain		
2	О	238	75%	18%	
2	Р	238	72%	20%	• • •
2	Q	238	76%	18%	• • •
2	R	238	72%	22%	



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 21930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called PROTOCATECHUATE 3,4-DIOXYGENASE.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	200	Total	С	N	О	S	0	0	0
1	Λ	200	1571	993	276	299	3	0	0	U
1	В	200	Total	С	N	О	S	0	0	0
1	Ъ	200	1571	993	276	299	3	0	0	U
1	С	200	Total	С	N	О	S	0	0	0
1		200	1571	993	276	299	3	0		U
1	D	200	Total	С	N	О	S	0	0	0
1	D	200	1571	993	276	299	3	0	0	0
1	E	200	Total	С	N	Ο	S	0	0	0
1	ш	200	1571	993	276	299	3	0	U	U
1	F	200	Total	С	N	О	S	0	0	0
1	I.	200	1571	993	276	299	3	U	U	U

• Molecule 2 is a protein called PROTOCATECHUATE 3,4-DIOXYGENASE.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
2	M	233	Total	С	N	О	S	0	0	0
2	101	255	1838	1168	336	327	7	0	U	
2	N	233	Total	С	N	О	S	0	0	0
2	11	200	1838	1168	336	327	7	U	U	U
2	0	233	Total	С	N	O	S	0	0	0
2		200	1838	1168	336	327	7	U	U	
2	Р	233	Total	С	N	O	S	0	0	0
2	1	200	1838	1168	336	327	7	0	U	U
2	Q	233	Total	С	N	O	S	0	0	0
2	Q .	200	1838	1168	336	327	7	U	U	U
2	R	233	Total	С	N	О	S	0	0	0
	16	200	1838	1168	336	327	7	0	U	U

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	447	HIS	TYR	engineered mutation	UNP P00437






 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
N	447	HIS	TYR	engineered mutation	UNP P00437
О	447	HIS	TYR	engineered mutation	UNP P00437
Р	447	HIS	TYR	engineered mutation	UNP P00437
Q	447	HIS	TYR	engineered mutation	UNP P00437
R	447	HIS	TYR	engineered mutation	UNP P00437

• Molecule 3 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	M	1	Total C O 4 1 3	0	0
3	N	1	Total C O 4 1 3	0	0
3	О	1	Total C O 4 1 3	0	0
3	Р	1	Total C O 4 1 3	0	0
3	Q	1	Total C O 4 1 3	0	0
3	R	1	Total C O 4 1 3	0	0

• Molecule 4 is FE (III) ION (three-letter code: FE) (formula: Fe).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	M	1	Total Fe 1 1	0	0
4	N	1	Total Fe 1 1	0	0
4	О	1	Total Fe 1 1	0	0
4	Р	1	Total Fe 1 1	0	0
4	Q	1	Total Fe 1 1	0	0
4	R	1	Total Fe 1 1	0	0

 \bullet Molecule 5 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: $\mathrm{C_2H_6OS}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	M	1	Total C O S 4 2 1 1	0	0
5	N	1	Total C O S 4 2 1 1	0	0
5	О	1	Total C O S 4 2 1 1	0	0
5	Р	1	Total C O S 4 2 1 1	0	0
5	Q	1	Total C O S 4 2 1 1	0	0
5	R	1	Total C O S 4 2 1 1	0	0



• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	73	Total O 73 73	0	0
6	M	161	Total O 161 161	0	0
6	В	79	Total O 79 79	0	0
6	N	163	Total O 163 163	0	0
6	С	78	Total O 78 78	0	0
6	О	156	Total O 156 156	0	0
6	D	81	Total O 81 81	0	0
6	Р	151	Total O 151 151	0	0
6	Е	82	Total O 82 82	0	0
6	Q	159	Total O 159 159	0	0
6	F	80	Total O 80 80	0	0
6	R	159	Total O 159 159	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: PROTOCATECHUATE 3,4-DIOXYGENASE




• Molecule 1: PROTOCATECHUATE 3,4-DIOXYGENASE



• Molecule 1: PROTOCATECHUATE 3,4-DIOXYGENASE




• Molecule 1: PROTOCATECHUATE 3,4-DIOXYGENASE









• Molecule 1: PROTOCATECHUATE 3,4-DIOXYGENASE




• Molecule 1: PROTOCATECHUATE 3,4-DIOXYGENASE




• Molecule 2: PROTOCATECHUATE 3,4-DIOXYGENASE




• Molecule 2: PROTOCATECHUATE 3,4-DIOXYGENASE




• Molecule 2: PROTOCATECHUATE 3,4-DIOXYGENASE







• Molecule 2: PROTOCATECHUATE 3,4-DIOXYGENASE



• Molecule 2: PROTOCATECHUATE 3,4-DIOXYGENASE



• Molecule 2: PROTOCATECHUATE 3,4-DIOXYGENASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	I 1 2 1	Depositor	
Cell constants	197.40Å 127.20Å 134.60Å	Depositor	
a, b, c, α , β , γ	90.00° 97.70° 90.00°	Depositor	
Resolution (Å)	6.00 - 2.10	Depositor	
% Data completeness	78.0 (6.00-2.10)	Depositor	
(in resolution range)	10.0 (0.00 2.10)	Depositor	
R_{merge}	(Not available)	Depositor	
R_{sym}	0.11	Depositor	
Refinement program	PROLSQ	Depositor	
R, R_{free}	0.180 , (Not available)	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	21930	wwPDB-VP	
Average B, all atoms (Å ²)	19.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, BME, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	ond lengths	E	Bond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5
1	A	1.05	1/1611 (0.1%)	1.70	30/2195 (1.4%)
1	В	1.06	2/1611 (0.1%)	1.79	30/2195 (1.4%)
1	С	1.04	0/1611	1.71	26/2195 (1.2%)
1	D	1.06	2/1611 (0.1%)	1.78	28/2195 (1.3%)
1	Е	1.01	1/1611 (0.1%)	1.60	28/2195 (1.3%)
1	F	1.03	0/1611	1.65	25/2195 (1.1%)
2	M	1.12	1/1893 (0.1%)	1.67	35/2577 (1.4%)
2	N	1.08	2/1893~(0.1%)	1.63	36/2577 (1.4%)
2	O	1.11	2/1893 (0.1%)	1.56	32/2577 (1.2%)
2	Р	1.12	1/1893 (0.1%)	1.75	34/2577 (1.3%)
2	Q	1.14	3/1893 (0.2%)	1.70	27/2577 (1.0%)
2	R	1.14	2/1893 (0.1%)	1.62	30/2577 (1.2%)
All	All	1.08	17/21024 (0.1%)	1.68	361/28632 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1
2	Р	0	1
All	All	0	2

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
1	D	94	ARG	CD-NE	-7.01	1.34	1.46
1	A	94	ARG	CD-NE	-6.94	1.34	1.46
2	R	452	GLY	N-CA	6.50	1.55	1.46
2	Q	466	SER	CB-OG	6.26	1.50	1.42



Continued from previous page...

Mol	Chain	Res	Type	Atoms	${f Z}$	$\operatorname{Observed}(\text{\AA})$	Ideal(A)	
2	Q	440	ARG	CD-NE	-6.24	1.35	1.46	

The worst 5 of 361 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
2	Р	440	ARG	NE-CZ-NH2	-31.69	104.46	120.30
1	D	133	ARG	CD-NE-CZ	30.52	166.32	123.60
1	В	133	ARG	CD-NE-CZ	29.97	165.56	123.60
2	N	440	ARG	NE-CZ-NH2	-22.92	108.84	120.30
2	Р	440	ARG	NE-CZ-NH1	22.20	131.40	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	184	ARG	Sidechain
2	Р	440	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1571	0	1499	28	0
1	В	1571	0	1499	26	0
1	С	1571	0	1499	33	0
1	D	1571	0	1499	23	0
1	Е	1571	0	1499	42	0
1	F	1571	0	1499	44	0
2	M	1838	0	1791	44	0
2	N	1838	0	1791	21	0
2	О	1838	0	1791	31	0
2	Р	1838	0	1791	42	0
2	Q	1838	0	1791	35	0
2	R	1838	0	1791	37	0
3	M	4	0	0	1	0
3	N	4	0	0	0	0
3	О	4	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H		H(added)	Clashes	Symm-Clashes
3	Р	4	0	0	1	0
3	Q	4	0	0	0	0
3	R	4	0	0	1	0
4	M	1	0	0	0	0
4	N	1	0	0	0	0
4	О	1	0	0	0	0
4	Р	1	0	0	0	0
4	Q	1	0	0	0	0
4	R	1	0	0	0	0
5	M	4	0	5	0	0
5	N	4	0	5	0	0
5	O	4	0	5	1	0
5	Р	4	0	5	0	0
5	Q	4	0	5	2	0
5	R	4	0	5	0	0
6	A	73	0	0	0	0
6	В	79	0	0	0	0
6	С	78	0	0	0	0
6	D	81	0	0	0	0
6	Е	82	0	0	1	0
6	F	80	0	0	0	0
6	M	161	0	0	4	0
6	N	163	0	0	2	0
6	O	156	0	0	2	0
6	Р	151	0	0	3	0
6	Q	159	0	0	2	0
6	R	159	0	0	2	0
All	All	21930	0	19770	377	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 377 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:F:165:GLN:H	1:F:165:GLN:HE21	0.97	0.93
1:E:165:GLN:H	1:E:165:GLN:NE2	1.64	0.93
1:F:165:GLN:H	1:F:165:GLN:NE2	1.66	0.93
1:E:165:GLN:H	1:E:165:GLN:HE21	1.05	0.92
2:M:497:ASN:HD22	2:M:499:GLU:H	1.22	0.87



There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	198/200 (99%)	190 (96%)	8 (4%)	0	100	100
1	В	198/200 (99%)	193 (98%)	5 (2%)	0	100	100
1	C	198/200 (99%)	193 (98%)	5 (2%)	0	100	100
1	D	198/200 (99%)	191 (96%)	7 (4%)	0	100	100
1	E	198/200 (99%)	189 (96%)	9 (4%)	0	100	100
1	F	198/200 (99%)	191 (96%)	7 (4%)	0	100	100
2	M	229/238 (96%)	222 (97%)	7 (3%)	0	100	100
2	N	229/238 (96%)	223 (97%)	6 (3%)	0	100	100
2	О	229/238 (96%)	221 (96%)	8 (4%)	0	100	100
2	Р	229/238~(96%)	224 (98%)	5 (2%)	0	100	100
2	Q	229/238 (96%)	221 (96%)	8 (4%)	0	100	100
2	R	229/238 (96%)	224 (98%)	5 (2%)	0	100	100
All	All	$2562/2628 \; (98\%)$	2482 (97%)	80 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	A	162/163~(99%)	156 (96%)	6 (4%)	34	35
1	В	162/163 (99%)	157 (97%)	5 (3%)	40	43
1	C	162/163 (99%)	156 (96%)	6 (4%)	34	35
1	D	162/163 (99%)	155 (96%)	7 (4%)	29	29
1	E	162/163~(99%)	154 (95%)	8 (5%)	25	23
1	F	162/163 (99%)	155 (96%)	7 (4%)	29	29
2	M	196/202 (97%)	185 (94%)	11 (6%)	21	18
2	N	196/202 (97%)	187 (95%)	9 (5%)	27	26
2	О	196/202 (97%)	188 (96%)	8 (4%)	30	31
2	Р	196/202 (97%)	183 (93%)	13 (7%)	16	14
2	Q	196/202 (97%)	185 (94%)	11 (6%)	21	18
2	R	196/202 (97%)	184 (94%)	12 (6%)	18	16
All	All	2148/2190 (98%)	2045 (95%)	103 (5%)	25	24

5 of 103 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
2	Р	416	LEU
1	Е	165	GLN
2	R	478	LEU
2	Р	434	ASP
1	Е	4	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
2	О	497	ASN
1	F	165	GLN
1	D	165	GLN
2	R	497	ASN
2	Q	497	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 6 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	В	ond leng	$_{ m gths}$	Bond angles		
MIOI	$egin{array}{c c} \operatorname{Mol} & \operatorname{Type} & \operatorname{Chain} & \end{array}$	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	CO3	О	550	4	2,3,3	0.61	0	2,3,3	1.43	0
5	BME	Q	601	2	3,3,3	0.57	0	1,2,2	1.15	0
3	CO3	R	550	4	2,3,3	0.63	0	2,3,3	1.77	1 (50%)
3	CO3	M	550	4	2,3,3	0.39	0	2,3,3	1.05	0
5	BME	Р	601	2	3,3,3	0.44	0	1,2,2	1.15	0
5	BME	N	601	2	3,3,3	0.25	0	1,2,2	0.51	0
3	CO3	P	550	4	2,3,3	0.46	0	2,3,3	0.69	0
5	BME	О	601	2	3,3,3	0.35	0	1,2,2	1.00	0
3	CO3	N	550	4	2,3,3	0.32	0	2,3,3	1.11	0
5	BME	R	601	2	3,3,3	0.30	0	1,2,2	0.07	0
3	CO3	Q	550	4	2,3,3	0.78	0	2,3,3	3.22	2 (100%)
5	BME	M	601	2	3,3,3	0.38	0	1,2,2	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BME	Q	601	2	-	0/1/1/1	-
5	BME	Р	601	2	-	0/1/1/1	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BME	N	601	2	-	0/1/1/1	-
5	BME	О	601	2	-	0/1/1/1	-
5	BME	R	601	2	-	0/1/1/1	-
5	BME	M	601	2	-	1/1/1/1	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
3	Q	550	CO3	O3-C-O1	-3.66	110.06	119.55
3	Q	550	CO3	O2-C-O1	-2.70	112.54	119.55
3	R	550	CO3	O3-C-O1	-2.43	113.25	119.55

There are no chirality outliers.

All (1) torsion outliers are listed below:

\mathbf{N}	Iol	Chain	Res	Type	Atoms
	5	M	601	BME	O1-C1-C2-S2

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	Q	601	BME	2	0
3	R	550	CO3	1	0
3	M	550	CO3	1	0
3	Р	550	CO3	1	0
5	О	601	BME	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

