



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 17, 2024 – 03:18 PM EST

PDB ID : 3SEY
Title : Zn-mediated Polymer of Maltose-binding Protein A216H/K220H by Synthetic Symmetrization (Form II)
Authors : Zhao, M.; Soriaga, A.B.; Laganowsky, A.; Sawaya, M.R.; Cascio, D.; Yeates, T.O.
Deposited on : 2011-06-11
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	
1	C	372	
1	E	372	
2	B	2	
2	D	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 50% 50%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9260 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	371	Total 2879	C 1852	N 470	O 550	S 7	0	4	0
1	C	371	Total 2879	C 1856	N 470	O 546	S 7	0	5	0
1	E	370	Total 2845	C 1831	N 465	O 543	S 6	0	1	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P0AEX9
A	216	HIS	ALA	engineered mutation	UNP P0AEX9
A	220	HIS	LYS	engineered mutation	UNP P0AEX9
A	360	ALA	-	expression tag	UNP P0AEX9
A	361	ALA	-	expression tag	UNP P0AEX9
A	362	LEU	-	expression tag	UNP P0AEX9
A	363	ALA	-	expression tag	UNP P0AEX9
A	364	ALA	-	expression tag	UNP P0AEX9
A	365	ALA	-	expression tag	UNP P0AEX9
A	366	GLN	-	expression tag	UNP P0AEX9
A	367	THR	-	expression tag	UNP P0AEX9
A	368	ASN	-	expression tag	UNP P0AEX9
A	369	ALA	-	expression tag	UNP P0AEX9
A	370	ALA	-	expression tag	UNP P0AEX9
A	371	ALA	-	expression tag	UNP P0AEX9
A	372	ALA	-	expression tag	UNP P0AEX9
C	1	MET	-	initiating methionine	UNP P0AEX9
C	216	HIS	ALA	engineered mutation	UNP P0AEX9
C	220	HIS	LYS	engineered mutation	UNP P0AEX9
C	360	ALA	-	expression tag	UNP P0AEX9
C	361	ALA	-	expression tag	UNP P0AEX9
C	362	LEU	-	expression tag	UNP P0AEX9
C	363	ALA	-	expression tag	UNP P0AEX9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	364	ALA	-	expression tag	UNP P0AEX9
C	365	ALA	-	expression tag	UNP P0AEX9
C	366	GLN	-	expression tag	UNP P0AEX9
C	367	THR	-	expression tag	UNP P0AEX9
C	368	ASN	-	expression tag	UNP P0AEX9
C	369	ALA	-	expression tag	UNP P0AEX9
C	370	ALA	-	expression tag	UNP P0AEX9
C	371	ALA	-	expression tag	UNP P0AEX9
C	372	ALA	-	expression tag	UNP P0AEX9
E	1	MET	-	initiating methionine	UNP P0AEX9
E	216	HIS	ALA	engineered mutation	UNP P0AEX9
E	220	HIS	LYS	engineered mutation	UNP P0AEX9
E	360	ALA	-	expression tag	UNP P0AEX9
E	361	ALA	-	expression tag	UNP P0AEX9
E	362	LEU	-	expression tag	UNP P0AEX9
E	363	ALA	-	expression tag	UNP P0AEX9
E	364	ALA	-	expression tag	UNP P0AEX9
E	365	ALA	-	expression tag	UNP P0AEX9
E	366	GLN	-	expression tag	UNP P0AEX9
E	367	THR	-	expression tag	UNP P0AEX9
E	368	ASN	-	expression tag	UNP P0AEX9
E	369	ALA	-	expression tag	UNP P0AEX9
E	370	ALA	-	expression tag	UNP P0AEX9
E	371	ALA	-	expression tag	UNP P0AEX9
E	372	ALA	-	expression tag	UNP P0AEX9

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	2	0
			23	12	11			
2	D	2	Total	C	O	0	2	0
			23	12	11			
2	F	2	Total	C	O	0	2	0
			46	24	22			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total	Zn	0	0
			10	10		
3	C	7	Total	Zn	0	0
			7	7		
3	E	6	Total	Zn	0	0
			6	6		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	220	Total O 220 220	0	0
6	C	200	Total O 200 200	0	0
6	E	60	Total O 60 60	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Maltose-binding periplasmic protein



- Molecule 1: Maltose-binding periplasmic protein



- Molecule 1: Maltose-binding periplasmic protein



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain B:  50% 50%


GLC1
GLC2

- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain D:  50% 50%


GLC1
GLC2

- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain F:  50% 50%


GLC1
GLC2

4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	100.66Å 63.65Å 221.83Å 90.00° 90.06° 90.00°	Depositor
Resolution (Å)	49.09 – 1.85 73.94 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.09-1.85) 99.8 (73.94-1.85)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 1.84Å)	Xtrriage
Refinement program	PHENIX 1.7.1_743	Depositor
R, R_{free}	0.208 , 0.247 0.206 , 0.242	Depositor DCC
R_{free} test set	5993 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtrriage
Anisotropy	0.170	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.018 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9260	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, ACT, ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2963	0.54	1/4030 (0.0%)
1	C	0.41	0/2966	0.54	0/4033
1	E	0.28	0/2919	0.42	0/3973
All	All	0.37	0/8848	0.50	1/12036 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	143	LYS	N-CA-C	5.41	125.62	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2879	0	2821	36	0
1	C	2879	0	2827	36	0
1	E	2845	0	2774	33	0
2	B	23	0	21	0	0
2	D	23	0	21	0	0
2	F	46	0	42	1	0
3	A	10	0	0	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	7	0	0	0	1
3	E	6	0	0	0	0
4	A	12	0	9	2	0
4	C	16	0	12	0	1
4	E	4	0	3	1	0
5	A	18	0	24	4	0
5	C	12	0	16	4	0
6	A	220	0	0	5	1
6	C	200	0	0	5	0
6	E	60	0	0	2	0
All	All	9260	0	8570	105	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:ALA:HA	1:A:144:GLY:H	1.10	1.12
1:A:142:ALA:HA	1:A:144:GLY:N	1.66	1.10
1:A:302:ALA:HB1	5:A:386:GOL:H32	1.43	1.01
1:A:142:ALA:CA	1:A:144:GLY:H	1.80	0.94
1:E:69:GLY:HA3	1:E:333:ASN:O	1.79	0.83

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:374:ZN:ZN	4:C:380:ACT:O[3_455]	1.46	0.74
3:A:375:ZN:ZN	6:A:401:HOH:O[2_555]	1.51	0.69

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	373/372 (100%)	361 (97%)	8 (2%)	4 (1%)	14	4
1	C	374/372 (100%)	364 (97%)	7 (2%)	3 (1%)	19	7
1	E	369/372 (99%)	361 (98%)	8 (2%)	0	100	100
All	All	1116/1116 (100%)	1086 (97%)	23 (2%)	7 (1%)	25	12

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	142	ALA
1	A	143	LYS
1	A	141	LYS
1	A	174	ASN
1	C	174	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/293 (100%)	287 (98%)	6 (2%)	55	40
1	C	291/293 (99%)	288 (99%)	3 (1%)	76	69
1	E	286/293 (98%)	281 (98%)	5 (2%)	60	47
All	All	870/879 (99%)	856 (98%)	14 (2%)	62	49

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	259	PHE
1	C	338	SER
1	E	329	GLU
1	E	101	ASN
1	E	259	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	202	ASN
1	E	283	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	1[A]	2	12,12,12	0.53	0	17,17,17	1.00	1 (5%)
2	GLC	B	2[A]	2	11,11,12	0.59	0	15,15,17	0.91	0
2	GLC	D	1[A]	2	12,12,12	0.48	0	17,17,17	0.91	0
2	GLC	D	2[A]	2	11,11,12	0.55	0	15,15,17	1.16	2 (13%)
2	GLC	F	1[A]	2	12,12,12	0.52	0	17,17,17	0.77	0
2	GLC	F	1[B]	2	12,12,12	0.59	0	17,17,17	1.54	3 (17%)
2	GLC	F	2[A]	2	11,11,12	0.63	0	15,15,17	0.93	1 (6%)
2	GLC	F	2[B]	2	11,11,12	0.64	0	15,15,17	0.92	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1[A]	2	-	1/2/22/22	0/1/1/1
2	GLC	B	2[A]	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1[A]	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2[A]	2	-	0/2/19/22	0/1/1/1
2	GLC	F	1[A]	2	-	1/2/22/22	0/1/1/1
2	GLC	F	1[B]	2	-	0/2/22/22	0/1/1/1
2	GLC	F	2[A]	2	-	0/2/19/22	0/1/1/1
2	GLC	F	2[B]	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1[B]	GLC	O5-C1-C2	4.32	118.00	110.28
2	F	1[B]	GLC	C1-O5-C5	3.28	119.85	113.66
2	D	2[A]	GLC	C1-O5-C5	3.09	116.38	112.19
2	F	1[B]	GLC	C1-C2-C3	2.62	115.75	110.31
2	B	1[A]	GLC	C1-O5-C5	2.34	118.07	113.66

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	1[A]	GLC	C4-C5-C6-O6
2	B	1[A]	GLC	C4-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	2[A]	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 23 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACT	A	384	3	3,3,3	1.01	0	3,3,3	1.44	1 (33%)
4	ACT	E	379	3	3,3,3	0.76	0	3,3,3	1.33	0
5	GOL	C	384	-	5,5,5	0.38	0	5,5,5	0.21	0
4	ACT	A	383	3	3,3,3	0.91	0	3,3,3	1.09	0
4	ACT	A	385	3	3,3,3	0.91	0	3,3,3	1.55	1 (33%)
4	ACT	C	382	-	3,3,3	0.87	0	3,3,3	1.42	0
4	ACT	C	381	-	3,3,3	0.60	0	3,3,3	1.39	0
5	GOL	A	386	-	5,5,5	0.45	0	5,5,5	0.35	0
4	ACT	C	385	3	3,3,3	0.70	0	3,3,3	1.36	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACT	C	380	-	3,3,3	0.67	0	3,3,3	1.66	1 (33%)
5	GOL	A	388	-	5,5,5	0.37	0	5,5,5	0.29	0
5	GOL	C	383	-	5,5,5	0.38	0	5,5,5	0.53	0
5	GOL	A	387	-	5,5,5	0.33	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	C	384	-	-	2/4/4/4	-
5	GOL	A	386	-	-	4/4/4/4	-
5	GOL	A	388	-	-	2/4/4/4	-
5	GOL	C	383	-	-	2/4/4/4	-
5	GOL	A	387	-	-	1/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	380	ACT	OXT-C-CH3	2.16	124.09	115.18
4	A	385	ACT	OXT-C-O	-2.07	114.41	122.05
4	A	384	ACT	OXT-C-O	-2.01	114.63	122.05

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	386	GOL	C1-C2-C3-O3
5	A	388	GOL	O1-C1-C2-C3
5	A	386	GOL	O1-C1-C2-C3
5	C	383	GOL	O1-C1-C2-C3
5	A	386	GOL	O1-C1-C2-O2

There are no ring outliers.

8 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	384	ACT	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	379	ACT	1	0
5	C	384	GOL	1	0
4	A	383	ACT	1	0
5	A	386	GOL	3	0
4	C	380	ACT	0	1
5	C	383	GOL	3	0
5	A	387	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	371/372 (99%)	0.01	4 (1%) 80 81	11, 23, 43, 73	0
1	C	371/372 (99%)	0.06	2 (0%) 91 91	10, 23, 45, 65	0
1	E	370/372 (99%)	0.60	18 (4%) 29 28	26, 43, 64, 88	0
All	All	1112/1116 (99%)	0.22	24 (2%) 62 61	10, 30, 58, 88	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	269	ALA	3.9
1	C	371	ALA	3.8
1	A	1	MET	3.8
1	E	291	LEU	3.5
1	A	371	ALA	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	B	1[A]	12/12	0.96	0.10	7,16,21,23	0
2	GLC	F	1[A]	12/12	0.96	0.11	24,29,32,34	12
2	GLC	F	1[B]	12/12	0.96	0.11	25,30,33,33	12

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	B	2[A]	11/12	0.97	0.11	13,17,19,20	0
2	GLC	F	2[A]	11/12	0.97	0.11	26,28,30,31	11
2	GLC	F	2[B]	11/12	0.97	0.11	26,28,30,30	11
2	GLC	D	1[A]	12/12	0.98	0.11	10,15,22,22	0
2	GLC	D	2[A]	11/12	0.98	0.10	16,17,19,21	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACT	E	379	4/4	0.68	0.18	52,52,55,55	0
5	GOL	C	384	6/6	0.83	0.19	44,52,53,53	0
5	GOL	A	386	6/6	0.85	0.23	35,41,41,43	0
4	ACT	C	385	4/4	0.85	0.12	40,49,50,56	0
4	ACT	A	383	4/4	0.87	0.18	23,25,32,35	0
3	ZN	E	378	1/1	0.88	0.39	46,46,46,46	1
3	ZN	A	380	1/1	0.90	0.11	85,85,85,85	0
3	ZN	A	379	1/1	0.91	0.11	51,51,51,51	0
5	GOL	A	387	6/6	0.91	0.16	26,46,48,51	0
3	ZN	C	379	1/1	0.91	0.08	53,53,53,53	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ZN	A	381	1/1	0.92	0.12	45,45,45,45	1
4	ACT	A	384	4/4	0.92	0.14	19,21,24,26	0
4	ACT	C	381	4/4	0.92	0.16	20,21,32,42	0
3	ZN	C	375	1/1	0.92	0.12	44,44,44,44	0
5	GOL	C	383	6/6	0.93	0.15	23,40,40,42	0
5	GOL	A	388	6/6	0.93	0.12	54,61,63,65	0
4	ACT	A	385	4/4	0.94	0.16	19,25,28,38	0
4	ACT	C	382	4/4	0.96	0.15	22,22,27,34	0
3	ZN	A	375	1/1	0.96	0.19	21,21,21,21	1
4	ACT	C	380	4/4	0.96	0.14	9,17,22,29	0
3	ZN	C	376	1/1	0.96	0.10	45,45,45,45	0
3	ZN	E	373	1/1	0.97	0.09	34,34,34,34	0
3	ZN	C	377	1/1	0.97	0.09	33,33,33,33	1
3	ZN	A	374	1/1	0.97	0.14	25,25,25,25	0
3	ZN	E	377	1/1	0.98	0.11	50,50,50,50	1
3	ZN	C	378	1/1	0.98	0.11	39,39,39,39	1
3	ZN	A	382	1/1	0.98	0.05	38,38,38,38	1
3	ZN	A	378	1/1	0.98	0.05	43,43,43,43	0
3	ZN	E	376	1/1	0.98	0.05	60,60,60,60	0
3	ZN	A	377	1/1	0.99	0.08	30,30,30,30	1
3	ZN	E	375	1/1	0.99	0.04	54,54,54,54	0
3	ZN	A	373	1/1	0.99	0.13	18,18,18,18	0
3	ZN	A	376	1/1	0.99	0.07	30,30,30,30	1
3	ZN	C	373	1/1	0.99	0.11	17,17,17,17	0
3	ZN	C	374	1/1	0.99	0.10	24,24,24,24	0
3	ZN	E	374	1/1	1.00	0.08	39,39,39,39	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.