



# wwPDB X-ray Structure Validation Summary Report i

Sep 3, 2023 – 08:40 PM EDT

PDB ID : 3SWD  
Title : E. coli MurA in complex with UDP-N-acetylmuramic acid and covalent adduct of PEP with Cys115  
Authors : Zhu, J.-Y.; Schonbrunn, E.  
Deposited on : 2011-07-13  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

*Continued from previous page...*



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	IAS	E	67	-	X	-	-
1	IAS	I	67	-	-	X	-
1	IAS	K	67	-	-	X	-

## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 39035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-N-acetylglucosamine 1-carboxyvinyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	B	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	C	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	D	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	E	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	F	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	G	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	H	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	I	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	J	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	K	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			
1	L	418	Total	C	N	O	P	S	0	0	0
			3143	1970	556	600	1	16			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	67	IAS	ASN	engineered mutation	UNP P0A749
B	67	IAS	ASN	engineered mutation	UNP P0A749
C	67	IAS	ASN	engineered mutation	UNP P0A749
D	67	IAS	ASN	engineered mutation	UNP P0A749
E	67	IAS	ASN	engineered mutation	UNP P0A749

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	67	IAS	ASN	engineered mutation	UNP P0A749
G	67	IAS	ASN	engineered mutation	UNP P0A749
H	67	IAS	ASN	engineered mutation	UNP P0A749
I	67	IAS	ASN	engineered mutation	UNP P0A749
J	67	IAS	ASN	engineered mutation	UNP P0A749
K	67	IAS	ASN	engineered mutation	UNP P0A749
L	67	IAS	ASN	engineered mutation	UNP P0A749

- Molecule 2 is (2R)-2-{{[(2R,3R,4R,5S,6R)-3-(acetylamino)-2-{{[(S)-{[(R)-{[(2R,3S,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3,4-dihydroxytetrahydrofuran-2-yl]methoxy}(hydroxy)phosphoryl]oxy}(hydroxy)phosphoryl]oxy}-5-hydroxy-6-(hydroxymethyl)tetrahydro-2H-pyran-4-yl]oxy}propanoic acid (three-letter code: EPZ) (formula: C<sub>20</sub>H<sub>31</sub>N<sub>3</sub>O<sub>19</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	20	3	19	2		
2	B	1	Total	C	N	O	P	0	0
			44	20	3	19	2		
2	C	1	Total	C	N	O	P	0	0
			44	20	3	19	2		
2	D	1	Total	C	N	O	P	0	0
			44	20	3	19	2		
2	E	1	Total	C	N	O	P	0	0
			44	20	3	19	2		
2	F	1	Total	C	N	O	P	0	0
			44	20	3	19	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total C N O P 44 20 3 19 2	0	0
2	H	1	Total C N O P 44 20 3 19 2	0	0
2	I	1	Total C N O P 44 20 3 19 2	0	0
2	J	1	Total C N O P 44 20 3 19 2	0	0
2	K	1	Total C N O P 44 20 3 19 2	0	0
2	L	1	Total C N O P 44 20 3 19 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	76	Total O 76 76	0	0
3	B	58	Total O 58 58	0	0
3	C	105	Total O 105 105	0	0
3	D	104	Total O 104 104	0	0
3	E	86	Total O 86 86	0	0
3	F	63	Total O 63 63	0	0
3	G	47	Total O 47 47	0	0
3	H	73	Total O 73 73	0	0
3	I	42	Total O 42 42	0	0
3	J	45	Total O 45 45	0	0
3	K	60	Total O 60 60	0	0
3	L	32	Total O 32 32	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase



- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase



- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase





- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase

Chain D:




- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase

Chain E:



- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase

Chain F:




- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase


Chain G:



- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase



- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase




- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase








- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase



- Molecule 1: UDP-N-acetylglucosamine 1-carboxyvinyltransferase





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.51 Å    120.91 Å    139.73 Å 111.52°    104.44°    90.19°	Depositor
Resolution (Å)	47.48 – 2.50 47.48 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.5 (47.48-2.50) 97.6 (47.48-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.49 (at 2.51 Å)	Xtriage
Refinement program	CNS 1.3	Depositor
$R$ , $R_{free}$	0.226 , 0.280 0.219 , 0.272	Depositor DCC
$R_{free}$ test set	8367 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.9	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 54.8	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	39035	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: QPA, IAS, EPZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.56	0/3161	0.97	3/4278 (0.1%)
1	B	0.55	0/3161	0.90	2/4278 (0.0%)
1	C	0.61	1/3161 (0.0%)	0.92	4/4278 (0.1%)
1	D	0.62	0/3161	1.09	6/4278 (0.1%)
1	E	0.58	0/3161	0.91	2/4278 (0.0%)
1	F	0.59	3/3161 (0.1%)	0.92	2/4278 (0.0%)
1	G	0.56	0/3161	0.90	2/4278 (0.0%)
1	H	0.59	0/3161	0.92	2/4278 (0.0%)
1	I	0.55	0/3161	0.91	2/4278 (0.0%)
1	J	0.70	2/3161 (0.1%)	1.01	0/4278
1	K	0.72	0/3161	1.00	2/4278 (0.0%)
1	L	0.52	0/3161	1.07	7/4278 (0.2%)
All	All	0.60	6/37932 (0.0%)	0.96	34/51336 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	K	0	2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	J	381	CYS	CB-SG	-5.45	1.73	1.81
1	J	330	ASN	CG-OD1	5.32	1.35	1.24
1	C	350	ASN	CG-OD1	5.14	1.35	1.24
1	F	381	CYS	CB-SG	-5.10	1.73	1.81
1	F	184	ASN	CG-OD1	5.04	1.35	1.24

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	75	ARG	NE-CZ-NH1	-20.39	110.11	120.30
1	D	120	ARG	NE-CZ-NH1	-20.14	110.23	120.30
1	D	212	ARG	NE-CZ-NH1	-19.88	110.36	120.30
1	L	75	ARG	NE-CZ-NH2	19.34	129.97	120.30
1	L	212	ARG	NE-CZ-NH1	-19.15	110.73	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	K	187	ARG	Sidechain
1	K	340	ARG	Sidechain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3143	0	3210	66	1
1	B	3143	0	3211	77	0
1	C	3143	0	3210	64	0
1	D	3143	0	3210	60	0
1	E	3143	0	3211	76	0
1	F	3143	0	3211	82	0
1	G	3143	0	3211	66	0
1	H	3143	0	3210	73	0
1	I	3143	0	3211	74	1
1	J	3143	0	3211	171	0
1	K	3143	0	3211	152	0
1	L	3143	0	3210	65	0
2	A	44	0	28	0	0
2	B	44	0	28	2	0
2	C	44	0	28	1	0
2	D	44	0	28	0	0
2	E	44	0	28	0	0
2	F	44	0	28	0	0
2	G	44	0	28	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	44	0	28	0	0
2	I	44	0	28	2	0
2	J	44	0	28	2	0
2	K	44	0	28	3	0
2	L	44	0	28	1	0
3	A	76	0	0	4	0
3	B	58	0	0	7	0
3	C	105	0	0	4	0
3	D	104	0	0	7	0
3	E	86	0	0	5	0
3	F	63	0	0	4	0
3	G	47	0	0	5	0
3	H	73	0	0	5	0
3	I	42	0	0	4	0
3	J	45	0	0	10	0
3	K	60	0	0	5	0
3	L	32	0	0	5	0
All	All	39035	0	38863	984	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 984 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:11:LYS:NZ	1:C:13:GLN:OE1	1.75	1.18
1:J:405:LYS:O	1:J:409:LEU:HD12	1.46	1.13
1:E:403:GLU:HG3	3:E:660:HOH:O	1.46	1.13
1:K:66:ARG:HG3	1:K:67:IAS:OXT	1.51	1.09
1:K:370:LEU:HD11	1:K:397:ARG:HB2	1.39	1.04

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:414:GLU:OE1	1:I:46:LYS:NZ[1_666]	2.06	0.14

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	413/418 (99%)	396 (96%)	15 (4%)	2 (0%)	29 48
1	B	413/418 (99%)	400 (97%)	13 (3%)	0	100 100
1	C	413/418 (99%)	398 (96%)	15 (4%)	0	100 100
1	D	413/418 (99%)	397 (96%)	15 (4%)	1 (0%)	47 68
1	E	413/418 (99%)	396 (96%)	17 (4%)	0	100 100
1	F	413/418 (99%)	396 (96%)	16 (4%)	1 (0%)	47 68
1	G	413/418 (99%)	397 (96%)	16 (4%)	0	100 100
1	H	413/418 (99%)	397 (96%)	15 (4%)	1 (0%)	47 68
1	I	413/418 (99%)	394 (95%)	18 (4%)	1 (0%)	47 68
1	J	413/418 (99%)	368 (89%)	43 (10%)	2 (0%)	29 48
1	K	413/418 (99%)	378 (92%)	33 (8%)	2 (0%)	29 48
1	L	413/418 (99%)	397 (96%)	15 (4%)	1 (0%)	47 68
All	All	4956/5016 (99%)	4714 (95%)	231 (5%)	11 (0%)	47 68

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	91	ARG
1	A	91	ARG
1	D	91	ARG
1	H	91	ARG
1	K	329	GLU

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	329/329 (100%)	317 (96%)	12 (4%)	35 61
1	B	329/329 (100%)	317 (96%)	12 (4%)	35 61
1	C	329/329 (100%)	317 (96%)	12 (4%)	35 61
1	D	329/329 (100%)	317 (96%)	12 (4%)	35 61
1	E	329/329 (100%)	317 (96%)	12 (4%)	35 61
1	F	329/329 (100%)	317 (96%)	12 (4%)	35 61
1	G	329/329 (100%)	316 (96%)	13 (4%)	31 56
1	H	329/329 (100%)	316 (96%)	13 (4%)	31 56
1	I	329/329 (100%)	317 (96%)	12 (4%)	35 61
1	J	329/329 (100%)	304 (92%)	25 (8%)	13 25
1	K	329/329 (100%)	313 (95%)	16 (5%)	25 47
1	L	329/329 (100%)	316 (96%)	13 (4%)	31 56
All	All	3948/3948 (100%)	3784 (96%)	164 (4%)	30 54

5 of 164 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	108	GLN
1	K	160	LYS
1	J	188	GLU
1	J	349	SER
1	K	416	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	307	GLN
1	L	155	HIS
1	F	155	HIS
1	G	155	HIS
1	H	155	HIS

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

24 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	IAS	E	67	1	6,7,8	1.80	2 (33%)	6,8,10	3.66	3 (50%)
1	QPA	B	115	1	10,15,16	3.79	4 (40%)	9,22,24	1.43	1 (11%)
1	IAS	L	67	1	6,7,8	0.84	0	6,8,10	1.18	0
1	QPA	F	115	1	10,15,16	3.85	4 (40%)	9,22,24	1.54	1 (11%)
1	QPA	J	115	1	10,15,16	3.74	4 (40%)	9,22,24	1.68	1 (11%)
1	QPA	L	115	1	10,15,16	3.62	4 (40%)	9,22,24	1.45	1 (11%)
1	QPA	A	115	1	10,15,16	3.99	4 (40%)	9,22,24	1.58	1 (11%)
1	IAS	F	67	1	6,7,8	1.51	1 (16%)	6,8,10	3.60	2 (33%)
1	IAS	B	67	1	6,7,8	0.85	0	6,8,10	1.24	0
1	IAS	A	67	1	6,7,8	0.92	0	6,8,10	1.21	0
1	IAS	H	67	1	6,7,8	1.00	0	6,8,10	1.27	0
1	QPA	C	115	1	10,15,16	3.54	4 (40%)	9,22,24	1.42	1 (11%)
1	QPA	G	115	1	10,15,16	3.60	5 (50%)	9,22,24	1.53	1 (11%)
1	QPA	I	115	1	10,15,16	4.12	4 (40%)	9,22,24	1.40	1 (11%)
1	IAS	K	67	1	6,7,8	1.86	2 (33%)	6,8,10	3.84	4 (66%)
1	IAS	C	67	1	6,7,8	0.97	0	6,8,10	1.25	0
1	IAS	G	67	1	6,7,8	0.93	0	6,8,10	1.14	0
1	IAS	I	67	1	6,7,8	1.59	1 (16%)	6,8,10	3.68	3 (50%)
1	QPA	D	115	1	10,15,16	3.76	5 (50%)	9,22,24	1.58	1 (11%)
1	QPA	E	115	1	10,15,16	3.92	4 (40%)	9,22,24	1.40	1 (11%)
1	QPA	K	115	1	10,15,16	3.87	5 (50%)	9,22,24	1.65	1 (11%)
1	IAS	J	67	1	6,7,8	1.02	0	6,8,10	1.18	0
1	QPA	H	115	1	10,15,16	3.65	4 (40%)	9,22,24	1.44	1 (11%)
1	IAS	D	67	1	6,7,8	1.07	0	6,8,10	1.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	IAS	E	67	1	-	4/7/7/8	-
1	QPA	B	115	1	-	1/9/20/22	-
1	IAS	L	67	1	-	2/7/7/8	-
1	QPA	F	115	1	-	1/9/20/22	-
1	QPA	J	115	1	-	1/9/20/22	-
1	QPA	L	115	1	-	0/9/20/22	-
1	QPA	A	115	1	-	1/9/20/22	-
1	IAS	F	67	1	-	3/7/7/8	-
1	IAS	B	67	1	-	2/7/7/8	-
1	IAS	A	67	1	-	2/7/7/8	-
1	IAS	H	67	1	-	0/7/7/8	-
1	QPA	C	115	1	-	1/9/20/22	-
1	QPA	G	115	1	-	1/9/20/22	-
1	QPA	I	115	1	-	1/9/20/22	-
1	IAS	K	67	1	-	1/7/7/8	-
1	IAS	C	67	1	-	2/7/7/8	-
1	IAS	G	67	1	-	2/7/7/8	-
1	IAS	I	67	1	-	2/7/7/8	-
1	QPA	D	115	1	-	1/9/20/22	-
1	QPA	E	115	1	-	1/9/20/22	-
1	QPA	K	115	1	-	0/9/20/22	-
1	IAS	J	67	1	-	4/7/7/8	-
1	QPA	H	115	1	-	1/9/20/22	-
1	IAS	D	67	1	-	1/7/7/8	-

The worst 5 of 57 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	115	QPA	P2-O14	9.75	1.77	1.59
1	A	115	QPA	P2-O14	9.56	1.77	1.59
1	E	115	QPA	P2-O14	8.79	1.75	1.59
1	J	115	QPA	P2-O14	8.67	1.75	1.59
1	F	115	QPA	P2-O14	8.60	1.75	1.59

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	I	67	IAS	OXT-C-O	7.77	141.73	124.09
1	K	67	IAS	OXT-C-O	7.39	140.87	124.09

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	67	IAS	OXT-C-O	7.23	140.50	124.09
1	E	67	IAS	OXT-C-O	6.98	139.94	124.09
1	F	67	IAS	O-C-CA	-4.97	104.61	122.14

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	E	67	IAS	N-CA-CB-CG
1	J	67	IAS	O-C-CA-N
1	J	67	IAS	OXT-C-CA-N
1	F	67	IAS	OXT-C-CA-CB
1	J	67	IAS	O-C-CA-CB

There are no ring outliers.

5 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	J	115	QPA	1	0
1	F	67	IAS	3	0
1	K	67	IAS	5	0
1	I	67	IAS	8	0
1	K	115	QPA	1	0

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EPZ	J	501	-	43,46,46	1.33	7 (16%)	60,69,69	1.62	7 (11%)
2	EPZ	G	501	-	43,46,46	1.33	6 (13%)	60,69,69	1.63	9 (15%)
2	EPZ	D	501	-	43,46,46	1.48	10 (23%)	60,69,69	1.64	9 (15%)
2	EPZ	I	501	-	43,46,46	1.31	8 (18%)	60,69,69	1.49	7 (11%)
2	EPZ	H	501	-	43,46,46	1.44	6 (13%)	60,69,69	1.55	8 (13%)
2	EPZ	E	501	-	43,46,46	1.39	7 (16%)	60,69,69	1.59	11 (18%)
2	EPZ	K	501	-	43,46,46	1.41	7 (16%)	60,69,69	1.65	6 (10%)
2	EPZ	B	501	-	43,46,46	1.38	6 (13%)	60,69,69	1.64	8 (13%)
2	EPZ	A	501	-	43,46,46	1.46	9 (20%)	60,69,69	1.56	9 (15%)
2	EPZ	L	501	-	43,46,46	1.32	4 (9%)	60,69,69	1.64	9 (15%)
2	EPZ	F	501	-	43,46,46	1.62	9 (20%)	60,69,69	1.59	8 (13%)
2	EPZ	C	501	-	43,46,46	1.50	8 (18%)	60,69,69	1.66	8 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EPZ	J	501	-	-	5/34/71/71	0/3/3/3
2	EPZ	G	501	-	-	5/34/71/71	0/3/3/3
2	EPZ	D	501	-	-	4/34/71/71	0/3/3/3
2	EPZ	I	501	-	-	5/34/71/71	0/3/3/3
2	EPZ	H	501	-	-	7/34/71/71	0/3/3/3
2	EPZ	E	501	-	-	4/34/71/71	0/3/3/3
2	EPZ	K	501	-	-	6/34/71/71	0/3/3/3
2	EPZ	B	501	-	-	7/34/71/71	0/3/3/3
2	EPZ	A	501	-	-	5/34/71/71	0/3/3/3
2	EPZ	L	501	-	-	5/34/71/71	0/3/3/3
2	EPZ	F	501	-	-	7/34/71/71	0/3/3/3
2	EPZ	C	501	-	-	4/34/71/71	0/3/3/3

The worst 5 of 87 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	K	501	EPZ	C2U-N1U	4.49	1.45	1.38
2	B	501	EPZ	C2U-N1U	4.31	1.45	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	501	EPZ	C2U-N1U	4.13	1.45	1.38
2	H	501	EPZ	C2U-N1U	3.85	1.44	1.38
2	A	501	EPZ	C2U-N1U	3.49	1.44	1.38

The worst 5 of 99 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	501	EPZ	C4U-N3U-C2U	-5.34	119.53	126.58
2	B	501	EPZ	C4U-N3U-C2U	-5.09	119.86	126.58
2	K	501	EPZ	C4U-N3U-C2U	-5.07	119.90	126.58
2	F	501	EPZ	C4U-N3U-C2U	-5.06	119.91	126.58
2	G	501	EPZ	C4U-N3U-C2U	-5.05	119.91	126.58

There are no chirality outliers.

5 of 64 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	EPZ	O1E-C1E-C2E-O3
2	A	501	EPZ	O2E-C1E-C2E-O3
2	F	501	EPZ	O4D-C4D-C5D-O5D
2	H	501	EPZ	C3E-C2E-O3-C3
2	L	501	EPZ	O1E-C1E-C2E-O3


There are no ring outliers.



7 monomers are involved in 12 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	J	501	EPZ	2	0
2	G	501	EPZ	1	0
2	I	501	EPZ	2	0
2	K	501	EPZ	3	0
2	B	501	EPZ	2	0
2	L	501	EPZ	1	0
2	C	501	EPZ	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.















## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	416/418 (99%)	-0.17	4 (0%) 82 84	36, 54, 76, 97	0
1	B	416/418 (99%)	0.23	18 (4%) 35 38	37, 61, 83, 100	0
1	C	416/418 (99%)	-0.23	1 (0%) 95 95	30, 47, 69, 95	0
1	D	416/418 (99%)	-0.31	0 100 100	25, 47, 68, 95	0
1	E	416/418 (99%)	-0.12	3 (0%) 87 89	30, 50, 72, 102	0
1	F	416/418 (99%)	-0.05	3 (0%) 87 89	29, 53, 77, 100	0
1	G	416/418 (99%)	0.11	13 (3%) 49 52	33, 60, 83, 104	0
1	H	416/418 (99%)	-0.24	3 (0%) 87 89	34, 50, 70, 95	0
1	I	416/418 (99%)	0.44	42 (10%) 7 6	39, 67, 93, 110	0
1	J	416/418 (99%)	-0.15	5 (1%) 79 80	37, 58, 79, 100	0
1	K	416/418 (99%)	-0.25	5 (1%) 79 80	29, 50, 73, 88	0
1	L	416/418 (99%)	0.69	55 (13%) 3 3	43, 72, 94, 120	0
All	All	4992/5016 (99%)	-0.00	152 (3%) 50 53	25, 55, 83, 120	0

The worst 5 of 152 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	221	LEU	8.8
1	L	222	GLY	8.2
1	L	215	ILE	7.0
1	I	215	ILE	6.8
1	I	81	CYS	5.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	IAS	I	67	8/9	0.65	0.27	79,89,111,120	0
1	IAS	L	67	8/9	0.82	0.26	81,89,99,102	0
1	IAS	A	67	8/9	0.89	0.15	50,56,66,69	0
1	IAS	G	67	8/9	0.90	0.18	64,67,75,77	0
1	QPA	B	115	16/17	0.91	0.12	46,64,92,95	0
1	QPA	A	115	16/17	0.92	0.22	49,62,92,92	0
1	IAS	C	67	8/9	0.92	0.15	61,69,72,72	0
1	IAS	D	67	8/9	0.92	0.15	42,54,62,62	0
1	QPA	I	115	16/17	0.93	0.15	46,65,92,92	0
1	IAS	F	67	8/9	0.93	0.13	52,64,76,80	0
1	QPA	E	115	16/17	0.93	0.15	42,62,77,80	0
1	IAS	B	67	8/9	0.93	0.19	66,72,84,90	0
1	IAS	K	67	8/9	0.93	0.14	59,61,65,71	0
1	QPA	F	115	16/17	0.93	0.20	41,60,80,86	0
1	QPA	H	115	16/17	0.94	0.17	40,63,76,77	0
1	QPA	L	115	16/17	0.94	0.15	50,65,81,85	0
1	IAS	H	67	8/9	0.94	0.20	53,66,80,88	0
1	QPA	K	115	16/17	0.95	0.18	36,57,80,80	0
1	IAS	J	67	8/9	0.95	0.13	53,75,79,84	0
1	QPA	C	115	16/17	0.95	0.19	38,62,79,83	0
1	IAS	E	67	8/9	0.95	0.08	41,53,62,78	0
1	QPA	D	115	16/17	0.96	0.17	30,51,68,68	0
1	QPA	G	115	16/17	0.96	0.15	41,58,76,79	0
1	QPA	J	115	16/17	0.98	0.15	37,55,84,103	0

### 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EPZ	I	501	44/44	0.94	0.17	38,58,68,84	0
2	EPZ	G	501	44/44	0.96	0.18	31,48,65,67	0


*Continued on next page...*


*Continued from previous page...*


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EPZ	L	501	44/44	0.96	0.18	25,60,76,99	0
2	EPZ	D	501	44/44	0.97	0.14	25,43,58,70	0
2	EPZ	A	501	44/44	0.97	0.17	28,46,60,63	0
2	EPZ	H	501	44/44	0.97	0.14	27,44,58,66	0
2	EPZ	B	501	44/44	0.97	0.13	27,51,73,82	0
2	EPZ	J	501	44/44	0.97	0.14	30,45,64,70	0
2	EPZ	K	501	44/44	0.97	0.14	27,43,60,66	0
2	EPZ	C	501	44/44	0.97	0.16	28,40,57,66	0
2	EPZ	E	501	44/44	0.98	0.16	31,44,60,82	0
2	EPZ	F	501	44/44	0.98	0.16	20,38,59,78	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

















## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.