



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 3, 2023 – 08:20 AM EDT

PDB ID : 3VSV
Title : The complex structure of XylC with xylose
Authors : Huang, C.H.; Sun, Y.; Ko, T.P.; Ma, Y.; Chen, C.C.; Zheng, Y.; Chan, H.C.;
Pang, X.; Wiegel, J.; Shao, W.; Guo, R.T.
Deposited on : 2012-05-09
Resolution : 1.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4690 (1.50-1.46)
Clashscore	141614	4955 (1.50-1.46)
Ramachandran outliers	138981	4846 (1.50-1.46)
Sidechain outliers	138945	4844 (1.50-1.46)
RSRZ outliers	127900	4614 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	638	 3% 94% 6%
1	B	638	 2% 93% 6%
1	C	638	 3% 94% 6%
1	D	638	 2% 92% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	XYP	A	707	-	-	X	-
2	XYP	D	701	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 24551 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	638	Total 5147	C 3275	N 877	O 983	S 12	0	0	0
1	B	638	Total 5147	C 3275	N 877	O 983	S 12	0	0	0
1	C	638	Total 5147	C 3275	N 877	O 983	S 12	0	0	0
1	D	638	Total 5147	C 3275	N 877	O 983	S 12	0	0	0

- Molecule 2 is beta-D-xylopyranose (three-letter code: XYP) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 10	C 5	O 5	0	0
2	A	1	Total 10	C 5	O 5	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	5	5		
2	A	1	Total	C	O	0	0
			10	5	5		
2	A	1	Total	C	O	0	0
			10	5	5		
2	A	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			10	5	5		
2	C	1	Total	C	O	0	0
			10	5	5		
2	C	1	Total	C	O	0	0
			10	5	5		
2	C	1	Total	C	O	0	0
			10	5	5		
2	C	1	Total	C	O	0	0
			10	5	5		
2	C	1	Total	C	O	0	0
			10	5	5		
2	D	1	Total	C	O	0	0
			10	5	5		
2	D	1	Total	C	O	0	0
			10	5	5		
2	D	1	Total	C	O	0	0
			10	5	5		
2	D	1	Total	C	O	0	0
			10	5	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	O	0	0
			10	5	5		
2	D	1	Total	C	O	0	0
			10	5	5		
2	D	1	Total	C	O	0	0
			10	5	5		

- Molecule 3 is alpha-D-xylopyranose (three-letter code: XYZ) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	5	5		
3	A	1	Total	C	O	0	0
			10	5	5		
3	B	1	Total	C	O	0	0
			10	5	5		
3	B	1	Total	C	O	0	0
			10	5	5		
3	C	1	Total	C	O	0	0
			10	5	5		
3	C	1	Total	C	O	0	0
			10	5	5		
3	C	1	Total	C	O	0	0
			10	5	5		
3	D	1	Total	C	O	0	0
			10	5	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			10	5	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	912	Total	O	0	0
			912	912		
4	B	942	Total	O	0	0
			942	942		
4	C	842	Total	O	0	0
			842	842		
4	D	917	Total	O	0	0
			917	917		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Xylosidase



- Molecule 1: Xylosidase




- Molecule 1: Xylosidase



- Molecule 1: Xylosidase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	88.78Å 201.95Å 100.11Å 90.00° 99.18° 90.00°	Depositor
Resolution (Å)	25.00 – 1.48 24.89 – 1.48	Depositor EDS
% Data completeness (in resolution range)	(Not available) (25.00-1.48) 95.0 (24.89-1.48)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.50 (at 1.48Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.150 , 0.168 0.146 , 0.164	Depositor DCC
R_{free} test set	27272 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	10.9	Xtrriage
Anisotropy	0.254	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 54.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	24551	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, XYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	1/5276 (0.0%)	0.87	5/7151 (0.1%)
1	B	0.73	0/5276	0.88	8/7151 (0.1%)
1	C	0.71	0/5276	0.89	8/7151 (0.1%)
1	D	0.72	0/5276	0.88	5/7151 (0.1%)
All	All	0.72	1/21104 (0.0%)	0.88	26/28604 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	509	ASN	CB-CG	5.54	1.63	1.51

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	298	ARG	NE-CZ-NH2	-15.23	112.68	120.30
1	C	298	ARG	NE-CZ-NH1	14.98	127.79	120.30
1	D	589	ARG	NE-CZ-NH2	-9.58	115.51	120.30
1	D	589	ARG	NE-CZ-NH1	9.22	124.91	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	589	ARG	NE-CZ-NH2	-8.80	115.90	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	105	TYR	Sidechain
1	B	105	TYR	Sidechain
1	C	105	TYR	Sidechain
1	D	105	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5147	0	4962	39	0
1	B	5147	0	4962	35	0
1	C	5147	0	4962	41	0
1	D	5147	0	4962	41	0
2	A	70	0	0	9	0
2	B	60	0	0	0	0
2	C	50	0	0	2	0
2	D	80	0	0	11	0
3	A	20	0	20	4	0
3	B	20	0	20	0	0
3	C	30	0	30	2	0
3	D	20	0	20	3	0
4	A	912	0	0	8	0
4	B	942	0	0	16	0
4	C	842	0	0	6	0
4	D	917	0	0	15	0
All	All	24551	0	19938	148	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 148 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:509:ASN:HB3	4:A:1676:HOH:O	1.34	1.27
1:C:509:ASN:HB3	4:C:1589:HOH:O	1.32	1.23
2:D:701:XYP:C4	4:D:1665:HOH:O	1.86	1.20
1:C:337:ASN:HB2	4:C:1599:HOH:O	1.47	1.15
1:A:427:MET:CE	1:A:445:ARG:HH21	1.58	1.15

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	636/638 (100%)	612 (96%)	24 (4%)	0	100	100
1	B	636/638 (100%)	612 (96%)	24 (4%)	0	100	100
1	C	636/638 (100%)	610 (96%)	26 (4%)	0	100	100
1	D	636/638 (100%)	614 (96%)	22 (4%)	0	100	100
All	All	2544/2552 (100%)	2448 (96%)	96 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	550/550 (100%)	545 (99%)	5 (1%)	78	59
1	B	550/550 (100%)	547 (100%)	3 (0%)	88	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	550/550 (100%)	545 (99%)	5 (1%)	78	59
1	D	550/550 (100%)	547 (100%)	3 (0%)	88	77
All	All	2200/2200 (100%)	2184 (99%)	16 (1%)	84	68

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	403	PHE
1	D	277	ARG
1	C	18	LYS
1	C	422	TYR
1	B	422	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	619	ASN
1	C	619	ASN
1	B	444	GLN
1	B	366	GLN
1	C	444	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

35 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XYP	D	706	-	10,10,10	0.94	0	14,14,14	1.27	2 (14%)
2	XYP	C	705	-	10,10,10	1.10	1 (10%)	14,14,14	1.23	2 (14%)
2	XYP	B	701	-	10,10,10	2.03	4 (40%)	14,14,14	1.20	1 (7%)
2	XYP	D	702	-	10,10,10	2.15	4 (40%)	14,14,14	1.28	2 (14%)
3	XYS	C	706	-	10,10,10	1.21	1 (10%)	14,14,14	0.83	0
2	XYP	C	701	-	10,10,10	2.35	5 (50%)	14,14,14	1.25	2 (14%)
3	XYS	D	709	-	10,10,10	1.87	3 (30%)	14,14,14	0.91	0
2	XYP	D	704	-	10,10,10	0.92	0	14,14,14	1.05	1 (7%)
3	XYS	B	707	-	10,10,10	1.67	1 (10%)	14,14,14	0.69	0
2	XYP	A	706	-	10,10,10	1.26	0	14,14,14	1.25	2 (14%)
2	XYP	A	703	-	10,10,10	1.16	1 (10%)	14,14,14	0.88	1 (7%)
2	XYP	D	708	-	10,10,10	1.38	2 (20%)	14,14,14	1.08	1 (7%)
3	XYS	C	708	-	10,10,10	1.68	2 (20%)	14,14,14	1.63	3 (21%)
2	XYP	D	707	-	10,10,10	1.11	0	14,14,14	0.76	1 (7%)
3	XYS	B	708	-	10,10,10	1.43	1 (10%)	14,14,14	0.76	0
3	XYS	C	707	-	10,10,10	1.73	2 (20%)	14,14,14	0.72	0
2	XYP	A	702	-	10,10,10	1.09	1 (10%)	14,14,14	0.78	0
3	XYS	A	709	-	10,10,10	1.47	1 (10%)	14,14,14	0.91	1 (7%)
3	XYS	D	710	-	10,10,10	1.31	1 (10%)	14,14,14	0.78	0
2	XYP	A	707	-	10,10,10	0.97	1 (10%)	14,14,14	1.32	1 (7%)
2	XYP	D	703	-	10,10,10	1.06	1 (10%)	14,14,14	0.86	0
2	XYP	A	705	-	10,10,10	1.02	0	14,14,14	1.28	1 (7%)
2	XYP	B	704	-	10,10,10	1.08	1 (10%)	14,14,14	0.76	0
2	XYP	B	702	-	10,10,10	1.16	1 (10%)	14,14,14	1.04	2 (14%)
3	XYS	A	708	-	10,10,10	1.13	0	14,14,14	0.77	0
2	XYP	A	701	-	10,10,10	1.80	3 (30%)	14,14,14	1.23	2 (14%)
2	XYP	D	701	-	10,10,10	0.99	0	14,14,14	0.97	0
2	XYP	B	705	-	10,10,10	1.19	1 (10%)	14,14,14	0.91	0
2	XYP	C	704	-	10,10,10	1.51	2 (20%)	14,14,14	1.12	2 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XYP	A	704	-	10,10,10	1.02	1 (10%)	14,14,14	0.70	0
2	XYP	D	705	-	10,10,10	1.08	0	14,14,14	0.65	0
2	XYP	C	703	-	10,10,10	1.26	1 (10%)	14,14,14	0.86	0
2	XYP	B	706	-	10,10,10	1.12	1 (10%)	14,14,14	0.78	0
2	XYP	B	703	-	10,10,10	0.88	0	14,14,14	0.83	1 (7%)
2	XYP	C	702	-	10,10,10	1.04	0	14,14,14	0.75	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYP	D	706	-	-	-	0/1/1/1
2	XYP	C	705	-	-	-	0/1/1/1
2	XYP	B	701	-	-	-	0/1/1/1
2	XYP	D	702	-	-	-	0/1/1/1
3	XYS	C	706	-	-	-	0/1/1/1
2	XYP	C	701	-	-	-	0/1/1/1
3	XYS	D	709	-	-	-	0/1/1/1
2	XYP	D	704	-	-	-	0/1/1/1
3	XYS	B	707	-	-	-	0/1/1/1
2	XYP	A	706	-	-	-	0/1/1/1
2	XYP	A	703	-	-	-	0/1/1/1
2	XYP	D	708	-	-	-	0/1/1/1
3	XYS	C	708	-	-	-	0/1/1/1
2	XYP	D	707	-	-	-	0/1/1/1
3	XYS	B	708	-	-	-	0/1/1/1
3	XYS	C	707	-	-	-	0/1/1/1
2	XYP	A	702	-	-	-	0/1/1/1
3	XYS	A	709	-	-	-	0/1/1/1
3	XYS	D	710	-	-	-	0/1/1/1
2	XYP	A	707	-	-	-	0/1/1/1
2	XYP	D	703	-	-	-	0/1/1/1
2	XYP	A	705	-	-	-	0/1/1/1
2	XYP	B	704	-	-	-	0/1/1/1
2	XYP	B	702	-	-	-	0/1/1/1
3	XYS	A	708	-	-	-	0/1/1/1
2	XYP	A	701	-	-	-	0/1/1/1
2	XYP	D	701	-	-	-	0/1/1/1
2	XYP	B	705	-	-	-	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYP	C	704	-	-	-	0/1/1/1
2	XYP	A	704	-	-	-	0/1/1/1
2	XYP	D	705	-	-	-	0/1/1/1
2	XYP	C	703	-	-	-	0/1/1/1
2	XYP	B	706	-	-	-	0/1/1/1
2	XYP	B	703	-	-	-	0/1/1/1
2	XYP	C	702	-	-	-	0/1/1/1

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	701	XYP	C4-C3	4.59	1.59	1.52
3	D	709	XYS	O5-C1	4.27	1.49	1.43
3	C	708	XYS	O5-C1	3.81	1.48	1.43
2	D	702	XYP	O1-C1	3.67	1.51	1.39
2	D	702	XYP	C3-C2	3.56	1.61	1.52

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	705	XYP	C5-O5-C1	-3.58	106.69	112.71
3	C	708	XYS	O5-C1-C2	3.44	114.54	109.43
3	C	708	XYS	C1-C2-C3	3.28	117.12	110.31
2	A	706	XYP	C5-C4-C3	-3.22	105.70	109.67
2	D	706	XYP	C5-O5-C1	-3.12	107.47	112.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 31 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	706	XYP	1	0
2	C	705	XYP	2	0
3	D	709	XYS	3	0
2	D	704	XYP	2	0
3	C	707	XYS	2	0
3	A	709	XYS	4	0
2	A	707	XYP	7	0
2	D	703	XYP	1	0
2	A	705	XYP	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	701	XYP	7	0
2	A	704	XYP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	638/638 (100%)	-0.32	16 (2%) 57 61	6, 12, 26, 44	0
1	B	638/638 (100%)	-0.39	12 (1%) 66 70	7, 11, 25, 42	0
1	C	638/638 (100%)	-0.28	16 (2%) 57 61	7, 13, 26, 43	0
1	D	638/638 (100%)	-0.33	13 (2%) 65 69	7, 12, 24, 44	0
All	All	2552/2552 (100%)	-0.33	57 (2%) 62 66	6, 12, 25, 44	0

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	117	VAL	7.2
1	A	117	VAL	6.2
1	C	117	VAL	5.8
1	B	118	ASN	5.5
1	A	118	ASN	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	XYS	A	709	10/10	0.59	0.34	37,42,44,45	0
3	XYS	C	708	10/10	0.65	0.30	46,47,48,49	0
2	XYP	C	704	10/10	0.67	0.21	34,38,40,40	0
2	XYP	D	708	10/10	0.71	0.21	34,37,39,41	0
3	XYS	D	709	10/10	0.71	0.24	29,33,34,34	0
2	XYP	B	706	10/10	0.73	0.23	25,32,34,37	0
2	XYP	B	705	10/10	0.74	0.24	34,40,41,41	0
3	XYS	C	707	10/10	0.74	0.25	32,35,36,36	0
2	XYP	D	706	10/10	0.75	0.25	24,29,33,36	0
2	XYP	B	702	10/10	0.75	0.22	33,34,37,38	0
2	XYP	D	704	10/10	0.76	0.31	51,52,53,53	0
2	XYP	C	703	10/10	0.77	0.18	31,37,38,39	0
2	XYP	A	704	10/10	0.77	0.21	39,43,45,45	0
2	XYP	A	702	10/10	0.77	0.21	40,42,44,44	0
2	XYP	D	705	10/10	0.78	0.20	32,37,39,40	0
2	XYP	D	707	10/10	0.78	0.20	35,40,42,42	0
3	XYS	B	707	10/10	0.78	0.20	26,29,30,30	0
2	XYP	C	702	10/10	0.80	0.21	39,41,42,42	0
2	XYP	C	705	10/10	0.81	0.26	25,33,35,39	0
2	XYP	D	703	10/10	0.81	0.19	33,35,37,37	0
2	XYP	A	703	10/10	0.82	0.26	41,43,43,44	0
2	XYP	A	705	10/10	0.82	0.26	24,32,34,37	0
2	XYP	A	706	10/10	0.82	0.20	29,33,36,36	0
2	XYP	A	707	10/10	0.84	0.17	16,19,21,24	0
2	XYP	D	701	10/10	0.84	0.15	16,19,21,22	0
3	XYS	C	706	10/10	0.84	0.21	34,37,38,38	0
3	XYS	D	710	10/10	0.84	0.21	34,37,38,39	0
2	XYP	B	703	10/10	0.85	0.30	42,43,44,44	0
2	XYP	B	704	10/10	0.85	0.18	31,35,36,37	0
3	XYS	A	708	10/10	0.88	0.18	32,34,36,36	0
3	XYS	B	708	10/10	0.90	0.17	24,27,28,28	0
2	XYP	C	701	10/10	0.94	0.11	10,14,20,21	0
2	XYP	B	701	10/10	0.94	0.10	10,14,20,20	0
2	XYP	A	701	10/10	0.94	0.10	10,14,20,21	0
2	XYP	D	702	10/10	0.95	0.09	10,13,19,20	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.