



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 07:05 am BST

PDB ID : 3WGD
Title : Crystal structure of ERp46 Trx1
Authors : Inaba, K.; Suzuki, M.; Kojima, R.
Deposited on : 2013-08-04
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	113	
1	B	113	
1	C	113	
1	D	113	
1	E	113	
1	F	113	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	113	 4% 88% 8% ••
1	H	113	 4% 92% 5% •
1	I	113	 7% 90% 7% ••

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8449 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thioredoxin domain-containing protein 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	111	892	567	154	162	9	0	0	0
1	B	111	892	567	154	162	9	0	0	0
1	C	112	898	570	155	164	9	0	0	0
1	D	111	892	567	154	162	9	0	0	0
1	E	111	892	567	154	162	9	0	0	0
1	F	109	881	562	153	157	9	0	1	0
1	G	109	879	560	152	158	9	0	0	0
1	H	110	882	561	151	161	9	0	0	0
1	I	111	892	567	154	162	9	0	0	0

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	58	GLY	-	EXPRESSION TAG	UNP Q8NBS9
A	59	SER	-	EXPRESSION TAG	UNP Q8NBS9
A	60	HIS	-	EXPRESSION TAG	UNP Q8NBS9
A	61	MET	-	EXPRESSION TAG	UNP Q8NBS9
B	58	GLY	-	EXPRESSION TAG	UNP Q8NBS9
B	59	SER	-	EXPRESSION TAG	UNP Q8NBS9
B	60	HIS	-	EXPRESSION TAG	UNP Q8NBS9
B	61	MET	-	EXPRESSION TAG	UNP Q8NBS9
C	58	GLY	-	EXPRESSION TAG	UNP Q8NBS9
C	59	SER	-	EXPRESSION TAG	UNP Q8NBS9
C	60	HIS	-	EXPRESSION TAG	UNP Q8NBS9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	61	MET	-	EXPRESSION TAG	UNP Q8NBS9
D	58	GLY	-	EXPRESSION TAG	UNP Q8NBS9
D	59	SER	-	EXPRESSION TAG	UNP Q8NBS9
D	60	HIS	-	EXPRESSION TAG	UNP Q8NBS9
D	61	MET	-	EXPRESSION TAG	UNP Q8NBS9
E	58	GLY	-	EXPRESSION TAG	UNP Q8NBS9
E	59	SER	-	EXPRESSION TAG	UNP Q8NBS9
E	60	HIS	-	EXPRESSION TAG	UNP Q8NBS9
E	61	MET	-	EXPRESSION TAG	UNP Q8NBS9
F	58	GLY	-	EXPRESSION TAG	UNP Q8NBS9
F	59	SER	-	EXPRESSION TAG	UNP Q8NBS9
F	60	HIS	-	EXPRESSION TAG	UNP Q8NBS9
F	61	MET	-	EXPRESSION TAG	UNP Q8NBS9
G	58	GLY	-	EXPRESSION TAG	UNP Q8NBS9
G	59	SER	-	EXPRESSION TAG	UNP Q8NBS9
G	60	HIS	-	EXPRESSION TAG	UNP Q8NBS9
G	61	MET	-	EXPRESSION TAG	UNP Q8NBS9
H	58	GLY	-	EXPRESSION TAG	UNP Q8NBS9
H	59	SER	-	EXPRESSION TAG	UNP Q8NBS9
H	60	HIS	-	EXPRESSION TAG	UNP Q8NBS9
H	61	MET	-	EXPRESSION TAG	UNP Q8NBS9
I	58	GLY	-	EXPRESSION TAG	UNP Q8NBS9
I	59	SER	-	EXPRESSION TAG	UNP Q8NBS9
I	60	HIS	-	EXPRESSION TAG	UNP Q8NBS9
I	61	MET	-	EXPRESSION TAG	UNP Q8NBS9

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	G	1	Total	O	P	0	0
			5	4	1		
2	G	1	Total	O	P	0	0
			5	4	1		
2	H	1	Total	O	P	0	0
			5	4	1		
2	I	1	Total	O	P	0	0
			5	4	1		
2	I	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	E	1	Total C O 6 3 3	0	0
3	F	1	Total C O 6 3 3	0	0
3	G	1	Total C O 6 3 3	0	0
3	H	1	Total C O 6 3 3	0	0
3	I	1	Total C O 6 3 3	0	0

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	I	1	Total K 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	2	Total K 2 2	0	0
4	F	1	Total K 1 1	0	0
4	E	1	Total K 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	53	Total O 54 54	0	1
5	B	31	Total O 31 31	0	0
5	C	56	Total O 56 56	0	0
5	D	25	Total O 25 25	0	0
5	E	18	Total O 18 18	0	0
5	F	53	Total O 53 53	0	0
5	G	22	Total O 22 22	0	0
5	H	16	Total O 16 16	0	0
5	I	28	Total O 28 28	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thioredoxin domain-containing protein 5




- Molecule 1: Thioredoxin domain-containing protein 5





- Molecule 1: Thioredoxin domain-containing protein 5




- Molecule 1: Thioredoxin domain-containing protein 5



- Molecule 1: Thioredoxin domain-containing protein 5




- Molecule 1: Thioredoxin domain-containing protein 5

Chain F:  85% 12% .




- Molecule 1: Thioredoxin domain-containing protein 5

Chain G:  4% 88% 8% ..




- Molecule 1: Thioredoxin domain-containing protein 5

Chain H:  4% 92% 5% .



- Molecule 1: Thioredoxin domain-containing protein 5

Chain I:  7% 90% 7% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.44Å 94.68Å 142.02Å 90.00° 90.85° 90.00°	Depositor
Resolution (Å)	34.77 – 2.50 34.77 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.2 (34.77-2.50) 98.3 (34.77-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.51Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.188 , 0.235 0.188 , 0.233	Depositor DCC
R_{free} test set	2305 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	27.5	Xtrriage
Anisotropy	0.066	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.034 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8449	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7070e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, K, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/919	0.39	0/1245
1	B	0.25	0/919	0.40	0/1245
1	C	0.24	0/925	0.41	0/1253
1	D	0.24	0/919	0.39	0/1245
1	E	0.23	0/919	0.38	0/1245
1	F	0.25	0/910	0.40	0/1231
1	G	0.23	0/905	0.40	0/1224
1	H	0.23	0/908	0.40	0/1230
1	I	0.23	0/919	0.37	0/1245
All	All	0.24	0/8243	0.39	0/11163

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	892	0	838	6	0
1	B	892	0	839	5	0
1	C	898	0	845	2	0
1	D	892	0	838	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	892	0	839	4	0
1	F	881	0	836	8	0
1	G	879	0	830	5	0
1	H	882	0	833	3	0
1	I	892	0	840	6	0
2	A	20	0	0	0	0
2	B	5	0	0	1	0
2	C	10	0	0	0	0
2	D	10	0	0	0	0
2	E	5	0	0	0	0
2	G	10	0	0	1	0
2	H	5	0	0	0	0
2	I	10	0	0	0	0
3	A	18	0	24	0	0
3	B	12	0	16	0	0
3	C	6	0	8	0	0
3	E	6	0	8	1	0
3	F	6	0	8	0	0
3	G	6	0	8	1	0
3	H	6	0	8	0	0
3	I	6	0	8	3	0
4	C	2	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	I	1	0	0	0	0
5	A	54	0	0	1	0
5	B	31	0	0	0	0
5	C	56	0	0	0	0
5	D	25	0	0	1	0
5	E	18	0	0	0	0
5	F	53	0	0	2	0
5	G	22	0	0	0	0
5	H	16	0	0	0	0
5	I	28	0	0	0	0
All	All	8449	0	7626	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:VAL:HB	1:A:168:LEU:HD11	1.69	0.74
1:A:154:PRO:HG2	1:A:159:THR:HG21	1.69	0.72
1:F:76[A]:GLN:NE2	5:F:339:HOH:O	2.23	0.71
1:D:154:PRO:HG2	1:D:159:THR:HG21	1.73	0.70
1:H:114:VAL:HB	1:H:168:LEU:HD11	1.78	0.66
1:F:112:ALA:N	5:F:342:HOH:O	2.31	0.64
1:I:67:THR:HG22	3:I:203:GOL:H11	1.81	0.63
1:F:113:LYS:HD2	1:F:113:LYS:H	1.64	0.62
1:D:60:HIS:N	5:D:309:HOH:O	2.33	0.61
1:I:124:HIS:HE1	3:I:203:GOL:H12	1.68	0.59
1:I:124:HIS:CE1	3:I:203:GOL:H12	2.38	0.58
1:G:67:THR:HG22	3:G:203:GOL:H12	1.85	0.58
1:F:109:MET:HG2	1:F:112:ALA:HB2	1.85	0.58
1:D:110:GLU:CD	1:D:110:GLU:H	2.09	0.55
1:F:154:PRO:HG2	1:F:159:THR:HG21	1.90	0.53
1:H:110:GLU:H	1:H:110:GLU:CD	2.13	0.52
1:G:92:CYS:O	1:G:96:GLN:HG2	2.11	0.51
1:A:92:CYS:O	1:A:96:GLN:HG2	2.12	0.49
1:E:110:GLU:CD	1:E:110:GLU:H	2.15	0.49
1:I:154:PRO:HB2	1:I:159:THR:HG21	1.94	0.49
1:E:92:CYS:O	1:E:96:GLN:HG2	2.13	0.48
1:D:114:VAL:HB	1:D:168:LEU:HD11	1.96	0.48
1:C:92:CYS:O	1:C:96:GLN:HG2	2.14	0.48
1:G:115:TYR:CE1	1:H:88:TRP:HB3	2.51	0.46
1:E:65:LEU:HB3	3:E:202:GOL:H11	1.99	0.45
1:B:124:HIS:NE2	2:B:201:PO4:O4	2.49	0.45
1:F:100:ASN:HD21	1:F:118:LYS:NZ	2.15	0.45
1:I:60:HIS:HB2	1:I:100:ASN:ND2	2.32	0.45
1:B:68:ALA:HB2	1:B:124:HIS:CD2	2.53	0.43
1:C:126:ASP:OD1	1:C:126:ASP:N	2.51	0.43
1:B:158:GLN:O	1:B:162:ASN:ND2	2.51	0.43
1:D:92:CYS:O	1:D:96:GLN:HG2	2.19	0.43
1:A:60:HIS:N	5:A:327:HOH:O	2.53	0.42
1:B:92:CYS:O	1:B:96:GLN:HG2	2.19	0.42
1:G:60:HIS:CD2	1:G:100:ASN:HD22	2.38	0.42
1:A:88:TRP:HB3	1:F:115:TYR:CE1	2.56	0.41
1:B:86:ALA:HA	1:B:87:PRO:HD3	1.95	0.41
1:G:94:ARG:NH2	2:G:201:PO4:O3	2.54	0.41
1:I:143:LYS:HA	1:I:144:PRO:HD3	1.97	0.41
1:A:103:GLY:HA2	1:A:116:VAL:HB	2.03	0.41
1:F:92:CYS:O	1:F:96:GLN:HG2	2.21	0.41
1:E:126:ASP:N	1:E:126:ASP:OD1	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/113 (96%)	108 (99%)	1 (1%)	0	100	100
1	B	109/113 (96%)	107 (98%)	2 (2%)	0	100	100
1	C	110/113 (97%)	109 (99%)	1 (1%)	0	100	100
1	D	109/113 (96%)	107 (98%)	2 (2%)	0	100	100
1	E	109/113 (96%)	106 (97%)	3 (3%)	0	100	100
1	F	106/113 (94%)	105 (99%)	1 (1%)	0	100	100
1	G	105/113 (93%)	103 (98%)	2 (2%)	0	100	100
1	H	108/113 (96%)	105 (97%)	3 (3%)	0	100	100
1	I	109/113 (96%)	108 (99%)	1 (1%)	0	100	100
All	All	974/1017 (96%)	958 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/96 (99%)	92 (97%)	3 (3%)	39	65
1	B	95/96 (99%)	90 (95%)	5 (5%)	22	43
1	C	96/96 (100%)	94 (98%)	2 (2%)	53	78

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	95/96 (99%)	94 (99%)	1 (1%)	73	89
1	E	95/96 (99%)	92 (97%)	3 (3%)	39	65
1	F	94/96 (98%)	92 (98%)	2 (2%)	53	78
1	G	94/96 (98%)	90 (96%)	4 (4%)	29	53
1	H	94/96 (98%)	92 (98%)	2 (2%)	53	78
1	I	95/96 (99%)	93 (98%)	2 (2%)	53	78
All	All	853/864 (99%)	829 (97%)	24 (3%)	43	70

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	HIS
1	A	62	SER
1	A	81	PHE
1	B	65	LEU
1	B	81	PHE
1	B	91	HIS
1	B	102	LEU
1	B	150	LYS
1	C	81	PHE
1	C	170	GLU
1	D	81	PHE
1	E	81	PHE
1	E	102	LEU
1	E	126	ASP
1	F	81	PHE
1	F	152	GLN
1	G	60	HIS
1	G	65	LEU
1	G	81	PHE
1	G	102	LEU
1	H	81	PHE
1	H	147	GLU
1	I	60	HIS
1	I	81	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN
1	B	76	GLN
1	B	93	GLN
1	B	166	GLN
1	C	131	GLN
1	E	80	HIS
1	E	124	HIS
1	E	146	GLN
1	F	100	ASN
1	F	131	GLN
1	G	60	HIS
1	I	64	HIS
1	I	162	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 31 ligands modelled in this entry, 5 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	204	-	4,4,4	0.90	0	6,6,6	0.45	0
2	PO4	A	203	-	4,4,4	0.90	0	6,6,6	0.45	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	D	201	-	4,4,4	0.94	0	6,6,6	0.46	0
3	GOL	E	202	-	5,5,5	0.36	0	5,5,5	0.30	0
3	GOL	H	202	-	5,5,5	0.34	0	5,5,5	0.37	0
2	PO4	D	202	-	4,4,4	0.89	0	6,6,6	0.48	0
3	GOL	A	205	-	5,5,5	0.35	0	5,5,5	0.40	0
2	PO4	G	201	-	4,4,4	0.91	0	6,6,6	0.39	0
3	GOL	B	202	-	5,5,5	0.39	0	5,5,5	0.18	0
3	GOL	A	207	-	5,5,5	0.37	0	5,5,5	0.32	0
2	PO4	G	202	-	4,4,4	0.91	0	6,6,6	0.43	0
2	PO4	E	201	-	4,4,4	0.93	0	6,6,6	0.46	0
2	PO4	I	201	-	4,4,4	0.91	0	6,6,6	0.45	0
2	PO4	H	201	-	4,4,4	0.91	0	6,6,6	0.46	0
2	PO4	I	202	-	4,4,4	0.88	0	6,6,6	0.43	0
3	GOL	G	203	-	5,5,5	0.36	0	5,5,5	0.33	0
2	PO4	B	201	-	4,4,4	0.94	0	6,6,6	0.46	0
2	PO4	A	201	-	4,4,4	0.93	0	6,6,6	0.46	0
2	PO4	C	201	-	4,4,4	0.94	0	6,6,6	0.48	0
2	PO4	A	202	-	4,4,4	0.91	0	6,6,6	0.57	0
3	GOL	C	203	-	5,5,5	0.36	0	5,5,5	0.29	0
3	GOL	B	203	-	5,5,5	0.36	0	5,5,5	0.31	0
3	GOL	A	206	-	5,5,5	0.36	0	5,5,5	0.27	0
3	GOL	I	203	-	5,5,5	0.34	0	5,5,5	0.41	0
2	PO4	C	202	-	4,4,4	0.90	0	6,6,6	0.46	0
3	GOL	F	201	-	5,5,5	0.38	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	203	-	-	2/4/4/4	-
3	GOL	A	207	-	-	0/4/4/4	-
3	GOL	B	203	-	-	4/4/4/4	-
3	GOL	G	203	-	-	3/4/4/4	-
3	GOL	E	202	-	-	2/4/4/4	-
3	GOL	H	202	-	-	2/4/4/4	-
3	GOL	A	206	-	-	0/4/4/4	-
3	GOL	A	205	-	-	2/4/4/4	-
3	GOL	F	201	-	-	4/4/4/4	-
3	GOL	I	203	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	202	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	202	GOL	O1-C1-C2-C3
3	G	203	GOL	O1-C1-C2-C3
3	C	203	GOL	O1-C1-C2-C3
3	B	203	GOL	C1-C2-C3-O3
3	B	203	GOL	O2-C2-C3-O3
3	I	203	GOL	O1-C1-C2-C3
3	F	201	GOL	C1-C2-C3-O3
3	E	202	GOL	O1-C1-C2-O2
3	B	203	GOL	O1-C1-C2-O2
3	E	202	GOL	O1-C1-C2-C3
3	H	202	GOL	O1-C1-C2-C3
3	A	205	GOL	O1-C1-C2-C3
3	B	203	GOL	O1-C1-C2-C3
3	F	201	GOL	O1-C1-C2-C3
3	H	202	GOL	O1-C1-C2-O2
3	B	202	GOL	O1-C1-C2-O2
3	G	203	GOL	O1-C1-C2-O2
3	C	203	GOL	O1-C1-C2-O2
3	I	203	GOL	O1-C1-C2-O2
3	F	201	GOL	O2-C2-C3-O3
3	F	201	GOL	O1-C1-C2-O2
3	A	205	GOL	O1-C1-C2-O2
3	G	203	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	202	GOL	1	0
2	G	201	PO4	1	0
3	G	203	GOL	1	0
2	B	201	PO4	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	203	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	111/113 (98%)	-0.28	1 (0%) 84 86	22, 32, 62, 75	0
1	B	111/113 (98%)	-0.22	1 (0%) 84 86	26, 39, 69, 106	0
1	C	112/113 (99%)	-0.24	1 (0%) 84 86	24, 30, 54, 73	0
1	D	111/113 (98%)	-0.15	1 (0%) 84 86	29, 48, 78, 86	0
1	E	111/113 (98%)	0.17	2 (1%) 68 71	29, 53, 85, 103	0
1	F	109/113 (96%)	-0.33	0 100 100	21, 32, 53, 81	0
1	G	109/113 (96%)	-0.10	5 (4%) 32 34	30, 47, 73, 80	0
1	H	110/113 (97%)	0.13	5 (4%) 33 36	35, 59, 85, 106	0
1	I	111/113 (98%)	0.14	8 (7%) 15 16	30, 49, 75, 81	0
All	All	995/1017 (97%)	-0.10	24 (2%) 59 62	21, 43, 77, 106	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	145	GLY	4.0
1	H	111	ASP	3.8
1	H	145	GLY	3.7
1	B	60	HIS	3.6
1	I	152	GLN	3.5
1	C	147	GLU	3.0
1	I	110	GLU	2.8
1	E	97	PRO	2.8
1	I	153	GLY	2.8
1	G	146	GLN	2.8
1	I	60	HIS	2.7
1	H	147	GLU	2.6
1	G	60	HIS	2.6
1	I	154	PRO	2.4
1	A	147	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	I	97	PRO	2.3
1	H	155	ARG	2.2
1	H	169	ASN	2.1
1	E	111	ASP	2.1
1	I	151	TYR	2.1
1	D	147	GLU	2.1
1	G	144	PRO	2.1
1	G	147	GLU	2.1
1	I	169	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	206	6/6	0.75	0.27	73,74,74,74	0
3	GOL	B	202	6/6	0.81	0.33	75,76,76,76	0
2	PO4	D	202	5/5	0.83	0.20	107,107,107,108	0
2	PO4	A	204	5/5	0.87	0.33	105,105,105,105	0
3	GOL	A	207	6/6	0.88	0.17	57,57,59,60	0
2	PO4	A	203	5/5	0.89	0.18	94,94,94,94	0
3	GOL	E	202	6/6	0.89	0.19	42,43,44,45	0
3	GOL	I	203	6/6	0.90	0.21	54,55,56,56	0
2	PO4	C	202	5/5	0.91	0.23	89,90,90,90	0
3	GOL	B	203	6/6	0.92	0.21	46,48,48,48	0
3	GOL	G	203	6/6	0.92	0.17	51,52,52,53	0
2	PO4	G	201	5/5	0.93	0.15	78,78,78,79	0
4	K	C	205	1/1	0.93	0.10	67,67,67,67	0
3	GOL	C	203	6/6	0.94	0.16	34,35,37,37	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	F	201	6/6	0.94	0.17	45,47,49,49	0
2	PO4	G	202	5/5	0.95	0.14	53,53,54,55	0
2	PO4	I	201	5/5	0.95	0.18	72,73,73,73	0
2	PO4	H	201	5/5	0.95	0.16	62,63,63,64	0
2	PO4	I	202	5/5	0.95	0.14	67,67,68,69	0
3	GOL	A	205	6/6	0.95	0.15	40,43,45,46	0
3	GOL	H	202	6/6	0.95	0.25	40,43,43,44	0
2	PO4	E	201	5/5	0.96	0.20	64,65,65,65	0
4	K	I	204	1/1	0.97	0.15	73,73,73,73	0
4	K	E	203	1/1	0.97	0.10	64,64,64,64	0
4	K	C	204	1/1	0.97	0.11	63,63,63,63	0
2	PO4	C	201	5/5	0.97	0.14	53,53,53,54	0
2	PO4	A	202	5/5	0.98	0.16	39,39,40,41	0
2	PO4	A	201	5/5	0.98	0.15	39,40,41,42	0
2	PO4	D	201	5/5	0.98	0.13	56,56,58,58	0
2	PO4	B	201	5/5	0.99	0.12	53,53,53,54	0
4	K	F	202	1/1	0.99	0.18	57,57,57,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.