



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 17, 2022 – 07:20 PM EDT

PDB ID : 3WW1  
Title : X-ray structure of Cellulomonas parahominis L-ribose isomerase with L-ribose  
Authors : Terami, Y.; Yoshida, H.; Takata, G.; Kamitori, S.  
Deposited on : 2014-06-13  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.29  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	
1	B	256	

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-ribose isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	242	Total	C	N	O	S	0	0	0
			1882	1193	323	357	9			
1	B	242	Total	C	N	O	S	0	0	0
			1882	1193	323	357	9			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	HIS	-	expression tag	UNP L0N3Y0
A	-5	HIS	-	expression tag	UNP L0N3Y0
A	-4	HIS	-	expression tag	UNP L0N3Y0
A	-3	HIS	-	expression tag	UNP L0N3Y0
A	-2	HIS	-	expression tag	UNP L0N3Y0
A	-1	HIS	-	expression tag	UNP L0N3Y0
A	0	GLY	-	expression tag	UNP L0N3Y0
A	1	SER	-	expression tag	UNP L0N3Y0
A	119	LEU	PHE	engineered mutation	UNP L0N3Y0
A	125	PHE	LEU	engineered mutation	UNP L0N3Y0
B	-6	HIS	-	expression tag	UNP L0N3Y0
B	-5	HIS	-	expression tag	UNP L0N3Y0
B	-4	HIS	-	expression tag	UNP L0N3Y0
B	-3	HIS	-	expression tag	UNP L0N3Y0
B	-2	HIS	-	expression tag	UNP L0N3Y0
B	-1	HIS	-	expression tag	UNP L0N3Y0
B	0	GLY	-	expression tag	UNP L0N3Y0
B	1	SER	-	expression tag	UNP L0N3Y0
B	119	LEU	PHE	engineered mutation	UNP L0N3Y0
B	125	PHE	LEU	engineered mutation	UNP L0N3Y0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mn 1 1	0	0
2	B	1	Total Mn 1 1	0	0

- Molecule 3 is beta-L-ribofuranose (three-letter code: 0MK) (formula: C<sub>5</sub>H<sub>10</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 10 5 5	0	0
3	A	1	Total C O 10 5 5	0	0
3	B	1	Total C O 10 5 5	0	0

- Molecule 4 is L-ribose (three-letter code: ROR) (formula: C<sub>5</sub>H<sub>10</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	C O	0	0
			10	5 5		


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	120	Total	O	0	0
			120	120		
5	B	90	Total	O	0	0
			90	90		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-ribose isomerase

Chain A: 



- Molecule 1: L-ribose isomerase

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.73Å 88.55Å 152.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.33 – 1.95 19.33 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.5 (19.33-1.95) 98.6 (19.33-1.95)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 1.94Å)	Xtrriage
Refinement program	CNS 1.3	Depositor
R, $R_{free}$	0.202 , 0.242 0.193 , 0.234	Depositor DCC
$R_{free}$ test set	3758 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.3	Xtrriage
Anisotropy	0.153	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 50.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4016	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0MK, MN, ROR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/1939	0.64	1/2646 (0.0%)
1	B	0.33	0/1939	0.62	1/2646 (0.0%)
All	All	0.33	0/3878	0.63	2/5292 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	103	CYS	N-CA-C	-7.50	90.74	111.00
1	B	103	CYS	N-CA-C	-6.68	92.97	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1882	0	1790	35	0
1	B	1882	0	1790	35	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	20	0	20	0	0
3	B	10	0	10	0	0
4	B	10	0	8	1	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	120	0	0	0	0
5	B	90	0	0	2	0
All	All	4016	0	3618	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:VAL:CG1	1:A:96:ILE:HB	2.09	0.82
1:B:239:ALA:O	1:B:243:ARG:HB2	1.83	0.79
1:A:64:VAL:HG13	1:A:96:ILE:HB	1.65	0.77
1:A:243:ARG:O	1:A:243:ARG:HD3	1.85	0.77
1:A:160:ARG:HH12	1:A:239:ALA:HB2	1.53	0.73
1:A:121:GLU:HG2	1:A:174:ARG:HG2	1.72	0.72
1:A:212:PRO:HG3	1:A:230:HIS:ND1	2.09	0.67
1:A:9:ARG:O	1:A:13:GLU:HG3	1.97	0.65
1:B:57:ARG:HD3	1:B:194:ALA:O	1.97	0.63
1:A:77:LEU:HD22	1:A:78:PRO:HD2	1.79	0.63
1:A:160:ARG:NH1	1:A:239:ALA:HB2	2.15	0.62
1:A:141:PHE:CD1	1:B:222:PRO:HA	2.35	0.61
1:B:40:ILE:O	1:B:62:VAL:HG23	2.00	0.61
1:A:8:ARG:HH21	1:A:8:ARG:HG3	1.64	0.60
1:A:15:LEU:HD21	1:A:38:ALA:HB3	1.83	0.60
1:A:61:GLU:OE1	1:A:160:ARG:NH1	2.35	0.59
1:B:7:SER:OG	1:B:10:GLU:HG3	2.04	0.57
1:B:109:GLN:HG3	1:B:133:VAL:HG21	1.86	0.57
1:B:9:ARG:O	1:B:13:GLU:HG3	2.04	0.57
1:B:61:GLU:OE1	1:B:160:ARG:NH1	2.37	0.56
1:A:111:LYS:HE3	1:A:113:GLU:OE2	2.07	0.55
1:B:167:LEU:N	1:B:167:LEU:HD12	2.22	0.55
1:A:226:TRP:HA	1:A:229:LEU:HD22	1.89	0.54
1:A:91:CYS:SG	1:A:93:LYS:HE3	2.48	0.54
1:A:167:LEU:HD12	1:A:167:LEU:N	2.24	0.53
1:B:159:GLY:N	1:B:161:GLU:OE1	2.42	0.53
1:B:212:PRO:HG2	1:B:230:HIS:CE1	2.44	0.52
1:A:221:ALA:CB	1:A:227:ARG:HG2	2.41	0.51
1:A:8:ARG:HG3	1:A:8:ARG:NH2	2.26	0.50
1:A:102:PHE:CD2	1:A:191:ARG:HB3	2.46	0.50
1:B:95:MET:HB2	1:B:202:VAL:HG13	1.94	0.50

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:PHE:CD2	1:B:191:ARG:HB3	2.48	0.49
1:B:57:ARG:NH1	5:B:513:HOH:O	2.45	0.49
1:B:107:PHE:HB3	1:B:187:LEU:HD23	1.94	0.49
1:B:243:ARG:C	1:B:243:ARG:HD3	2.33	0.48
1:A:234:PHE:CD1	1:A:240:ASN:HA	2.49	0.48
1:B:11:TYR:CE2	1:B:15:LEU:HD22	2.50	0.47
1:A:221:ALA:HB2	1:A:227:ARG:HG2	1.96	0.46
1:A:61:GLU:CD	1:A:104:PRO:HB3	2.35	0.46
1:A:107:PHE:CD1	1:A:107:PHE:C	2.89	0.46
1:A:42:PHE:HB3	1:A:239:ALA:HB1	1.97	0.45
1:B:25:LEU:O	1:B:26:ARG:HB2	2.16	0.45
1:B:211:GLU:O	1:B:213:THR:N	2.50	0.45
1:A:121:GLU:CG	1:A:174:ARG:HG2	2.44	0.45
1:A:243:ARG:HD3	1:A:243:ARG:C	2.37	0.44
1:A:69:LEU:HB2	1:B:179:LYS:HG3	1.99	0.44
1:B:244:LEU:O	1:B:245:ALA:C	2.56	0.44
1:B:107:PHE:CZ	1:B:233:THR:HG21	2.53	0.44
1:B:14:TRP:CE2	1:B:64:VAL:HG23	2.53	0.43
1:B:37:SER:HB3	1:B:213:THR:CG2	2.48	0.43
1:B:233:THR:HG22	5:B:562:HOH:O	2.17	0.43
1:A:77:LEU:CD2	1:A:78:PRO:HD2	2.46	0.43
1:B:18:ALA:HB1	1:B:96:ILE:HD13	2.01	0.42
1:A:30:THR:OG1	1:A:32:GLU:OE2	2.37	0.42
1:A:214:GLU:OE2	1:A:214:GLU:N	2.44	0.42
1:A:123:GLU:O	1:A:190:PHE:HA	2.20	0.41
1:B:167:LEU:N	1:B:167:LEU:CD1	2.82	0.41
1:B:104:PRO:HA	1:B:105:PRO:HD3	1.91	0.41
1:B:160:ARG:HH12	1:B:239:ALA:HB2	1.85	0.41
1:A:25:LEU:O	1:A:26:ARG:HB2	2.20	0.41
1:B:243:ARG:HD3	1:B:243:ARG:O	2.21	0.41
1:A:167:LEU:N	1:A:167:LEU:CD1	2.84	0.41
1:B:243:ARG:HH11	4:B:402:ROR:C5	2.33	0.41
1:B:14:TRP:CH2	1:B:62:VAL:CG2	3.04	0.41
1:A:124:VAL:HB	1:A:171:VAL:HG12	2.03	0.40
1:B:26:ARG:HG2	1:B:26:ARG:HH11	1.87	0.40
1:B:160:ARG:NH1	1:B:239:ALA:HB2	2.36	0.40
1:B:172:ARG:HG2	1:B:174:ARG:NH1	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/256 (93%)	230 (97%)	7 (3%)	1 (0%)	34	22
1	B	238/256 (93%)	224 (94%)	8 (3%)	6 (2%)	5	1
All	All	476/512 (93%)	454 (95%)	15 (3%)	7 (2%)	10	3

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	133	VAL
1	B	244	LEU
1	B	245	ALA
1	A	244	LEU
1	B	135	ASP
1	B	136	ASP
1	B	212	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/209 (95%)	192 (97%)	6 (3%)	41	30
1	B	198/209 (95%)	191 (96%)	7 (4%)	36	24
All	All	396/418 (95%)	383 (97%)	13 (3%)	38	26

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ARG
1	A	54	LEU
1	A	62	VAL
1	A	77	LEU
1	A	240	ASN
1	A	243	ARG
1	B	54	LEU
1	B	136	ASP
1	B	162	ASP
1	B	214	GLU
1	B	233	THR
1	B	237	GLU
1	B	244	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	240	ASN
1	B	52	HIS
1	B	230	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	OMK	B	403	-	10,10,10	0.64	0	14,14,14	0.41	0
3	OMK	A	403	-	10,10,10	0.68	0	14,14,14	0.44	0
4	ROR	B	402	2	8,9,9	0.44	0	10,11,11	0.59	0
3	OMK	A	402	-	10,10,10	0.67	0	14,14,14	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OMK	B	403	-	-	-	0/1/1/1
3	OMK	A	403	-	-	-	0/1/1/1
3	OMK	A	402	-	-	-	0/1/1/1
4	ROR	B	402	2	-	6/10/12/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	402	ROR	O4-C4-C5-O5
4	B	402	ROR	O2-C2-C3-C4
4	B	402	ROR	C1-C2-C3-C4
4	B	402	ROR	O2-C2-C3-O3
4	B	402	ROR	C3-C4-C5-O5
4	B	402	ROR	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	402	ROR	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	242/256 (94%)	0.47	24 (9%) <b>7</b> <b>12</b>	22, 30, 59, 82	0
1	B	242/256 (94%)	0.47	23 (9%) <b>8</b> <b>13</b>	22, 32, 62, 82	0
All	All	484/512 (94%)	0.47	47 (9%) <b>7</b> <b>12</b>	22, 31, 62, 82	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	242	GLY	8.5
1	A	213	THR	7.5
1	B	213	THR	7.4
1	B	135	ASP	7.2
1	B	241	SER	7.1
1	A	244	LEU	7.0
1	A	214	GLU	6.7
1	A	241	SER	6.4
1	A	240	ASN	6.0
1	A	249	ALA	5.8
1	B	2	THR	5.6
1	B	136	ASP	5.4
1	B	214	GLU	5.2
1	A	135	ASP	5.2
1	A	2	THR	5.2
1	A	230	HIS	4.8
1	B	249	ALA	4.8
1	A	237	GLU	4.6
1	A	227	ARG	4.5
1	B	137	ASP	4.4
1	B	134	GLY	4.4
1	B	133	VAL	4.2
1	B	212	PRO	4.0
1	B	240	ASN	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	134	GLY	3.9
1	A	242	GLY	3.8
1	B	245	ALA	3.8
1	A	224	PRO	3.7
1	B	237	GLU	3.6
1	B	244	LEU	3.5
1	A	243	ARG	3.4
1	A	136	ASP	3.4
1	A	62	VAL	3.2
1	A	212	PRO	2.9
1	B	243	ARG	2.9
1	A	239	ALA	2.8
1	B	152	GLU	2.7
1	A	79	ALA	2.6
1	B	238	ALA	2.5
1	B	165	ALA	2.4
1	B	239	ALA	2.3
1	B	80	ALA	2.3
1	A	228	GLY	2.2
1	A	245	ALA	2.2
1	A	137	ASP	2.1
1	B	146	GLU	2.1
1	A	3	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	OMK	A	403	10/10	0.76	0.39	72,74,75,75	0
3	OMK	A	402	10/10	0.77	0.16	78,79,79,79	0
4	ROR	B	402	10/10	0.87	0.15	66,68,69,71	0
3	OMK	B	403	10/10	0.90	0.25	73,74,74,75	0
2	MN	A	401	1/1	0.99	0.06	40,40,40,40	0
2	MN	B	401	1/1	0.99	0.05	37,37,37,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.