



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 11:11 am BST

PDB ID : 3ZHQ  
Title : Crystal structure of the H747A mutant of the SucA domain of Mycobacterium smegmatis KGD  
Authors : Wagner, T.; Barilone, N.; Bellinzoni, M.; Alzari, P.M.  
Deposited on : 2012-12-24  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	868	 2% 88% 5% 7%
1	B	868	 4% 87% 6% 7%
1	C	868	 4% 88% 5% 6%
1	D	868	 4% 89% 5% 7%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 25781 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called MULTIFUNCTIONAL 2-OXOGLUTARATE METABOLISM ENZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	811	Total 6283	C 3959	N 1108	O 1194	S 22	0	1	0
1	B	808	Total 6208	C 3913	N 1103	O 1168	S 24	0	0	0
1	C	812	Total 6301	C 3971	N 1110	O 1197	S 23	0	0	0
1	D	810	Total 6234	C 3933	N 1097	O 1180	S 24	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	360	GLY	-	expression tag	UNP A0R2B1
A	747	ALA	HIS	engineered mutation	UNP A0R2B1
B	360	GLY	-	expression tag	UNP A0R2B1
B	747	ALA	HIS	engineered mutation	UNP A0R2B1
C	360	GLY	-	expression tag	UNP A0R2B1
C	747	ALA	HIS	engineered mutation	UNP A0R2B1
D	360	GLY	-	expression tag	UNP A0R2B1
D	747	ALA	HIS	engineered mutation	UNP A0R2B1

- Molecule 2 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: C<sub>12</sub>H<sub>19</sub>N<sub>4</sub>O<sub>7</sub>P<sub>2</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
2	B	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
2	C	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
2	D	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
4	B	1	Total	Ca	0	0
			1	1		

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0
4	D	1	Total Ca 1 1	0	0
4	C	1	Total Ca 1 1	0	0



- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	197	Total O 197 197	0	0
5	B	154	Total O 154 154	0	0
5	C	165	Total O 165 165	0	0
5	D	127	Total O 127 127	0	0





● Molecule 1: MULTIFUNCTIONAL 2-OXOGLUTARATE METABOLISM ENZYME



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.67Å 83.74Å 159.58Å 99.86° 98.95° 100.44°	Depositor
Resolution (Å)	48.97 – 2.50 48.97 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.8 (48.97-2.50) 96.7 (48.97-2.50)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 2.51Å)	Xtrriage
Refinement program	BUSTER 2.11.2	Depositor
R, $R_{free}$	0.194 , 0.223 0.203 , 0.238	Depositor DCC
$R_{free}$ test set	6718 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.9	Xtrriage
Anisotropy	0.401	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 54.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.012 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	25781	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MG, TPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/6411	0.65	0/8699
1	B	0.51	0/6335	0.65	0/8600
1	C	0.50	0/6427	0.65	0/8712
1	D	0.50	0/6359	0.65	0/8625
All	All	0.50	0/25532	0.65	0/34636

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6283	0	6051	23	0
1	B	6208	0	5963	27	0
1	C	6301	0	6108	27	0
1	D	6234	0	6017	23	0
2	A	26	0	16	2	0
2	B	26	0	16	2	0
2	C	26	0	16	2	0
2	D	26	0	16	2	0
3	A	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	197	0	0	2	0
5	B	154	0	0	1	0
5	C	165	0	0	0	0
5	D	127	0	0	4	0
All	All	25781	0	24203	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 99 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:705:PRO:HG2	1:C:735:VAL:HG13	1.69	0.73
1:C:373:LEU:CD1	1:C:436:LEU:HD21	2.20	0.71
1:C:692:SER:HB2	1:C:697:ASP:OD2	1.93	0.67
1:C:373:LEU:HD11	1:C:436:LEU:CD2	2.27	0.65
1:A:692:SER:HB2	1:A:697:ASP:OD2	1.98	0.63

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	802/868 (92%)	779 (97%)	20 (2%)	3 (0%)	<span style="border: 1px solid red; padding: 2px;">34</span> <span style="border: 1px solid gray; padding: 2px;">54</span>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	798/868 (92%)	779 (98%)	18 (2%)	1 (0%)	51	73
1	C	802/868 (92%)	779 (97%)	21 (3%)	2 (0%)	47	68
1	D	800/868 (92%)	777 (97%)	22 (3%)	1 (0%)	51	73
All	All	3202/3472 (92%)	3114 (97%)	81 (2%)	7 (0%)	47	68

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	605	HIS
1	A	605	HIS
1	A	636	SER
1	A	682	PHE
1	B	682	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	645/725 (89%)	637 (99%)	8 (1%)	71	88
1	B	629/725 (87%)	622 (99%)	7 (1%)	73	89
1	C	651/725 (90%)	644 (99%)	7 (1%)	73	89
1	D	637/725 (88%)	634 (100%)	3 (0%)	88	96
All	All	2562/2900 (88%)	2537 (99%)	25 (1%)	76	90

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	950	LEU
1	B	979	ASP
1	D	950	LEU
1	B	953	PHE
1	B	1187	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	464	GLN
1	D	464	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	TPP	A	2001	3	22,27,27	1.65	3 (13%)	29,40,40	1.87	9 (31%)
2	TPP	D	2001	3	22,27,27	2.05	3 (13%)	29,40,40	1.79	8 (27%)
2	TPP	B	2001	3	22,27,27	2.08	3 (13%)	29,40,40	1.81	8 (27%)
2	TPP	C	2001	3	22,27,27	2.13	4 (18%)	29,40,40	1.91	8 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPP	A	2001	3	-	4/16/17/17	0/2/2/2
2	TPP	D	2001	3	-	2/16/17/17	0/2/2/2
2	TPP	B	2001	3	-	4/16/17/17	0/2/2/2
2	TPP	C	2001	3	-	5/16/17/17	0/2/2/2

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2001	TPP	C4-N3	-7.49	1.33	1.39
2	C	2001	TPP	C4-N3	-7.38	1.33	1.39
2	B	2001	TPP	C4-N3	-7.34	1.33	1.39
2	A	2001	TPP	C4-N3	-5.16	1.35	1.39
2	B	2001	TPP	PB-O1B	3.66	1.62	1.50

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	TPP	CM4-C4-C5	-4.46	117.84	127.60
2	B	2001	TPP	CM4-C4-C5	-4.31	118.19	127.60
2	A	2001	TPP	C5-C4-N3	4.24	116.06	107.57
2	C	2001	TPP	CM4-C4-C5	-4.12	118.59	127.60
2	D	2001	TPP	CM4-C4-C5	-3.88	119.11	127.60

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	2001	TPP	C7-O7-PA-O1A
2	D	2001	TPP	C7-O7-PA-O1A
2	B	2001	TPP	C7-O7-PA-O1A
2	C	2001	TPP	C7-O7-PA-O1A
2	C	2001	TPP	C5-C6-C7-O7



There are no ring outliers.


4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	TPP	2	0
2	D	2001	TPP	2	0
2	B	2001	TPP	2	0
2	C	2001	TPP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	811/868 (93%)	0.01	20 (2%) 57 61	22, 39, 68, 97	0
1	B	808/868 (93%)	0.11	38 (4%) 31 33	20, 40, 75, 103	0
1	C	812/868 (93%)	0.15	32 (3%) 39 42	22, 42, 74, 98	0
1	D	810/868 (93%)	0.18	38 (4%) 31 33	23, 42, 75, 114	0
All	All	3241/3472 (93%)	0.11	128 (3%) 39 42	20, 41, 73, 114	0

The worst 5 of 128 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	368	ALA	5.9
1	B	472	ASP	5.8
1	C	779	ILE	5.4
1	B	810	LEU	5.4
1	D	472	ASP	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CA	C	2003	1/1	0.95	0.04	42,42,42,42	0
3	MG	A	2002	1/1	0.97	0.15	20,20,20,20	0
3	MG	D	2002	1/1	0.97	0.12	23,23,23,23	0
2	TPP	A	2001	26/26	0.97	0.15	16,26,33,34	0
2	TPP	D	2001	26/26	0.97	0.18	23,33,36,38	0
2	TPP	B	2001	26/26	0.98	0.16	20,31,37,38	0
2	TPP	C	2001	26/26	0.98	0.15	23,34,39,40	0
4	CA	B	2003	1/1	0.99	0.06	38,38,38,38	0
3	MG	B	2002	1/1	0.99	0.08	18,18,18,18	0
4	CA	D	2003	1/1	0.99	0.06	40,40,40,40	0
3	MG	C	2002	1/1	0.99	0.14	15,15,15,15	0
4	CA	A	2003	1/1	0.99	0.09	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around TPP D 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around TPP B 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.