



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 9, 2020 – 02:11 PM BST

PDB ID : 4HJ1
Title : Crystal structure of glycoprotein C from Rift Valley Fever Virus (glycosylated)
Authors : Dessau, M.; Modis, Y.
Deposited on : 2012-10-12
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	 6% 87% 11% .
1	B	432	 26% 82% 15% ..
1	C	432	 14% 85% 13% .
1	D	432	 8% 86% 13% .
2	E	4	 75% 25%
3	F	3	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	G	5	 100%
4	H	5	 60% 40%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BMA	G	5	-	-	-	X
7	GOL	D	1210	-	-	X	-

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 14382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ENVELOPE GLYCOPROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	431	Total 3279	C 2036	N 561	O 653	S 29	0	2	0
1	B	427	Total 3242	C 2011	N 557	O 645	S 29	0	0	0
1	C	430	Total 3271	C 2031	N 561	O 649	S 30	0	1	0
1	D	431	Total 3296	C 2046	N 562	O 658	S 30	0	5	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	688	ASP	-	expression tag	UNP A2T075
A	689	PRO	-	expression tag	UNP A2T075
A	690	GLY	-	expression tag	UNP A2T075
B	688	ASP	-	expression tag	UNP A2T075
B	689	PRO	-	expression tag	UNP A2T075
B	690	GLY	-	expression tag	UNP A2T075
C	688	ASP	-	expression tag	UNP A2T075
C	689	PRO	-	expression tag	UNP A2T075
C	690	GLY	-	expression tag	UNP A2T075
D	688	ASP	-	expression tag	UNP A2T075
D	689	PRO	-	expression tag	UNP A2T075
D	690	GLY	-	expression tag	UNP A2T075

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	4	50	28	2	20	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	3	39	22	2	15	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	5	61	34	2	25	0	0	0
4	H	5	61	34	2	25	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	Total 14	C 8	N 1	O 5	0	0
5	B	1	Total 14	C 8	N 1	O 5	0	0
5	C	1	Total 14	C 8	N 1	O 5	0	0
5	D	1	Total 14	C 8	N 1	O 5	0	0

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	332	Total 332	O 332	0	0
8	B	111	Total 111	O 111	0	0
8	C	170	Total 170	O 170	0	0
8	D	324	Total 324	O 324	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ENVELOPE GLYCOPROTEIN




- Molecule 1: ENVELOPE GLYCOPROTEIN





- Molecule 1: ENVELOPE GLYCOPROTEIN





- Molecule 1: ENVELOPE GLYCOPROTEIN




- Molecule 2: beta-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 4: beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:



- EMG1
- EMG2
- EMG3
- EMG4
- EMG5

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	128.68Å 56.37Å 140.19Å 90.00° 96.55° 90.00°	Depositor
Resolution (Å)	29.76 – 1.90 29.75 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.3 (29.76-1.90) 97.4 (29.75-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.214 , 0.255 0.215 , 0.257	Depositor DCC
R_{free} test set	7722 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	33.0	Xtrriage
Anisotropy	0.330	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14382	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.9715e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BMA, NAG, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/3345	0.69	1/4523 (0.0%)
1	B	0.39	1/3302 (0.0%)	0.56	0/4463
1	C	0.40	0/3331	0.59	0/4502
1	D	0.51	0/3368	0.66	2/4553 (0.0%)
All	All	0.45	1/13346 (0.0%)	0.63	3/18041 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1097	PHE	CG-CD2	-6.11	1.29	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1047	ARG	NE-CZ-NH2	-6.47	117.06	120.30
1	D	772	LEU	CA-CB-CG	5.33	127.55	115.30
1	D	881	ARG	NE-CZ-NH1	5.02	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3279	0	3155	39	0
1	B	3242	0	3111	39	0
1	C	3271	0	3145	36	0
1	D	3296	0	3168	52	0
2	E	50	0	43	1	0
3	F	39	0	34	0	0
4	G	61	0	52	0	0
4	H	61	0	52	3	0
5	A	14	0	13	0	0
5	B	14	0	13	0	0
5	C	14	0	13	0	0
5	D	14	0	13	0	0
6	A	12	0	13	1	0
6	B	12	0	13	0	0
6	C	12	0	13	0	0
6	D	24	0	26	3	0
7	A	6	0	8	0	0
7	B	6	0	8	0	0
7	C	6	0	8	0	0
7	D	12	0	16	10	0
8	A	332	0	0	12	0
8	B	111	0	0	1	0
8	C	170	0	0	7	0
8	D	324	0	0	11	0
All	All	14382	0	12917	166	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 166 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:912:GLY:HA3	7:D:1210:GOL:H11	1.41	1.02
1:B:1060:LEU:HA	1:B:1101:CYS:HB3	1.53	0.89
1:B:1104:ASP:HB3	1:B:1105:GLU:HA	1.53	0.89
1:B:1102:ASP:HB3	1:B:1104:ASP:N	1.90	0.83
1:A:753:GLU:HG2	1:A:1002:PHE:CE1	2.13	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	431/432 (100%)	417 (97%)	10 (2%)	4 (1%)	17	7
1	B	425/432 (98%)	404 (95%)	17 (4%)	4 (1%)	17	7
1	C	429/432 (99%)	413 (96%)	13 (3%)	3 (1%)	22	12
1	D	434/432 (100%)	422 (97%)	11 (2%)	1 (0%)	47	38
All	All	1719/1728 (100%)	1656 (96%)	51 (3%)	12 (1%)	22	12

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	960	ILE
1	B	960	ILE
1	C	785	VAL
1	C	960	ILE
1	D	960	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/372 (100%)	356 (95%)	17 (5%)	27	17
1	B	368/372 (99%)	345 (94%)	23 (6%)	18	8
1	C	371/372 (100%)	359 (97%)	12 (3%)	39	30
1	D	376/372 (101%)	361 (96%)	15 (4%)	31	22
All	All	1488/1488 (100%)	1421 (96%)	67 (4%)	27	18

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	997	ARG
1	B	1085	ILE
1	D	1017	ASN
1	B	1016	ASP
1	B	1030	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	741	GLN
1	C	1008	GLN
1	D	829	ASN
1	B	1064	ASN
1	D	784	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	E	1	1,2	14,14,15	0.64	0	17,19,21	1.25	3 (17%)
2	NAG	E	2	2	14,14,15	0.74	0	17,19,21	1.31	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	E	3	2	11,11,12	0.82	0	15,15,17	1.48	3 (20%)
2	BMA	E	4	2	11,11,12	0.75	1 (9%)	15,15,17	1.94	4 (26%)
3	NAG	F	1	1,3	14,14,15	0.59	0	17,19,21	1.35	2 (11%)
3	NAG	F	2	3	14,14,15	0.43	0	17,19,21	1.72	6 (35%)
3	BMA	F	3	3	11,11,12	1.10	1 (9%)	15,15,17	1.25	2 (13%)
4	NAG	G	1	1,4	14,14,15	0.49	0	17,19,21	1.46	4 (23%)
4	NAG	G	2	4	14,14,15	0.58	0	17,19,21	1.28	1 (5%)
4	BMA	G	3	4	11,11,12	0.51	0	15,15,17	1.10	1 (6%)
4	BMA	G	4	4	11,11,12	0.49	0	15,15,17	1.33	1 (6%)
4	BMA	G	5	4	11,11,12	0.52	0	15,15,17	2.09	3 (20%)
4	NAG	H	1	1,4	14,14,15	0.52	0	17,19,21	1.21	3 (17%)
4	NAG	H	2	4	14,14,15	0.80	0	17,19,21	1.39	1 (5%)
4	BMA	H	3	4	11,11,12	0.92	1 (9%)	15,15,17	3.40	6 (40%)
4	BMA	H	4	4	11,11,12	0.67	0	15,15,17	2.69	6 (40%)
4	BMA	H	5	4	11,11,12	0.51	0	15,15,17	2.00	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	BMA	E	3	2	-	2/2/19/22	0/1/1/1
2	BMA	E	4	2	-	2/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	4/6/23/26	0/1/1/1
3	NAG	F	2	3	-	1/6/23/26	0/1/1/1
3	BMA	F	3	3	-	2/2/19/22	0/1/1/1
4	NAG	G	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	BMA	G	4	4	-	2/2/19/22	1/1/1/1
4	BMA	G	5	4	-	2/2/19/22	0/1/1/1
4	NAG	H	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1
4	BMA	H	3	4	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BMA	H	4	4	-	2/2/19/22	0/1/1/1
4	BMA	H	5	4	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	3	BMA	O5-C1	3.17	1.48	1.43
4	H	3	BMA	O5-C1	2.32	1.47	1.43
2	E	4	BMA	O4-C4	2.00	1.47	1.43

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	3	BMA	O5-C1-C2	6.69	121.10	110.77
4	H	3	BMA	O5-C5-C6	6.26	117.03	107.20
4	G	5	BMA	C1-O5-C5	5.80	120.05	112.19
4	H	5	BMA	C1-O5-C5	-5.74	104.42	112.19
4	H	4	BMA	C1-O5-C5	5.69	119.90	112.19

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	3	BMA	C4-C5-C6-O6
2	E	3	BMA	O5-C5-C6-O6
4	G	5	BMA	O5-C5-C6-O6
4	H	3	BMA	O5-C5-C6-O6
4	G	4	BMA	O5-C5-C6-O6


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	G	4	BMA	C1-C2-C3-C4-C5-O5


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	5	BMA	2	0
4	H	1	NAG	1	0
2	E	4	BMA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
7	GOL	B	1206	-	5,5,5	0.33	0	5,5,5	0.34	0
6	MES	C	1207	-	12,12,12	2.39	1 (8%)	14,16,16	6.09	5 (35%)
5	NAG	C	1201	1	14,14,15	0.54	0	17,19,21	1.08	0
5	NAG	A	1201	1	14,14,15	0.52	0	17,19,21	1.26	2 (11%)
7	GOL	C	1208	-	5,5,5	0.20	0	5,5,5	0.44	0
6	MES	B	1205	-	12,12,12	2.09	1 (8%)	14,16,16	1.71	1 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MES	D	1208	-	12,12,12	2.14	1 (8%)	14,16,16	7.11	5 (35%)
7	GOL	D	1210	-	5,5,5	0.09	0	5,5,5	0.56	0
7	GOL	A	1207	-	5,5,5	0.21	0	5,5,5	0.62	0
5	NAG	D	1201	1	14,14,15	0.48	0	17,19,21	0.96	1 (5%)
7	GOL	D	1209	-	5,5,5	0.38	0	5,5,5	0.50	0
5	NAG	B	1201	1	14,14,15	0.49	0	17,19,21	1.10	0
6	MES	A	1206	-	12,12,12	2.26	1 (8%)	14,16,16	7.00	9 (64%)
6	MES	D	1207	-	12,12,12	2.40	1 (8%)	14,16,16	6.49	8 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	B	1206	-	-	1/4/4/4	-
6	MES	C	1207	-	-	2/6/14/14	0/1/1/1
5	NAG	C	1201	1	-	2/6/23/26	0/1/1/1
5	NAG	A	1201	1	-	2/6/23/26	0/1/1/1
7	GOL	C	1208	-	-	0/4/4/4	-
6	MES	B	1205	-	-	4/6/14/14	0/1/1/1
6	MES	D	1208	-	-	4/6/14/14	0/1/1/1
7	GOL	D	1210	-	-	2/4/4/4	-
7	GOL	A	1207	-	-	0/4/4/4	-
5	NAG	D	1201	1	-	2/6/23/26	0/1/1/1
7	GOL	D	1209	-	-	0/4/4/4	-
5	NAG	B	1201	1	-	4/6/23/26	0/1/1/1
6	MES	A	1206	-	-	3/6/14/14	0/1/1/1
6	MES	D	1207	-	-	5/6/14/14	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	1207	MES	C8-S	-8.11	1.66	1.77
6	D	1207	MES	C8-S	-8.00	1.66	1.77
6	A	1206	MES	C8-S	-7.49	1.66	1.77
6	D	1208	MES	C8-S	-7.08	1.67	1.77
6	B	1205	MES	C8-S	-6.84	1.67	1.77

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1206	MES	O3S-S-C8	-13.41	84.08	105.77
6	D	1208	MES	O3S-S-C8	-13.28	84.28	105.77
6	C	1207	MES	O2S-S-C8	13.00	122.57	106.92
6	D	1208	MES	O2S-S-C8	12.75	122.27	106.92
6	D	1207	MES	O3S-S-O1S	-11.99	81.97	111.27

There are no chirality outliers.

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	1207	MES	C7-C8-S-O2S
6	B	1205	MES	C7-C8-S-O2S
6	B	1205	MES	C7-C8-S-O3S
7	D	1210	GOL	O1-C1-C2-C3
6	A	1206	MES	N4-C7-C8-S

There are no ring outliers.

3 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	1208	MES	3	0
7	D	1210	GOL	10	0
6	A	1206	MES	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	431/432 (99%)	0.33	24 (5%) 24 27	22, 37, 68, 100	0
1	B	427/432 (98%)	1.57	113 (26%) 0 0	35, 70, 169, 214	0
1	C	430/432 (99%)	0.67	62 (14%) 2 2	28, 52, 110, 139	0
1	D	431/432 (99%)	0.45	34 (7%) 12 14	21, 39, 71, 128	0
All	All	1719/1728 (99%)	0.75	233 (13%) 3 3	21, 49, 120, 214	0

The worst 5 of 233 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	706	THR	13.7
1	B	709	VAL	13.1
1	B	1023	VAL	12.2
1	B	1067	GLY	10.6
1	B	737	VAL	10.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	F	3	11/12	0.59	0.14	88,93,96,99	0
4	BMA	G	4	11/12	0.63	0.29	105,110,113,115	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	BMA	G	5	11/12	0.64	0.44	108,113,115,116	0
4	BMA	H	4	11/12	0.66	0.21	64,73,76,76	0
2	BMA	E	3	11/12	0.79	0.20	101,104,105,106	0
4	BMA	H	5	11/12	0.79	0.35	94,103,107,109	0
3	NAG	F	1	14/15	0.80	0.13	79,83,91,92	0
4	BMA	G	3	11/12	0.81	0.27	98,102,107,109	0
4	NAG	G	2	14/15	0.81	0.14	69,78,84,92	0
4	BMA	H	3	11/12	0.81	0.17	51,64,73,84	0
2	BMA	E	4	11/12	0.83	0.18	102,103,104,104	0
4	NAG	G	1	14/15	0.83	0.17	75,78,90,93	0
3	NAG	F	2	14/15	0.87	0.12	75,81,83,88	0
2	NAG	E	2	14/15	0.92	0.11	34,38,46,51	0
4	NAG	H	2	14/15	0.94	0.10	35,36,44,44	0
4	NAG	H	1	14/15	0.94	0.10	37,38,48,53	0
2	NAG	E	1	14/15	0.95	0.10	37,39,52,57	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NAG	B	1201	14/15	0.64	0.33	129,141,158,160	0
5	NAG	C	1201	14/15	0.71	0.37	103,110,114,115	0
7	GOL	D	1210	6/6	0.79	0.33	33,39,47,47	0
5	NAG	D	1201	14/15	0.81	0.28	79,89,94,95	0
5	NAG	A	1201	14/15	0.83	0.31	82,96,99,100	0
7	GOL	B	1206	6/6	0.90	0.13	69,70,71,71	0
6	MES	A	1206	12/12	0.91	0.18	52,57,60,60	12
6	MES	D	1208	12/12	0.93	0.19	55,63,70,71	12
6	MES	C	1207	12/12	0.94	0.15	33,40,49,49	12
7	GOL	D	1209	6/6	0.94	0.09	35,39,40,43	0
7	GOL	A	1207	6/6	0.94	0.08	35,36,39,41	0
7	GOL	C	1208	6/6	0.95	0.08	47,54,57,62	0
6	MES	D	1207	12/12	0.95	0.12	69,76,81,81	0
6	MES	B	1205	12/12	0.96	0.17	59,67,74,75	0

6.5 Other polymers

There are no such residues in this entry.