



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 04:56 am BST

PDB ID : 4HVM
Title : Crystal structure of tallysomycin biosynthesis protein TlmII
Authors : Chang, C.; Bigelow, L.; Bearden, J.; Babnigg, G.; Bingman, C.A.; Yennamalli, R.; Lohman, J.; Ma, M.; Shen, B.; Phillips Jr., G.N.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG); Enzyme Discovery for Natural Product Biosynthesis (NatPro)
Deposited on : 2012-11-06
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	493	 % 73% 7% 20%
1	B	493	 % 75% 8% 16%
1	C	493	 % 70% 11% 19%
1	D	493	 % 78% 7% 15%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	D	501	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12790 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TlmII.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	393	Total 3098	C 1954	N 575	O 561	S 1	Se 7	0	0	0
1	B	414	Total 3212	C 2017	N 594	O 592	S 2	Se 7	0	0	0
1	C	401	Total 3137	C 1980	N 580	O 568	S 2	Se 7	0	0	0
1	D	419	Total 3258	C 2046	N 603	O 600	S 2	Se 7	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP A4KUC0
A	-1	ASN	-	EXPRESSION TAG	UNP A4KUC0
A	0	ALA	-	EXPRESSION TAG	UNP A4KUC0
B	-2	SER	-	EXPRESSION TAG	UNP A4KUC0
B	-1	ASN	-	EXPRESSION TAG	UNP A4KUC0
B	0	ALA	-	EXPRESSION TAG	UNP A4KUC0
C	-2	SER	-	EXPRESSION TAG	UNP A4KUC0
C	-1	ASN	-	EXPRESSION TAG	UNP A4KUC0
C	0	ALA	-	EXPRESSION TAG	UNP A4KUC0
D	-2	SER	-	EXPRESSION TAG	UNP A4KUC0
D	-1	ASN	-	EXPRESSION TAG	UNP A4KUC0
D	0	ALA	-	EXPRESSION TAG	UNP A4KUC0


- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	16	Total O 16 16	0	0
3	B	16	Total O 16 16	0	0
3	C	18	Total O 18 18	0	0
3	D	15	Total O 15 15	0	0



- Molecule 1: TlmII

Chain D: 78% 7% 15%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	123.80Å 128.01Å 129.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.71 – 2.70 45.49 – 2.70	Depositor EDS
% Data completeness (in resolution range)	77.6 (44.71-2.70) 90.8 (45.49-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.21 (at 2.69Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.182 , 0.232 0.195 , 0.246	Depositor DCC
R_{free} test set	2634 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	28.7	Xtriage
Anisotropy	0.198	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.020 for -h,l,k 0.017 for -l,-k,-h 0.019 for k,h,-l 0.004 for k,l,h 0.004 for l,h,k	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	12790	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/3161	0.39	0/4297
1	B	0.20	0/3277	0.39	0/4459
1	C	0.20	0/3202	0.40	0/4357
1	D	0.21	0/3326	0.40	0/4528
All	All	0.20	0/12966	0.40	0/17641

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3098	0	3033	18	0
1	B	3212	0	3134	23	0
1	C	3137	0	3069	26	0
1	D	3258	0	3170	18	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
2	D	5	0	0	2	0
3	A	16	0	0	0	0
3	B	16	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	18	0	0	0	0
3	D	15	0	0	0	0
All	All	12790	0	12406	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:415:ASP:HB2	1:D:430:ASP:HB2	1.79	0.63
1:B:216:ARG:NH2	1:B:389:ALA:O	2.31	0.63
1:C:113:PRO:HA	1:C:118:LEU:HD21	1.81	0.63
1:B:247:ALA:HA	1:B:272:LEU:HD23	1.81	0.62
1:A:113:PRO:HA	1:A:118:LEU:HD21	1.82	0.61
1:C:73:LEU:HD21	1:C:140:MSE:HE1	1.83	0.61
1:D:274:LEU:HB3	1:D:356:VAL:HB	1.83	0.60
1:B:227:LEU:HD22	1:B:451:MSE:HE3	1.82	0.60
1:B:111:THR:OG1	1:B:120:ARG:NH2	2.35	0.60
1:C:227:LEU:HD22	1:C:451:MSE:HE3	1.84	0.59
1:A:415:ASP:HB3	1:A:430:ASP:HB2	1.85	0.59
1:B:168:ARG:HB3	1:B:289:VAL:HG22	1.86	0.58
1:A:73:LEU:HD21	1:A:140:MSE:HE1	1.87	0.57
1:B:414:THR:HG22	1:B:431:GLN:HG2	1.87	0.57
1:C:65:VAL:HG21	1:C:75:MSE:HE3	1.86	0.56
1:C:168:ARG:HB3	1:C:289:VAL:HG22	1.88	0.55
1:A:111:THR:OG1	1:A:120:ARG:NH2	2.43	0.52
1:D:397:ARG:NH2	2:D:501:SO4:O1	2.42	0.51
1:D:73:LEU:HD21	1:D:140:MSE:HE1	1.92	0.51
1:B:65:VAL:HG11	1:B:75:MSE:HE3	1.93	0.50
1:D:281:ARG:NH2	1:D:291:GLY:O	2.44	0.50
1:D:242:MSE:HE2	1:D:416:LEU:HD21	1.93	0.50
1:C:163:ARG:O	1:C:166:GLN:HG2	2.11	0.50
1:A:96:GLN:HG2	1:A:126:LEU:HD13	1.93	0.50
1:B:217:LEU:HD11	1:B:448:ASP:HB2	1.95	0.49
1:B:36:ARG:NH1	3:B:612:HOH:O	2.46	0.49
1:A:11:ARG:NH1	1:A:171:ALA:O	2.45	0.49
1:D:17:GLN:HB3	1:D:140:MSE:HE3	1.95	0.48
1:A:144:ALA:HA	1:A:147:ARG:NH1	2.28	0.48
1:B:113:PRO:HA	1:B:118:LEU:HD21	1.96	0.48
1:C:240:LEU:HB3	1:C:315:MSE:HE1	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:17:GLN:HB3	1:C:140:MSE:HE3	1.95	0.47
1:A:65:VAL:HG21	1:A:75:MSE:HE3	1.95	0.47
1:B:274:LEU:HB3	1:B:356:VAL:HB	1.95	0.47
1:D:217:LEU:HB2	1:D:444:LEU:HD13	1.96	0.47
1:B:67:PHE:HB3	1:B:73:LEU:HD11	1.96	0.47
1:C:415:ASP:HB3	1:C:430:ASP:HB2	1.96	0.47
1:A:190:ARG:HH21	1:A:194:ARG:HH12	1.64	0.46
1:A:306:LEU:HD22	1:A:309:PRO:HB3	1.97	0.46
1:C:144:ALA:HA	1:C:147:ARG:NH1	2.30	0.46
1:D:156:GLN:O	1:D:161:ARG:N	2.49	0.46
1:D:252:THR:O	1:D:253:GLN:HB2	2.16	0.46
1:A:302:ASP:O	1:A:317:ARG:HD3	2.16	0.46
1:C:64:ARG:HD3	1:C:81:ALA:HB2	1.98	0.45
1:B:163:ARG:O	1:B:166:GLN:HG2	2.16	0.45
1:A:215:SER:HB3	1:D:402:THR:HG22	1.99	0.45
1:A:89:LEU:HD13	1:A:98:ARG:HB2	1.98	0.45
1:A:24:LEU:HD22	1:A:32:ALA:HB2	1.98	0.44
1:C:177:HIS:HA	1:C:178:PRO:HD3	1.90	0.44
1:C:24:LEU:HD22	1:C:32:ALA:HB2	2.00	0.44
1:C:112:ASP:OD1	1:C:114:GLU:HG2	2.17	0.44
1:B:64:ARG:HD3	1:B:81:ALA:HB2	2.00	0.44
1:D:106:ASP:OD2	1:D:122:HIS:NE2	2.32	0.44
1:C:281:ARG:HA	1:C:286:LEU:HB2	2.00	0.44
1:D:418:VAL:HG12	1:D:427:ILE:HD13	1.99	0.43
1:C:28:PRO:HA	1:C:29:PRO:HD3	1.87	0.43
1:A:403:TRP:CZ3	1:B:401:ASP:HB2	2.54	0.43
1:B:229:GLU:HA	1:B:230:PRO:HD3	1.90	0.43
1:C:96:GLN:HG2	1:C:126:LEU:HD13	2.00	0.43
1:D:36:ARG:NH2	2:D:501:SO4:O4	2.49	0.43
1:A:313:ASP:O	1:A:317:ARG:HG3	2.18	0.43
1:B:223:ARG:HG2	1:B:455:VAL:HG21	2.01	0.43
1:C:94:ASP:O	1:C:98:ARG:HG2	2.19	0.42
1:D:182:GLU:O	1:D:186:VAL:HG13	2.19	0.42
1:B:105:ALA:HB1	1:C:452:ALA:O	2.19	0.42
1:B:210:GLY:O	1:B:432:ARG:NH1	2.45	0.42
1:D:163:ARG:O	1:D:166:GLN:HG2	2.19	0.42
1:C:182:GLU:HA	1:C:185:ARG:HG2	1.99	0.42
1:B:234:LEU:HB2	1:B:311:PHE:CE2	2.54	0.42
1:B:106:ASP:O	1:B:120:ARG:NH1	2.52	0.42
1:C:14:SER:O	1:C:18:ARG:HG3	2.19	0.42
1:C:45:ASP:HB3	1:C:48:ARG:HB3	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ASP:OD1	1:A:114:GLU:HG2	2.20	0.42
1:B:22:ARG:NH1	1:B:337:ASP:OD2	2.53	0.41
1:D:21:TRP:HE3	1:D:140:MSE:SE	2.54	0.41
1:C:92:HIS:HA	1:C:93:PRO:HD3	1.90	0.41
1:C:330:LEU:HD12	1:C:331:PRO:HD2	2.03	0.41
1:C:143:ASP:O	1:C:147:ARG:HG3	2.21	0.41
1:A:145:ARG:O	1:A:149:MSE:HG3	2.21	0.41
1:D:144:ALA:HA	1:D:147:ARG:NH1	2.36	0.40
1:B:415:ASP:HB2	1:B:430:ASP:OD1	2.21	0.40
1:C:22:ARG:NH1	1:C:333:HIS:HB2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/493 (77%)	367 (97%)	12 (3%)	0	100	100
1	B	404/493 (82%)	390 (96%)	14 (4%)	0	100	100
1	C	389/493 (79%)	374 (96%)	13 (3%)	2 (0%)	29	54
1	D	409/493 (83%)	394 (96%)	15 (4%)	0	100	100
All	All	1581/1972 (80%)	1525 (96%)	54 (3%)	2 (0%)	51	78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	304	THR
1	C	463	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/386 (83%)	319 (100%)	0	100	100
1	B	330/386 (86%)	323 (98%)	7 (2%)	53	80
1	C	321/386 (83%)	318 (99%)	3 (1%)	78	92
1	D	334/386 (86%)	332 (99%)	2 (1%)	86	95
All	All	1304/1544 (84%)	1292 (99%)	12 (1%)	78	92

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	101	GLU
1	B	229	GLU
1	B	234	LEU
1	B	274	LEU
1	B	282	ASP
1	B	401	ASP
1	B	430	ASP
1	C	82	ASP
1	C	180	ASP
1	C	285	GLN
1	D	10	LEU
1	D	282	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	502	-	4,4,4	0.14	0	6,6,6	0.11	0
2	SO4	A	501	-	4,4,4	0.14	0	6,6,6	0.19	0
2	SO4	D	501	-	4,4,4	0.15	0	6,6,6	0.16	0
2	SO4	B	501	-	4,4,4	0.14	0	6,6,6	0.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	501	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/493 (78%)	-0.29	6 (1%) 72 74	12, 30, 64, 92	0
1	B	407/493 (82%)	-0.07	6 (1%) 73 76	19, 47, 83, 115	0
1	C	394/493 (79%)	-0.21	7 (1%) 68 70	11, 35, 75, 104	0
1	D	412/493 (83%)	-0.24	1 (0%) 95 96	11, 38, 80, 110	0
All	All	1599/1972 (81%)	-0.20	20 (1%) 77 78	11, 37, 77, 115	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	201	VAL	4.0
1	A	71	ASP	3.5
1	C	393	GLY	3.3
1	B	403	TRP	3.2
1	C	71	ASP	3.0
1	A	360	LEU	2.9
1	C	9	GLY	2.9
1	B	201	VAL	2.7
1	B	402	THR	2.7
1	A	70	ASP	2.6
1	A	43	ARG	2.5
1	B	463	PRO	2.4
1	C	79	GLU	2.4
1	B	163	ARG	2.4
1	B	406	THR	2.3
1	C	200	PRO	2.2
1	A	393	GLY	2.2
1	A	161	ARG	2.2
1	D	11	ARG	2.1
1	C	10	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	501	5/5	0.91	0.20	67,74,80,82	0
2	SO4	A	502	5/5	0.97	0.15	75,76,78,82	0
2	SO4	D	501	5/5	0.97	0.15	39,42,45,52	0
2	SO4	B	501	5/5	0.98	0.13	49,51,54,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.