



Full wwPDB X-ray Structure Validation Report

Sep 10, 2023 – 11:48 PM EDT

PDB ID : 4KGL
Title : Crystal structure of human alpha-L-iduronidase complex with [2R,3R,4R,5S]-2-carboxy-3,4,5-trihydroxy-piperidine
Authors : Bie, H.; Yin, J.; He, X.; Kermode, A.R.; Goddard-Borger, E.D.; Withers, S.G.; James, M.N.G.
Deposited on : 2013-04-29
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	627	
1	B	627	
2	C	2	
2	E	2	
3	D	5	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	9	 11% 67% 22%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 9990 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-L-iduronidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	595	4734	3030	852	840	12	0	0	0
1	B	612	4888	3121	886	867	14	0	4	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	GLN	HIS	SEE REMARK 999	UNP P35475
A	63	PRO	GLN	SEE REMARK 999	UNP P35475
A	105	GLN	ARG	SEE REMARK 999	UNP P35475
B	33	GLN	HIS	SEE REMARK 999	UNP P35475
B	63	PRO	GLN	SEE REMARK 999	UNP P35475
B	105	GLN	ARG	SEE REMARK 999	UNP P35475

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	2	28	16	2	10	0	0	0
2	E	2	28	16	2	10	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	5	61	34	2	25	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	F	9	105	58	2	45	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	B	1	14	8	1	5	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	6	3	3	0	0
6	B	1	6	3	3	0	0
6	B	1	6	3	3	0	0
6	B	1	6	3	3	0	0
6	B	1	6	3	3	0	0

- Molecule 7 is (2R,3R,4R,5S)-3,4,5-trihydroxypiperidine-2-carboxylic acid (three-letter code: IDJ) (formula: C₆H₁₁NO₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
7	A	1	12	6	1	5	0	0
7	B	1	12	6	1	5	0	0

- Molecule 8 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: $C_4H_6O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	B	1	10	4	6	0	0

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	20	Total 20	O 20	0	0
9	B	34	Total 34	O 34	0	0

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50% 50%


MAG1
MAG2

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  20% 40% 40%

MAG1
MAG2
BMA3
MAN4
MAN5

- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  11% 67% 22%

MAG1
MAG2
BMA3
MAN4
MAN5
MAN6
MAN7
MAN8
MAN9

4 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	259.36Å 259.36Å 71.57Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.06 – 2.70 49.01 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.06-2.70) 99.4 (49.01-2.70)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.204 , 0.254 0.202 , 0.250	Depositor DCC
R_{free} test set	2479 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	48.2	Xtrriage
Anisotropy	0.797	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.007 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9990	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, TLA, BMA, GOL, NAG, IDJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/4879	0.55	0/6666
1	B	0.36	0/5039	0.57	0/6886
All	All	0.35	0/9918	0.56	0/13552

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4734	0	4632	34	0
1	B	4888	0	4786	37	0
2	C	28	0	25	0	0
2	E	28	0	25	1	0
3	D	61	0	52	2	0
4	F	105	0	88	2	0
5	A	14	0	13	0	0
5	B	14	0	13	0	0
6	A	6	0	8	3	0
6	B	24	0	32	2	0
7	A	12	0	10	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	12	0	10	0	0
8	B	10	0	4	0	0
9	A	20	0	0	0	0
9	B	34	0	0	1	0
All	All	9990	0	9698	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:577[A]:CYS:SG	1:B:577[A]:CYS:O	2.33	0.86
1:A:315:ASP:HB2	1:A:478:ASN:OD1	1.77	0.83
3:D:3:BMA:H61	3:D:5:MAN:H5	1.74	0.69
1:B:270:ILE:HD11	2:E:1:NAG:H82	1.80	0.63
1:B:334:LEU:HD21	1:B:343:TYR:HB3	1.80	0.63
1:B:320:ALA:HB1	1:B:443:SER:HB2	1.81	0.61
1:A:152:TRP:O	1:A:156:VAL:HG23	2.01	0.61
1:A:131:GLU:OE2	1:A:181:ASN:HB2	2.01	0.60
1:B:356:HIS:HB2	6:B:914:GOL:H31	1.82	0.60
1:A:273:LEU:HD11	1:A:333:LEU:HD13	1.84	0.59
1:A:298:ASP:HA	1:A:347:SER:HB3	1.84	0.59
1:B:359:PRO:HG3	6:B:914:GOL:H12	1.85	0.58
1:A:324:LYS:NZ	6:A:909:GOL:H11	2.18	0.58
1:A:378:HIS:CD2	1:A:560:THR:HB	2.38	0.58
1:A:36:ALA:HA	1:A:401:LEU:HD13	1.86	0.57
3:D:3:BMA:H61	3:D:5:MAN:C5	2.35	0.57
1:B:414:SER:C	1:B:416:HIS:H	2.11	0.54
1:A:166:ARG:HG3	1:A:166:ARG:HH11	1.75	0.52
1:A:476:LEU:HG	1:A:510:PRO:HB3	1.91	0.52
1:B:242:HIS:CD2	1:B:289:LYS:HB2	2.45	0.51
1:B:56:LEU:HD13	4:F:5:MAN:H3	1.91	0.51
1:A:242:HIS:CD2	1:A:289:LYS:HB2	2.46	0.51
1:A:261:LEU:HD13	1:A:263:ARG:HG2	1.92	0.50
1:B:315:ASP:HA	1:B:385:PRO:HG2	1.93	0.50
1:A:492:ARG:NH2	4:F:3:BMA:O2	2.46	0.49
1:A:368:ARG:HA	1:A:381:LEU:HD23	1.95	0.48
1:A:434:TRP:CH2	1:A:436:ALA:HB2	2.48	0.48
1:A:383:ARG:HG2	1:A:387:LEU:HD23	1.95	0.48
1:B:584:GLN:HB3	1:B:592:TYR:HB3	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:267:ARG:HG2	1:A:446:THR:HG21	1.95	0.48
1:B:98:THR:OG1	1:B:110:ASN:HB3	2.13	0.48
1:A:276:GLU:OE1	1:A:329:HIS:NE2	2.44	0.47
1:A:32:VAL:HG22	1:A:405:VAL:HG22	1.96	0.46
1:A:315:ASP:HA	1:A:385:PRO:HG2	1.97	0.46
1:B:242:HIS:NE2	1:B:289:LYS:HB2	2.31	0.46
1:B:401:LEU:HD21	1:B:424:ALA:HB2	1.97	0.46
1:A:365:LEU:O	1:A:384:LYS:HB2	2.16	0.45
1:B:386:VAL:HG13	1:B:387:LEU:N	2.31	0.45
1:B:73:ASN:HB2	1:B:353:LEU:HD21	1.97	0.45
1:B:314:ALA:HB1	1:B:360:PHE:HB3	1.98	0.45
1:A:419:GLY:HA3	1:A:441:TYR:CE1	2.52	0.45
1:A:171:HIS:CD2	1:A:175:TRP:HE1	2.35	0.45
1:A:509:ASP:HB2	1:A:598:LYS:HE3	2.00	0.44
1:A:86:LYS:HE2	1:A:127:LEU:HD11	2.00	0.44
1:A:154:ASP:O	1:A:158:SER:HB2	2.18	0.44
1:A:242:HIS:NE2	1:A:289:LYS:HB2	2.33	0.44
1:B:298:ASP:HA	1:B:347:SER:HB3	1.99	0.43
1:A:166:ARG:HD3	1:A:167:TYR:CZ	2.54	0.43
1:B:99:THR:HG21	1:B:139:HIS:HB2	2.01	0.43
1:B:475:TYR:HE1	1:B:530:LEU:HD22	1.84	0.43
1:A:324:LYS:HZ3	6:A:909:GOL:H11	1.81	0.43
1:B:91:HIS:ND1	1:B:131:GLU:OE1	2.52	0.43
1:B:378:HIS:CD2	1:B:560:THR:HB	2.54	0.43
1:B:456:VAL:O	1:B:528:PRO:HD2	2.18	0.43
1:B:182:GLU:HB3	1:B:185:HIS:HB2	2.01	0.42
1:B:48:ARG:HD2	1:B:83:ARG:O	2.18	0.42
1:B:395:LEU:HD22	1:B:576:LYS:HB3	2.01	0.42
1:A:320:ALA:HB1	1:A:443:SER:HB2	2.01	0.42
1:A:324:LYS:HZ1	6:A:909:GOL:H11	1.83	0.42
1:B:317:THR:HA	1:B:533:PRO:HG3	2.02	0.42
1:B:624:ASP:OD2	1:B:628:ARG:HB2	2.20	0.42
1:A:263:ARG:HD3	1:A:263:ARG:HA	1.87	0.42
1:B:372:ASN:HA	1:B:377:PRO:HB3	2.02	0.41
1:B:337:THR:HG21	9:B:1022:HOH:O	2.19	0.41
1:B:564:LEU:HD12	1:B:564:LEU:C	2.39	0.41
1:B:329:HIS:O	1:B:333:LEU:HB3	2.21	0.41
1:B:298:ASP:O	1:B:299:GLU:HG3	2.19	0.41
1:B:312:TRP:CD2	1:B:446:THR:HG22	2.56	0.41
1:A:423:SER:OG	1:A:437:ALA:HB3	2.21	0.41
1:B:103:THR:HG21	1:B:147:GLN:HE21	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:LEU:HA	1:A:309:PRO:HD3	1.91	0.40
1:B:177:PHE:HB2	1:B:217:ARG:O	2.21	0.40
1:B:261:LEU:O	1:B:297:ASN:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	587/627 (94%)	543 (92%)	41 (7%)	3 (0%)	29 54
1	B	612/627 (98%)	569 (93%)	36 (6%)	7 (1%)	14 34
All	All	1199/1254 (96%)	1112 (93%)	77 (6%)	10 (1%)	19 43

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	338	THR
1	A	339	SER
1	A	340	ALA
1	B	340	ALA
1	B	60	GLN
1	B	415	ASN
1	B	61	ALA
1	B	587	GLN
1	B	588	ASP
1	B	430	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	503/527 (95%)	494 (98%)	9 (2%)	59 83
1	B	520/527 (99%)	504 (97%)	16 (3%)	40 69
All	All	1023/1054 (97%)	998 (98%)	25 (2%)	47 77

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	TRP
1	A	98	THR
1	A	107	LEU
1	A	127	LEU
1	A	239	ARG
1	A	262	HIS
1	A	349	ASP
1	A	598	LYS
1	A	639	LEU
1	B	38	ARG
1	B	41	TRP
1	B	59	SER
1	B	69	ASP
1	B	91	HIS
1	B	130	PHE
1	B	147	GLN
1	B	245	THR
1	B	262	HIS
1	B	337	THR
1	B	349	ASP
1	B	417	THR
1	B	426	ARG
1	B	555	ARG
1	B	561	GLN
1	B	567	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	GLN
1	A	125	GLN
1	A	281	GLN
1	A	587	GLN
1	B	147	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.52	0	17,19,21	1.38	2 (11%)
2	NAG	C	2	2	14,14,15	0.58	0	17,19,21	0.94	1 (5%)
3	NAG	D	1	3,1	14,14,15	0.58	0	17,19,21	1.06	1 (5%)
3	NAG	D	2	3	14,14,15	0.54	0	17,19,21	1.45	1 (5%)
3	BMA	D	3	3	11,11,12	0.40	0	15,15,17	0.98	1 (6%)
3	MAN	D	4	3	11,11,12	0.58	0	15,15,17	0.68	0
3	MAN	D	5	3	11,11,12	0.57	0	15,15,17	1.98	3 (20%)
2	NAG	E	1	2,1	14,14,15	0.55	0	17,19,21	0.99	1 (5%)
2	NAG	E	2	2	14,14,15	0.51	0	17,19,21	1.34	2 (11%)
4	NAG	F	1	4,1	14,14,15	0.50	0	17,19,21	1.32	2 (11%)
4	NAG	F	2	4	14,14,15	0.57	0	17,19,21	1.10	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BMA	F	3	4	11,11,12	0.63	0	15,15,17	1.47	1 (6%)
4	MAN	F	4	4	11,11,12	0.66	0	15,15,17	0.93	1 (6%)
4	MAN	F	5	4	11,11,12	0.51	0	15,15,17	1.34	1 (6%)
4	MAN	F	6	4	11,11,12	0.57	0	15,15,17	0.78	1 (6%)
4	MAN	F	7	4	11,11,12	0.57	0	15,15,17	1.10	1 (6%)
4	MAN	F	8	4	11,11,12	0.56	0	15,15,17	0.80	0
4	MAN	F	9	4	11,11,12	0.74	0	15,15,17	1.49	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
3	NAG	D	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	BMA	D	3	3	-	2/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	MAN	D	5	3	-	2/2/19/22	0/1/1/1
2	NAG	E	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
4	NAG	F	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	F	2	4	-	0/6/23/26	0/1/1/1
4	BMA	F	3	4	-	0/2/19/22	0/1/1/1
4	MAN	F	4	4	-	0/2/19/22	0/1/1/1
4	MAN	F	5	4	-	2/2/19/22	0/1/1/1
4	MAN	F	6	4	-	0/2/19/22	0/1/1/1
4	MAN	F	7	4	-	2/2/19/22	0/1/1/1
4	MAN	F	8	4	-	2/2/19/22	0/1/1/1
4	MAN	F	9	4	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	5	MAN	C1-O5-C5	5.57	119.74	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	5	MAN	C1-O5-C5	4.43	118.19	112.19
2	C	1	NAG	C1-O5-C5	4.41	118.17	112.19
4	F	3	BMA	C1-C2-C3	4.19	114.82	109.67
3	D	2	NAG	C1-O5-C5	4.16	117.83	112.19
3	D	5	MAN	C1-C2-C3	3.92	114.49	109.67
4	F	9	MAN	C2-C3-C4	3.41	116.80	110.89
4	F	1	NAG	O5-C5-C6	3.22	112.25	107.20
4	F	9	MAN	C1-C2-C3	3.16	113.55	109.67
2	E	2	NAG	C1-O5-C5	3.14	116.45	112.19
4	F	1	NAG	O5-C1-C2	-2.96	106.62	111.29
4	F	2	NAG	C1-O5-C5	2.94	116.18	112.19
4	F	7	MAN	C1-O5-C5	2.84	116.04	112.19
2	E	2	NAG	C2-N2-C7	2.80	126.89	122.90
2	E	1	NAG	O5-C5-C6	2.77	111.55	107.20
4	F	9	MAN	C3-C4-C5	2.56	114.81	110.24
3	D	1	NAG	O5-C1-C2	-2.54	107.27	111.29
2	C	2	NAG	C1-O5-C5	2.50	115.58	112.19
3	D	5	MAN	O5-C1-C2	2.50	114.62	110.77
3	D	3	BMA	O5-C5-C6	2.49	111.11	107.20
4	F	6	MAN	C1-O5-C5	2.05	114.97	112.19
2	C	1	NAG	O5-C1-C2	-2.03	108.08	111.29
4	F	4	MAN	C1-C2-C3	2.00	112.12	109.67

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1	NAG	O5-C5-C6-O6
3	D	3	BMA	O5-C5-C6-O6
3	D	5	MAN	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
3	D	3	BMA	C4-C5-C6-O6
4	F	8	MAN	C4-C5-C6-O6
4	F	8	MAN	O5-C5-C6-O6
2	E	1	NAG	C4-C5-C6-O6
3	D	5	MAN	C4-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6
4	F	7	MAN	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
4	F	7	MAN	O5-C5-C6-O6
4	F	9	MAN	O5-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	1	NAG	C4-C5-C6-O6
4	F	5	MAN	O5-C5-C6-O6
4	F	9	MAN	C4-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
3	D	1	NAG	O5-C5-C6-O6
4	F	5	MAN	C4-C5-C6-O6


There are no ring outliers.


5 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	3	BMA	1	0
4	F	5	MAN	1	0
3	D	5	MAN	2	0
2	E	1	NAG	1	0
3	D	3	BMA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	A	909	-	5,5,5	0.23	0	5,5,5	0.19	0
5	NAG	A	901	1	14,14,15	0.54	0	17,19,21	1.22	2 (11%)
6	GOL	B	916	-	5,5,5	0.36	0	5,5,5	0.65	0
7	IDJ	A	910	-	12,12,12	1.15	0	12,17,17	2.27	2 (16%)
8	TLA	B	917	-	9,9,9	1.16	0	12,12,12	1.23	1 (8%)
6	GOL	B	914	-	5,5,5	0.35	0	5,5,5	0.22	0
7	IDJ	B	918	-	12,12,12	1.07	0	12,17,17	1.96	2 (16%)
6	GOL	B	913	-	5,5,5	0.36	0	5,5,5	0.25	0
5	NAG	B	901	1	14,14,15	0.47	0	17,19,21	0.91	1 (5%)
6	GOL	B	915	-	5,5,5	0.33	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	909	-	-	2/4/4/4	-
5	NAG	A	901	1	-	1/6/23/26	0/1/1/1
6	GOL	B	916	-	-	0/4/4/4	-
7	IDJ	A	910	-	-	1/3/21/21	0/1/1/1
8	TLA	B	917	-	-	8/12/12/12	-
6	GOL	B	914	-	-	4/4/4/4	-
7	IDJ	B	918	-	-	0/3/21/21	0/1/1/1
6	GOL	B	913	-	-	0/4/4/4	-
5	NAG	B	901	1	-	2/6/23/26	0/1/1/1
6	GOL	B	915	-	-	4/4/4/4	-

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	910	IDJ	C1-C2-C3	5.44	116.72	110.33
7	B	918	IDJ	C1-C2-C3	5.07	116.28	110.33
7	A	910	IDJ	C2-C3-C4	4.80	119.19	110.89
7	B	918	IDJ	C2-C3-C4	3.59	117.11	110.89
5	A	901	NAG	C1-O5-C5	3.20	116.53	112.19
8	B	917	TLA	O41-C4-C3	2.69	120.54	113.27
5	B	901	NAG	C1-O5-C5	2.37	115.41	112.19
5	A	901	NAG	O5-C5-C6	2.32	110.85	107.20

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	909	GOL	O1-C1-C2-O2
6	A	909	GOL	O1-C1-C2-C3
8	B	917	TLA	C2-C3-C4-O4
8	B	917	TLA	C2-C3-C4-O41
8	B	917	TLA	O3-C3-C4-O4
8	B	917	TLA	O3-C3-C4-O41
5	B	901	NAG	O5-C5-C6-O6
5	B	901	NAG	C4-C5-C6-O6
8	B	917	TLA	C1-C2-C3-O3
8	B	917	TLA	O2-C2-C3-C4
6	B	914	GOL	O1-C1-C2-O2
8	B	917	TLA	O2-C2-C3-O3
6	B	914	GOL	O1-C1-C2-C3
6	B	914	GOL	C1-C2-C3-O3
6	B	915	GOL	C1-C2-C3-O3
6	B	914	GOL	O2-C2-C3-O3
8	B	917	TLA	C1-C2-C3-C4
6	B	915	GOL	O2-C2-C3-O3
6	B	915	GOL	O1-C1-C2-O2
5	A	901	NAG	C4-C5-C6-O6
6	B	915	GOL	O1-C1-C2-C3
7	A	910	IDJ	C4-C5-C6-O6A

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	909	GOL	3	0
6	B	914	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	595/627 (94%)	0.05	22 (3%) 41 41	34, 52, 81, 105	0
1	B	612/627 (97%)	-0.14	8 (1%) 77 78	33, 47, 76, 112	0
All	All	1207/1254 (96%)	-0.04	30 (2%) 57 59	33, 49, 79, 112	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	339	SER	5.2
1	B	338	THR	3.8
1	B	588	ASP	3.7
1	A	639	LEU	3.4
1	B	641	VAL	3.3
1	A	338	THR	3.2
1	A	614	VAL	3.1
1	B	635	PRO	3.1
1	B	642	PRO	2.9
1	A	618	TYR	2.9
1	B	590	LYS	2.8
1	A	611	THR	2.7
1	A	585	PHE	2.7
1	A	524	LEU	2.7
1	A	623	LEU	2.7
1	A	588	ASP	2.6
1	B	229	PRO	2.5
1	A	411	VAL	2.4
1	A	64	TYR	2.3
1	A	526	LEU	2.3
1	A	638	TYR	2.2
1	A	620	VAL	2.2
1	A	523	ARG	2.2
1	A	335	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	536	LEU	2.1
1	A	336	ASN	2.1
1	B	589	GLY	2.1
1	A	473	THR	2.1
1	A	458	LEU	2.0
1	A	610	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
4	MAN	F	9	11/12	0.75	0.21	78,89,91,92	0
3	MAN	D	5	11/12	0.79	0.21	75,82,85,91	0
3	MAN	D	4	11/12	0.80	0.30	98,102,105,107	0
4	MAN	F	7	11/12	0.81	0.19	80,89,92,95	0
2	NAG	C	2	14/15	0.83	0.34	94,102,109,115	0
2	NAG	E	2	14/15	0.84	0.34	75,77,80,81	0
4	MAN	F	5	11/12	0.84	0.17	92,99,103,105	0
4	MAN	F	6	11/12	0.86	0.17	85,93,100,101	0
3	BMA	D	3	11/12	0.93	0.13	74,78,83,93	0
2	NAG	E	1	14/15	0.94	0.20	53,59,64,71	0
4	MAN	F	4	11/12	0.94	0.13	69,76,84,90	0
4	MAN	F	8	11/12	0.94	0.12	70,73,76,83	0
2	NAG	C	1	14/15	0.94	0.20	56,64,70,83	0
4	BMA	F	3	11/12	0.95	0.12	50,53,61,61	0
4	NAG	F	2	14/15	0.96	0.16	42,43,49,49	0
3	NAG	D	2	14/15	0.96	0.13	53,61,68,74	0
3	NAG	D	1	14/15	0.96	0.15	56,58,61,63	0
4	NAG	F	1	14/15	0.96	0.16	40,41,44,45	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	901	14/15	0.78	0.25	87,95,99,99	0
7	IDJ	A	910	12/12	0.86	0.23	82,86,88,89	0
6	GOL	B	913	6/6	0.88	0.28	55,62,64,65	0
6	GOL	B	916	6/6	0.89	0.20	41,45,47,51	0
8	TLA	B	917	10/10	0.91	0.23	63,66,70,71	0
6	GOL	B	914	6/6	0.92	0.20	51,57,57,58	0
7	IDJ	B	918	12/12	0.92	0.16	69,73,75,77	0
6	GOL	A	909	6/6	0.92	0.19	50,52,54,56	0
5	NAG	B	901	14/15	0.93	0.19	55,58,61,62	0
6	GOL	B	915	6/6	0.94	0.19	51,52,53,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.