



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 09:39 pm BST

PDB ID : 4QAX
Title : Crystal structure of post-catalytic binary complex of Phosphoglycerate mutase from *Staphylococcus aureus*
Authors : Roychowdhury, A.; Kundu, A.; Bose, M.; Gujar, A.; Das, A.K.
Deposited on : 2014-05-06
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	513	 % 84% 11% ...

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	2PG	A	603	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4235 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2,3-bisphosphoglycerate-independent phosphoglycerate mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	503	3934	2478	665	780	11	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	HIS	-	EXPRESSION TAG	UNP Q2G029
A	-6	HIS	-	EXPRESSION TAG	UNP Q2G029
A	-5	HIS	-	EXPRESSION TAG	UNP Q2G029
A	-4	HIS	-	EXPRESSION TAG	UNP Q2G029
A	-3	HIS	-	EXPRESSION TAG	UNP Q2G029
A	-2	HIS	-	EXPRESSION TAG	UNP Q2G029
A	-1	GLY	-	EXPRESSION TAG	UNP Q2G029
A	0	SER	-	EXPRESSION TAG	UNP Q2G029

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		

- Molecule 3 is 2-PHOSPHOGLYCERIC ACID (three-letter code: 2PG) (formula: C₃H₇O₇P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
3	A	1	11	3	7	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	288	288	288	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2,3-bisphosphoglycerate-independent phosphoglycerate mutase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.80Å 80.69Å 89.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.53 – 2.00 19.53 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (19.53-2.00) 99.5 (19.53-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.166 , 0.214 0.178 , 0.222	Depositor DCC
R_{free} test set	1787 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	22.2	Xtrriage
Anisotropy	0.350	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4235	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MN, 2PG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.91	0/4009	1.19	32/5432 (0.6%)

There are no bond length outliers.

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	ARG	NE-CZ-NH2	-16.74	111.93	120.30
1	A	185	ARG	NE-CZ-NH2	-16.63	111.98	120.30
1	A	17	ARG	NE-CZ-NH1	15.82	128.21	120.30
1	A	185	ARG	NE-CZ-NH1	15.74	128.17	120.30
1	A	206	ARG	NE-CZ-NH2	-15.30	112.65	120.30
1	A	260	ARG	NE-CZ-NH1	15.15	127.88	120.30
1	A	206	ARG	NE-CZ-NH1	13.84	127.22	120.30
1	A	350	ARG	NE-CZ-NH2	-13.50	113.55	120.30
1	A	17	ARG	NE-CZ-NH2	-13.34	113.63	120.30
1	A	350	ARG	NE-CZ-NH1	12.23	126.42	120.30
1	A	35	ARG	NE-CZ-NH2	9.75	125.17	120.30
1	A	35	ARG	NE-CZ-NH1	-9.59	115.51	120.30
1	A	334	VAL	CG1-CB-CG2	7.55	122.97	110.90
1	A	481	LEU	CB-CG-CD1	7.25	123.32	111.00
1	A	153	ARG	NE-CZ-NH2	-7.20	116.70	120.30
1	A	391	LEU	CB-CG-CD2	7.06	123.00	111.00
1	A	259	ASP	CB-CG-OD1	6.48	124.13	118.30
1	A	260	ARG	CD-NE-CZ	6.47	132.66	123.60
1	A	17	ARG	CD-NE-CZ	6.29	132.40	123.60
1	A	243	ASP	CB-CG-OD2	6.24	123.92	118.30
1	A	42	THR	N-CA-CB	-6.23	98.47	110.30
1	A	260	ARG	CG-CD-NE	-5.99	99.22	111.80
1	A	206	ARG	CD-NE-CZ	5.81	131.74	123.60
1	A	486	LEU	CB-CG-CD1	5.49	120.33	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	194	ARG	NE-CZ-NH2	-5.48	117.56	120.30
1	A	453	MET	CA-CB-CG	5.45	122.56	113.30
1	A	185	ARG	CD-NE-CZ	5.44	131.22	123.60
1	A	488	LEU	CB-CG-CD1	5.43	120.24	111.00
1	A	496	ASP	CB-CG-OD1	5.33	123.10	118.30
1	A	355	ASP	CB-CG-OD1	5.32	123.09	118.30
1	A	352	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	A	153	ARG	NE-CZ-NH1	5.19	122.90	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3934	0	3785	43	0
2	A	2	0	0	0	0
3	A	11	0	4	12	0
4	A	288	0	0	11	1
All	All	4235	0	3789	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:ARG:HD2	3:A:603:2PG:O1	1.26	1.23
1:A:257:ARG:NH2	3:A:603:2PG:O2	1.91	1.04
1:A:260:ARG:CD	3:A:603:2PG:O1	2.10	0.98
1:A:206:ARG:HD2	4:A:754:HOH:O	1.64	0.95
1:A:185:ARG:HD3	4:A:763:HOH:O	1.66	0.93
1:A:80:THR:HG21	4:A:859:HOH:O	1.77	0.84
1:A:16:ASN:HD22	1:A:44:GLN:HE21	1.26	0.84
1:A:193:LYS:H	1:A:263:GLN:HE22	1.27	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:THR:HG23	4:A:829:HOH:O	1.81	0.79
1:A:316:GLN:HE21	1:A:347:LYS:H	1.32	0.78
1:A:260:ARG:NH1	3:A:603:2PG:O4P	2.20	0.73
1:A:125:HIS:HD2	1:A:127:LYS:H	1.37	0.71
1:A:206:ARG:HD3	1:A:245:VAL:HB	1.72	0.70
1:A:260:ARG:HD2	3:A:603:2PG:C1	2.20	0.68
3:A:603:2PG:C1	3:A:603:2PG:O4P	2.39	0.67
1:A:212:THR:CG2	4:A:822:HOH:O	2.45	0.65
1:A:42:THR:CG2	4:A:829:HOH:O	2.42	0.63
1:A:141:VAL:O	1:A:142:GLU:CB	2.49	0.59
1:A:94:ASN:HD22	1:A:97:LEU:H	1.50	0.59
1:A:193:LYS:H	1:A:263:GLN:NE2	2.03	0.54
1:A:260:ARG:NH1	3:A:603:2PG:O1	2.39	0.51
1:A:185:ARG:CD	4:A:763:HOH:O	2.42	0.51
3:A:603:2PG:O1	3:A:603:2PG:O4P	2.29	0.51
1:A:212:THR:HG23	4:A:822:HOH:O	2.10	0.51
1:A:103:HIS:CE1	1:A:283:ASP:O	2.65	0.49
1:A:103:HIS:HE1	1:A:283:ASP:O	1.97	0.48
1:A:316:GLN:HE21	1:A:347:LYS:N	2.08	0.47
1:A:196:GLU:H	1:A:196:GLU:CD	2.17	0.46
1:A:32:ASN:HD22	1:A:35:ARG:HH21	1.64	0.46
1:A:16:ASN:HD22	1:A:44:GLN:NE2	2.05	0.45
1:A:42:THR:HA	1:A:463:VAL:O	2.17	0.45
1:A:149:PHE:HA	1:A:183:SER:O	2.17	0.44
1:A:21:HIS:HD2	4:A:845:HOH:O	2.00	0.44
1:A:32:ASN:ND2	1:A:35:ARG:HH21	2.16	0.44
1:A:6:THR:HG23	1:A:489:LEU:HD21	1.99	0.44
1:A:154:ASP:OD2	3:A:603:2PG:O3	2.20	0.44
1:A:156:ASP:HB3	4:A:870:HOH:O	2.18	0.44
1:A:80:THR:HG22	1:A:84:LYS:HE2	1.99	0.43
1:A:143:LYS:HG3	1:A:177:GLY:H	1.83	0.43
1:A:185:ARG:NH1	3:A:603:2PG:H32	2.34	0.42
1:A:6:THR:HG23	1:A:489:LEU:CD2	2.49	0.42
1:A:21:HIS:HE1	4:A:908:HOH:O	2.03	0.42
3:A:603:2PG:O2P	3:A:603:2PG:H31	2.21	0.41
1:A:59:MET:SD	1:A:456:HIS:HB2	2.61	0.41
1:A:185:ARG:HH12	3:A:603:2PG:H32	1.86	0.41
1:A:83:ASN:OD1	1:A:128:HIS:HE1	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:821:HOH:O	4:A:930:HOH:O[4_445]	1.39	0.81

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	501/513 (98%)	488 (97%)	9 (2%)	4 (1%)	19 13

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	142	GLU
1	A	439	HIS
1	A	441	ASN
1	A	208	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	417/433 (96%)	391 (94%)	26 (6%)	18 13

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	THR
1	A	18	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	35	ARG
1	A	42	THR
1	A	93	GLU
1	A	94	ASN
1	A	162	LYS
1	A	185	ARG
1	A	196	GLU
1	A	200	LYS
1	A	212	THR
1	A	253	PHE
1	A	270	ASN
1	A	294	ASP
1	A	305	VAL
1	A	320	LEU
1	A	334	VAL
1	A	374	LYS
1	A	388	LEU
1	A	391	LEU
1	A	425	LYS
1	A	443	ASP
1	A	470	VAL
1	A	481	LEU
1	A	486	LEU
1	A	488	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	HIS
1	A	32	ASN
1	A	44	GLN
1	A	58	GLN
1	A	77	GLN
1	A	94	ASN
1	A	103	HIS
1	A	107	HIS
1	A	125	HIS
1	A	128	HIS
1	A	178	GLN
1	A	207	ASN
1	A	241	GLN
1	A	263	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	270	ASN
1	A	316	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	2PG	A	603	-	7,10,10	2.21	2 (28%)	8,14,14	1.62	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2PG	A	603	-	-	4/7/11/11	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	2PG	O1P-C2	-4.83	1.39	1.45
3	A	603	2PG	P-O4P	-2.72	1.44	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	603	2PG	P-O1P-C2	-3.33	115.40	123.04
3	A	603	2PG	O3P-P-O2P	2.24	119.44	110.68

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	603	2PG	C1-C2-C3-O3
3	A	603	2PG	O1P-C2-C3-O3
3	A	603	2PG	C3-C2-O1P-P
3	A	603	2PG	C2-O1P-P-O2P

There are no ring outliers.

1 monomer is involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	603	2PG	12	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	503/513 (98%)	-0.20	5 (0%) 82 81	13, 22, 38, 50	2 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	305	VAL	3.1
1	A	142	GLU	2.6
1	A	294	ASP	2.4
1	A	88	ASP	2.4
1	A	140	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	2PG	A	603	11/11	0.76	0.32	17,29,34,35	11
2	MN	A	602	1/1	1.00	0.05	9,9,9,9	0
2	MN	A	601	1/1	1.00	0.02	15,15,15,15	0

6.5 Other polymers

There are no such residues in this entry.