

wwPDB X-ray Structure Validation Summary Report (i)

Jan 4, 2024 – 04:33 pm GMT

PDB ID	:	4YDU
Title	:	Crystal structure of E. coli YgjD-YeaZ heterodimer in complex with ADP
Authors	:	Zhang, W.; Collinet, B.; Perrochia, L.; Durand, D.; Van Tilbeurgh, H.
Deposited on		
Resolution	:	2.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	343	89%	9% •
1	В	343	.% 89%	9% ••
2	С	237	% 8 9%	7% • •
2	D	237	84%	10% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard



residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit cr	rite-
ria:	

Mol	Type	Chain	\mathbf{Res}	Chirality	Geometry	Clashes	Electron density
6	ACT	А	407	-	-	Х	-
6	ACT	D	302	-	-	Х	-
6	ACT	D	306	-	-	Х	-



 $\mathbf{2}$

Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 9012 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Δ	336	Total	С	Ν	0	\mathbf{S}	0	0	0
	A	550	2470	1573	423	460	14	0	0	0
1	р	338	Total	С	Ν	0	S	0	0	0
	D	000	2498	1587	429	468	14		U	U

• Molecule 1 is a protein called tRNA N6-adenosine threonylcarbamoyltransferase.

Chain	Residue	Modelled	Actual	Comment	Reference
A	338	HIS	-	expression tag	UNP P05852
А	339	HIS	-	expression tag	UNP P05852
A	340	HIS	-	expression tag	UNP P05852
А	341	HIS	-	expression tag	UNP P05852
А	342	HIS	-	expression tag	UNP P05852
А	343	HIS	-	expression tag	UNP P05852
В	338	HIS	-	expression tag	UNP P05852
В	339	HIS	-	expression tag	UNP P05852
В	340	HIS	-	expression tag	UNP P05852
В	341	HIS	-	expression tag	UNP P05852
В	342	HIS	-	expression tag	UNP P05852
В	343	HIS	-	expression tag	UNP P05852

There are 12 discrepancies between the modelled and reference sequences:

• Molecule 2 is a protein called tRNA threonylcarbamoyladenosine biosynthesis protein TsaB.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	С	230	Total 1718	C 1091	1,	0 321	S 12	0	0	0
2	D	229	Total 1724	C 1095	1,	0 318	S 12	0	0	0


There are 12 discrepancies between the modelled and reference sequences:



4YDU

Chain	Residue	Modelled	Actual	Comment	Reference
С	232	HIS	-	expression tag	UNP P76256
С	233	HIS	-	expression tag	UNP P76256
С	234	HIS	-	expression tag	UNP P76256
С	235	HIS	-	expression tag	UNP P76256
С	236	HIS	-	expression tag	UNP P76256
С	237	HIS	-	expression tag	UNP P76256
D	232	HIS	-	expression tag	UNP P76256
D	233	HIS	-	expression tag	UNP P76256
D	234	HIS	-	expression tag	UNP P76256
D	235	HIS	-	expression tag	UNP P76256
D	236	HIS	-	expression tag	UNP P76256
D	237	HIS	-	expression tag	UNP P76256

• Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
3	А	1	Total	С	Ν	Ο	Р	0	0	
5	Π	1	27	10	5	10	2	0	0	
3	В	1	Total	С	Ν	Ο	Р	0	0	
5	D	1	27	10	5	10	2	0	0	
3	С	1	Total	С	Ν	Ο	Р	0	0	
5	U	1	27	10	5	10	2	0	U	
3	Л	1	Total	С	Ν	Ο	Р	0	0	
0	D	1	27	10	5	10	2	0	0	

• Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total Mg 1 1	0	0
4	В	1	Total Mg 1 1	0	0

• Molecule 5 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total Fe 1 1	0	0
5	В	1	Total Fe 1 1	0	0

• Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	105	Total O 105 105	0	0
7	В	124	Total O 124 124	0	0
7	С	103	Total O 103 103	0	0
7	D	106	Total O 106 106	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: tRNA N6-adenosine threonylcarbamoyltransferase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	63.62Å 68.48Å 87.07Å	Deperitor
a, b, c, α , β , γ	109.38° 92.66° 117.66°	Depositor
Resolution (Å)	49.38 - 2.33	Depositor
Resolution (A)	49.38 - 2.25	EDS
% Data completeness	97.9(49.38-2.33)	Depositor
(in resolution range)	97.9(49.38-2.25)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	0.12	Depositor
$< I/\sigma(I) > 1$	$1.75 (at 2.25 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
B B.	0.202 , 0.253	Depositor
R, R_{free}	0.173 , 0.221	DCC
R_{free} test set	2783 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	28.6	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 36.0	EDS
L-test for twinning ²	$< L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	0.045 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9012	wwPDB-VP
Average B, all atoms $(Å^2)$	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.41% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, ACT, ADP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Chain Bond lengtl		B	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.68	0/2519	0.80	2/3423~(0.1%)
1	В	0.78	2/2548~(0.1%)	0.90	5/3462~(0.1%)
2	С	0.78	1/1755~(0.1%)	0.81	4/2394~(0.2%)
2	D	1.08	6/1761~(0.3%)	0.99	10/2398~(0.4%)
All	All	0.82	9/8583~(0.1%)	0.87	21/11677~(0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	С	0	1
2	D	0	1
All	All	0	2

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	D	117	ALA	C-N	-24.97	0.76	1.34
2	С	224	TRP	C-N	15.13	1.68	1.34
2	D	104	ARG	C-N	-14.44	1.00	1.34
1	В	291	PHE	C-N	-11.68	1.07	1.34
2	D	50	GLY	C-N	-10.90	1.08	1.34

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	291	PHE	O-C-N	-14.12	100.10	122.70
1	В	291	PHE	C-N-CA	13.82	156.25	121.70



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	D	117	ALA	C-N-CA	13.49	155.43	121.70
2	D	104	ARG	O-C-N	-10.62	105.71	122.70
1	В	291	PHE	CA-C-N	9.89	138.97	117.20

Continued from previous page...

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	С	223	ALA	Mainchain
2	D	118	ARG	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2470	0	2428	26	0
1	В	2498	0	2458	20	0
2	С	1718	0	1658	13	0
2	D	1724	0	1690	20	0
3	А	27	0	12	0	0
3	В	27	0	12	0	0
3	С	27	0	12	0	0
3	D	27	0	12	0	0
4	А	1	0	0	0	0
4	В	1	0	0	0	0
5	А	1	0	0	0	0
5	В	1	0	0	0	0
6	А	16	0	12	2	0
6	В	8	0	6	1	0
6	С	8	0	6	0	0
6	D	20	0	15	5	0
7	А	105	0	0	2	0
7	В	124	0	0	2	1
7	С	103	0	0	0	0
7	D	106	0	0	2	1
All	All	9012	0	8321	77	1



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 77 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:224:TRP:C	2:C:225:LYS:N	1.68	1.42
2:D:117:ALA:O	2:D:118:ARG:N	1.59	1.33
2:D:117:ALA:CA	2:D:118:ARG:N	2.10	1.14
2:D:117:ALA:C	2:D:118:ARG:CA	2.19	1.10
2:D:79:GLN:HE22	6:D:306:ACT:H1	1.43	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:517:HOH:O	7:D:403:HOH:O[1_554]	2.10	0.10

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	334/343~(97%)	326~(98%)	8 (2%)	0	100	100
1	В	336/343~(98%)	326~(97%)	9~(3%)	1 (0%)	41	47
2	С	228/237~(96%)	221 (97%)	7 (3%)	0	100	100
2	D	227/237~(96%)	219~(96%)	7 (3%)	1 (0%)	34	38
All	All	1125/1160 (97%)	1092 (97%)	31 (3%)	2~(0%)	47	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type						
1	В	292	TYR						



Continued from previous page...

Mol	Chain	Res	Type
2	D	228	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	239/264~(90%)	237~(99%)	2(1%)	81 89
1	В	245/264~(93%)	244 (100%)	1 (0%)	91 95
2	С	165/187~(88%)	164 (99%)	1 (1%)	86 92
2	D	170/187~(91%)	170 (100%)	0	100 100
All	All	819/902~(91%)	815 (100%)	4 (0%)	88 93

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	50	ASP
1	А	292	TYR
1	В	50	ASP
2	С	168	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	D	79	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 21 ligands modelled in this entry, 4 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	gles
MOI	туре	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
6	ACT	D	306	-	$3,\!3,\!3$	0.62	0	$3,\!3,\!3$	1.52	1 (33%)
6	ACT	А	404	-	3,3,3	1.20	0	3,3,3	0.65	0
6	ACT	D	305	-	3,3,3	0.63	0	3,3,3	1.49	0
3	ADP	В	401	5	24,29,29	0.83	1 (4%)	$29,\!45,\!45$	1.44	3 (10%)
6	ACT	А	406	-	3,3,3	0.89	0	3,3,3	0.69	0
6	ACT	D	302	-	3,3,3	0.88	0	3,3,3	0.25	0
6	ACT	С	303	-	3, 3, 3	0.98	0	3, 3, 3	0.79	0
6	ACT	А	407	-	$3,\!3,\!3$	0.65	0	$3,\!3,\!3$	1.77	1 (33%)
3	ADP	D	301	-	24,29,29	1.00	1 (4%)	$29,\!45,\!45$	1.52	6 (20%)
6	ACT	В	404	-	3,3,3	0.93	0	3,3,3	1.14	0
6	ACT	D	304	-	3,3,3	0.80	0	3,3,3	1.58	1 (33%)
6	ACT	А	405	-	3,3,3	0.63	0	3,3,3	1.25	0
6	ACT	С	302	-	3,3,3	0.70	0	3,3,3	0.86	0
3	ADP	С	301	-	24,29,29	0.84	1 (4%)	29,45,45	1.43	4 (13%)
3	ADP	А	401	-	24,29,29	0.99	1 (4%)	29,45,45	1.79	10 (34%)
6	ACT	D	303	-	3,3,3	0.59	0	3,3,3	1.03	0
6	ACT	В	405	-	$3,\!3,\!3$	0.63	0	$3,\!3,\!3$	1.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings		
3	ADP	А	401	-	-	1/12/32/32	0/3/3/3		



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	С	301	-	-	2/12/32/32	0/3/3/3
3	ADP	В	401	5	-	2/12/32/32	0/3/3/3
3	ADP	D	301	-	-	1/12/32/32	0/3/3/3

Continued from previous page...

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
3	D	301	ADP	C2-N3	2.31	1.35	1.32
3	В	401	ADP	C5-C4	2.18	1.46	1.40
3	А	401	ADP	C2-N3	2.11	1.35	1.32
3	С	301	ADP	C2-N3	2.03	1.35	1.32

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	D	301	ADP	N3-C2-N1	-4.32	121.92	128.68
3	С	301	ADP	N3-C2-N1	-4.00	122.43	128.68
3	А	401	ADP	N3-C2-N1	-3.81	122.73	128.68
3	В	401	ADP	N3-C2-N1	-3.72	122.86	128.68
3	А	401	ADP	O3A-PB-O1B	-3.42	92.23	111.19

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	301	ADP	PA-O3A-PB-O2B
3	С	301	ADP	PA-O3A-PB-O3B
3	D	301	ADP	PA-O3A-PB-O3B
3	В	401	ADP	C3'-C4'-C5'-O5'
3	В	401	ADP	O4'-C4'-C5'-O5'


There are no ring outliers.

5 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	306	ACT	2	0
6	D	305	ACT	1	0
6	D	302	ACT	2	0
6	А	407	ACT	2	0
6	В	404	ACT	1	0




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and similar rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	D	4
2	С	1
1	В	1

The worst 5 of 6 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	С	224:TRP	С	225:LYS	Ν	1.68
1	D	103:TRP	С	104:ARG	Ν	1.10
1	D	50:GLY	С	51:THR	Ν	1.09
1	В	291:PHE	С	292:TYR	N	1.07
1	D	104:ARG	С	105:LYS	Ν	1.00



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	336/343~(97%)	-0.42	1 (0%) 94 97	19, 37, 68, 86	0
1	В	338/343~(98%)	-0.56	3 (0%) 84 89	19, 30, 54, 73	0
2	С	230/237~(97%)	-0.54	3 (1%) 77 83	20, 32, 62, 84	0
2	D	229/237~(96%)	-0.62	1 (0%) 92 96	18, 30, 56, 75	0
All	All	1133/1160~(97%)	-0.53	8 (0%) 87 92	18, 32, 61, 86	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	RSRZ
2	С	229	GLY	2.8
1	А	226	GLY	2.7
2	С	223	ALA	2.3
2	С	224	TRP	2.3
2	D	229	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	ACT	В	404	4/4	0.67	0.20	$41,\!51,\!53,\!59$	0
6	ACT	А	407	4/4	0.78	0.20	40,41,48,51	0
6	ACT	В	405	4/4	0.79	0.24	49,50,56,60	0
6	ACT	D	306	4/4	0.84	0.21	32,47,49,53	0
6	ACT	D	304	4/4	0.85	0.14	48,50,54,61	0
6	ACT	С	302	4/4	0.86	0.14	46,48,50,51	0
6	ACT	С	303	4/4	0.89	0.25	42,45,47,47	0
6	ACT	D	305	4/4	0.90	0.17	37,37,45,53	0
6	ACT	D	303	4/4	0.91	0.13	51,52,52,54	0
6	ACT	А	405	4/4	0.92	0.14	39,41,42,44	0
6	ACT	А	404	4/4	0.92	0.09	39,41,42,49	0
3	ADP	С	301	27/27	0.94	0.08	30,37,61,70	0
6	ACT	D	302	4/4	0.94	0.18	41,44,45,47	0
6	ACT	А	406	4/4	0.95	0.08	45,48,56,61	0
4	MG	В	402	1/1	0.96	0.05	$35,\!35,\!35,\!35$	0
3	ADP	D	301	27/27	0.96	0.07	24,29,51,60	0
4	MG	А	402	1/1	0.97	0.03	33,33,33,33	0
3	ADP	В	401	27/27	0.98	0.09	19,23,32,38	0
3	ADP	А	401	27/27	0.98	0.07	23,25,36,38	0
5	FE	В	403	1/1	1.00	0.09	23,23,23,23	0
5	FE	А	403	1/1	1.00	0.07	$25,\!25,\!25,\!25$	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











6.5 Other polymers (i)

There are no such residues in this entry.

