



wwPDB X-ray Structure Validation Summary Report

Jan 2, 2024 – 06:04 pm GMT

PDB ID : 5AH5
Title : Crystal structure of the ternary complex of Agrobacterium radiobacter K84 agnB2 LeuRS-tRNA-LeuAMS
Authors : Palencia, A.; Chopra, S.; Virus, C.; Schulwitz, S.; Temple, B.R.; Cusack, S.; Reader, J.S.
Deposited on : 2015-02-05
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)
RNA backbone	3102	1000 (2.54-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	822	
1	B	822	
2	C	84	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	84	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into segments: a small red segment at the start, followed by a large green segment labeled '65%', then a yellow segment labeled '27%', and finally a small grey segment at the end. Above the bar, there is a '%' symbol. Below the bar, there are two dots '• •'.</p>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 16819 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LEUCINE-TRNA LIGASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	789	Total	C	N	O	S	0	5	0
			6288	3990	1093	1175	30			
1	B	771	Total	C	N	O	S	0	4	0
			6146	3899	1067	1150	30			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	815	LEU	-	expression tag	UNP B9JQP8
A	816	GLU	-	expression tag	UNP B9JQP8
A	817	HIS	-	expression tag	UNP B9JQP8
A	818	HIS	-	expression tag	UNP B9JQP8
A	819	HIS	-	expression tag	UNP B9JQP8
A	820	HIS	-	expression tag	UNP B9JQP8
A	821	HIS	-	expression tag	UNP B9JQP8
A	822	HIS	-	expression tag	UNP B9JQP8
B	815	LEU	-	expression tag	UNP B9JQP8
B	816	GLU	-	expression tag	UNP B9JQP8
B	817	HIS	-	expression tag	UNP B9JQP8
B	818	HIS	-	expression tag	UNP B9JQP8
B	819	HIS	-	expression tag	UNP B9JQP8
B	820	HIS	-	expression tag	UNP B9JQP8
B	821	HIS	-	expression tag	UNP B9JQP8
B	822	HIS	-	expression tag	UNP B9JQP8


- Molecule 2 is a RNA chain called TRNA-LEU TAA ISOACCEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	80	Total	C	N	O	P	0	0	0
			1702	759	304	559	80			
2	D	81	Total	C	N	O	P	0	0	0
			1721	768	305	567	81			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 5'-O-(L-leucylsulfamoyl)adenosine (three-letter code: LSS) (formula: C₁₆H₂₅N₇O₇S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	N	O	S	0	0
			31	16	7	7	1		
4	B	1	Total	C	N	O	S	0	0
			31	16	7	7	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0

- Molecule 6 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	1	Total Mn 1 1	0	0
6	D	1	Total Mn 1 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	322	Total O 322 322	0	0
7	B	384	Total O 384 384	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	C	89	Total O 89 89	0	0
7	D	76	Total O 76 76	0	0


3 Residue-property plots i


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LEUCINE-TRNA LIGASE




• Molecule 1: LEUCINE-TRNA LIGASE






- Molecule 2: TRNA-LEU TAA ISOACCEPTOR



- Molecule 2: TRNA-LEU TAA ISOACCEPTOR



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	169.91Å 50.32Å 170.52Å 90.00° 93.48° 90.00°	Depositor
Resolution (Å)	48.58 – 2.10 48.54 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.58-2.10) 99.5 (48.54-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.194 , 0.237 0.200 , 0.241	Depositor DCC
R_{free} test set	8487 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	37.7	Xtrriage
Anisotropy	0.085	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 43.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.014 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16819	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN, LSS, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.83	0/6447	0.93	17/8760 (0.2%)
1	B	0.90	3/6299 (0.0%)	0.94	22/8559 (0.3%)
2	C	0.67	1/1899 (0.1%)	1.00	5/2953 (0.2%)
2	D	0.69	3/1920 (0.2%)	0.97	6/2985 (0.2%)
All	All	0.83	7/16565 (0.0%)	0.95	50/23257 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	G	OP3-P	-10.10	1.49	1.61
2	C	1	G	OP3-P	-9.70	1.49	1.61
2	D	22	A	O3'-P	-6.16	1.53	1.61
2	D	23	C	O3'-P	-5.86	1.54	1.61
1	B	421	ARG	CD-NE	-5.30	1.37	1.46

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	194	ARG	NE-CZ-NH2	-10.88	114.86	120.30
1	B	421	ARG	NE-CZ-NH2	-10.76	114.92	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	713	ARG	NE-CZ-NH1	9.51	125.05	120.30
1	B	20	ARG	NE-CZ-NH1	9.20	124.90	120.30
2	C	4	C	O5'-P-OP2	-8.55	98.00	105.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	531	THR	Peptide
1	B	265	ILE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6288	0	6179	39	0
1	B	6146	0	6030	47	0
2	C	1702	0	868	6	0
2	D	1721	0	876	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	31	0	25	1	0
4	B	31	0	25	0	0
5	A	5	0	0	0	0
5	B	10	0	0	0	0
5	C	5	0	0	0	0
5	D	5	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	322	0	0	5	0
7	B	384	0	0	6	0
7	C	89	0	0	1	0
7	D	76	0	0	0	0
All	All	16819	0	14003	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:646:CYS:SG	7:B:2316:HOH:O	2.26	0.93
1:A:47[A]:HIS:HB2	7:A:2045:HOH:O	1.85	0.76
1:B:713:ARG:HD2	7:B:2343:HOH:O	1.87	0.75
1:A:257:ILE:HD12	1:A:308:VAL:HG11	1.73	0.70
1:B:754:ILE:HD12	1:B:772:VAL:HG22	1.74	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	786/822 (96%)	751 (96%)	33 (4%)	2 (0%)	41	41
1	B	767/822 (93%)	735 (96%)	30 (4%)	2 (0%)	41	41
All	All	1553/1644 (94%)	1486 (96%)	63 (4%)	4 (0%)	41	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	402	LEU
1	A	356	ARG
1	B	74	PRO
1	B	335	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	684/707 (97%)	670 (98%)	14 (2%)	55	60
1	B	667/707 (94%)	652 (98%)	15 (2%)	52	57
All	All	1351/1414 (96%)	1322 (98%)	29 (2%)	55	59

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	14	GLN
1	B	712	VAL
1	B	47[A]	HIS
1	B	453	LEU
1	B	39	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	79/84 (94%)	12 (15%)	7 (8%)
2	D	79/84 (94%)	12 (15%)	3 (3%)
All	All	158/168 (94%)	24 (15%)	10 (6%)

5 of 24 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	10	G
2	C	16	U
2	C	17	C
2	C	18	G
2	C	32	C

5 of 10 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	D	9	G
2	D	46(B)	U
2	D	75	C
2	C	37	A
2	C	46(B)	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	LSS	B	1818	-	30,33,33	1.92	6 (20%)	33,49,49	1.89	6 (18%)
5	SO4	C	1078	-	4,4,4	0.44	0	6,6,6	0.45	0
5	SO4	B	1820	-	4,4,4	0.48	0	6,6,6	0.52	0
5	SO4	A	1819	-	4,4,4	0.53	0	6,6,6	0.60	0
4	LSS	A	1818	-	30,33,33	1.56	5 (16%)	33,49,49	1.85	8 (24%)
5	SO4	B	1819	-	4,4,4	0.41	0	6,6,6	0.26	0
5	SO4	D	1078	-	4,4,4	0.43	0	6,6,6	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LSS	B	1818	-	-	0/18/39/39	0/3/3/3
4	LSS	A	1818	-	-	0/18/39/39	0/3/3/3

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1818	LSS	O1A-S1	5.85	1.47	1.42
4	B	1818	LSS	S1-N2	-5.68	1.49	1.59
4	A	1818	LSS	S1-N2	-5.28	1.50	1.59
4	B	1818	LSS	O2A-S1	3.21	1.45	1.42
4	A	1818	LSS	C1-N2	-3.10	1.31	1.37

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1818	LSS	O2A-S1-O1A	-7.59	108.94	120.76
4	A	1818	LSS	O2A-S1-O1A	-4.58	113.62	120.76
4	A	1818	LSS	N3-C2-N1	-4.23	122.06	128.68
4	A	1818	LSS	O5-C25-C24	3.89	114.88	107.62
4	A	1818	LSS	O4-C21-C22	-3.84	101.31	106.93

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1818	LSS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	789/822 (95%)	0.47	88 (11%) 5 6	24, 42, 115, 147	0
1	B	771/822 (93%)	0.46	84 (10%) 5 7	21, 38, 119, 146	0
2	C	80/84 (95%)	-0.51	1 (1%) 77 80	31, 52, 106, 141	0
2	D	81/84 (96%)	-0.57	1 (1%) 79 82	29, 48, 82, 136	0
All	All	1721/1812 (94%)	0.37	174 (10%) 7 9	21, 41, 115, 147	0

The worst 5 of 174 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	393	ILE	9.9
1	B	316	ARG	8.6
1	B	309	VAL	8.6
1	A	380	SER	7.9
1	B	312	LEU	7.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	SO4	A	1819	5/5	0.84	0.17	69,70,91,94	0
6	MN	D	1077	1/1	0.86	0.06	91,91,91,91	0
5	SO4	B	1819	5/5	0.88	0.15	105,113,114,115	0
5	SO4	D	1078	5/5	0.92	0.21	80,80,83,92	0
5	SO4	B	1820	5/5	0.92	0.19	61,61,79,93	0
5	SO4	C	1078	5/5	0.95	0.17	74,76,83,84	0
6	MN	C	1077	1/1	0.96	0.08	84,84,84,84	0
4	LSS	B	1818	31/31	0.98	0.13	20,26,30,32	0
3	ZN	A	900	1/1	0.98	0.03	73,73,73,73	0
3	ZN	B	900	1/1	0.98	0.04	63,63,63,63	0
4	LSS	A	1818	31/31	0.98	0.12	22,27,32,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.