

# wwPDB X-ray Structure Validation Summary Report (i)

May 16, 2020 – 03:22 pm BST

PDB ID : 5CVB

> Title Crystal structure of the type IX collagen NC2 hetero-trimerization domain

> > with a guest fragment a1a1a1 of type I collagen

: Boudko, S.P.; Bachinger, H.P. Authors

Deposited on 2015-07-25

2.25 Å(reported) Resolution

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity 4.02b-467

> 1.8.5 (274361), CSD as541be (2020) Mogul

Xtriage (Phenix) 1.13 EDS 2.11

Percentile statistics 20191225.v01 (using entries in the PDB archive December 25th 2019)

> Refmac 5.8.0158

7.0.044 (Gargrove) CCP4 Engh & Huber (2001)

Ideal geometry (proteins) Ideal geometry (DNA, RNA) Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) 2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{aligned}  ext{Similar resolution} \ (\# ext{Entries},  ext{resolution range}( ext{Å})) \end{aligned}$
$R_{free}$	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
			8%				
1	A	71	82%		11%		7%
			27%				
1	D	71	70%	23%	ó	•	6%
			14%				
2	В	71	79%	6%	ó	15%	
			25%				
2	E	71	69% 14	%	•	15%	_
			10%				
3	С	72	88%			6%	7%
			29%				
3	F	72	82%		139	6	6%



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2786 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Collagen alpha-1(I) chain, Collagen alpha-1(IX) chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	Λ	66	Total	С	N	О	S	Se	0	0	0
1	1 A 00	00	465	292	88	81	1	3	0		
1	D	67	Total	С	N	О	S	Se	0	0	0
1	D	07	471	295	89	83	1	3	0		U

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P02452
A	2	SER	_	expression tag	UNP P02452
A	3	GLY	-	expression tag	UNP P02452
A	4	PRO	-	expression tag	UNP P02452
A	5	PRO	-	expression tag	UNP P02452
A	6	GLY	-	expression tag	UNP P02452
A	7	PRO	-	expression tag	UNP P02452
A	8	PRO	_	expression tag	UNP P02452
A	9	GLY	-	expression tag	UNP P02452
A	10	PRO	_	expression tag	UNP P02452
A	11	PRO	-	expression tag	UNP P02452
A	12	GLY	-	expression tag	UNP P02452
A	13	PRO	_	expression tag	UNP P02452
A	14	PRO	-	expression tag	UNP P02452
A	27	GLY	-	linker	UNP P02452
A	28	PRO	-	linker	UNP P02452
A	29	PRO	-	linker	UNP P02452
A	30	GLY	-	linker	UNP P02452
A	31	PRO	-	linker	UNP P02452
A	32	PRO	=	linker	UNP P02452
A	33	GLY	-	linker	UNP P02452
A	34	PRO		linker	UNP P02452
A	35	PRO	-	linker	UNP P02452
D	1	GLY	-	expression tag	UNP P02452
D	2	SER	-	expression tag	UNP P02452



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
D	3	GLY	_	expression tag	UNP P02452
D	4	PRO	-	expression tag	UNP P02452
D	5	PRO	-	expression tag	UNP P02452
D	6	GLY	-	expression tag	UNP P02452
D	7	PRO	-	expression tag	UNP P02452
D	8	PRO	_	expression tag	UNP P02452
D	9	GLY	-	expression tag	UNP P02452
D	10	PRO	-	expression tag	UNP P02452
D	11	PRO	-	expression tag	UNP P02452
D	12	GLY	-	expression tag	UNP P02452
D	13	PRO	-	expression tag	UNP P02452
D	14	PRO	-	expression tag	UNP P02452
D	27	GLY	-	linker	UNP P02452
D	28	PRO	-	linker	UNP P02452
D	29	PRO	-	linker	UNP P02452
D	30	GLY	-	linker	UNP P02452
D	31	PRO	-	linker	UNP P02452
D	32	PRO	-	linker	UNP P02452
D	33	GLY	-	linker	UNP P02452
D	34	PRO	-	linker	UNP P02452
D	35	PRO	-	linker	UNP P02452

• Molecule 2 is a protein called Collagen alpha-1(I) chain, Collagen alpha-2(IX) chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	D	60	Total	С	N	О	S	0	0	0
	2 D		399	253	70	74	2			
2	Б	60	Total	С	N	О	S	0	0	0
	Ľ	00	393	250	66	75	2	0	U	U

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	1	GLY	_	expression tag	UNP P02452
В	2	SER	-	expression tag	UNP P02452
В	3	GLY	_	expression tag	UNP P02452
В	4	PRO	-	expression tag	UNP P02452
В	5	PRO	_	expression tag	UNP P02452
В	6	GLY	_	expression tag	UNP P02452
В	7	PRO	-	expression tag	UNP P02452
В	8	PRO	_	expression tag	UNP P02452
В	9	GLY	-	expression tag	UNP P02452



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
В	10	PRO	_	expression tag	UNP P02452
В	11	PRO	_	expression tag	UNP P02452
В	12	GLY	-	expression tag	UNP P02452
В	13	PRO	_	expression tag	UNP P02452
В	14	PRO	-	expression tag	UNP P02452
В	27	GLY	_	linker	UNP P02452
В	28	PRO	-	linker	UNP P02452
В	29	PRO	-	linker	UNP P02452
В	30	GLY	-	linker	UNP P02452
В	31	PRO	-	linker	UNP P02452
В	32	PRO	-	linker	UNP P02452
В	33	GLY	-	linker	UNP P02452
В	34	PRO	-	linker	UNP P02452
В	35	PRO	-	linker	UNP P02452
Е	1	GLY	-	expression tag	UNP P02452
Е	2	SER	-	expression tag	UNP P02452
Е	3	GLY	_	expression tag	UNP P02452
Е	4	PRO	_	expression tag	UNP P02452
Е	5	PRO	_	expression tag	UNP P02452
Е	6	GLY	-	expression tag	UNP P02452
Е	7	PRO	-	expression tag	UNP P02452
Е	8	PRO	-	expression tag	UNP P02452
Е	9	GLY	-	expression tag	UNP P02452
Е	10	PRO	_	expression tag	UNP P02452
Е	11	PRO	-	expression tag	UNP P02452
Е	12	GLY	-	expression tag	UNP P02452
Е	13	PRO	-	expression tag	UNP P02452
Е	14	PRO	-	expression tag	UNP P02452
Е	27	GLY	_	linker	UNP P02452
Е	28	PRO	-	linker	UNP P02452
Е	29	PRO	-	linker	UNP P02452
Е	30	GLY	-	linker	UNP P02452
Е	31	PRO	-	linker	UNP P02452
Е	32	PRO	-	linker	UNP P02452
Е	33	GLY	-	linker	UNP P02452
Е	34	PRO	-	linker	UNP P02452
Е	35	PRO	_	linker	UNP P02452

 $\bullet$  Molecule 3 is a protein called Collagen alpha-1(I) chain, Collagen alpha-3(IX) chain.  $Continued\ on\ next\ page...$ 



 $Continued\ from\ previous\ page...$ 

Mol	Chain Residues	Atoms	ZeroOcc	AltConf	Trace
-----	----------------	-------	---------	---------	-------

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	С	67	Total	С	N	О	S	0	1	0	
) 	3   0	07	462	290	87	82	3				
9	D.	69	Total	С	N	О	S	0	1	0	
3	3   F	F 68		295	88	83	4	0	1		

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
С	1	GLY	-	expression tag	UNP P02452
С	2	SER	_	expression tag	UNP P02452
С	3	GLY	-	expression tag	UNP P02452
С	4	PRO	-	expression tag	UNP P02452
С	5	PRO	-	expression tag	UNP P02452
С	6	GLY	-	expression tag	UNP P02452
С	7	PRO	_	expression tag	UNP P02452
С	8	PRO	-	expression tag	UNP P02452
С	9	GLY	-	expression tag	UNP P02452
С	10	PRO	-	expression tag	UNP P02452
С	11	PRO	-	expression tag	UNP P02452
С	12	GLY	-	expression tag	UNP P02452
С	13	PRO	-	expression tag	UNP P02452
С	14	PRO	-	expression tag	UNP P02452
С	27	GLY	-	linker	UNP P02452
С	28	PRO	-	linker	UNP P02452
С	29	PRO	_	linker	UNP P02452
С	30	GLY	-	linker	UNP P02452
С	31	PRO	-	linker	UNP P02452
С	32	PRO	-	linker	UNP P02452
С	33	GLY	-	linker	UNP P02452
С	34	PRO	-	linker	UNP P02452
С	35	PRO	-	linker	UNP P02452
F	1	GLY	-	expression tag	UNP P02452
F	2	SER	-	expression tag	UNP P02452
F	3	GLY	-	expression tag	UNP P02452
F	4	PRO	-	expression tag	UNP P02452
F	5	PRO	-	expression tag	UNP P02452
F	6	GLY	-	expression tag	UNP P02452
F	7	PRO	-	expression tag	UNP P02452
F	8	PRO	-	expression tag	UNP P02452
F	9	GLY	-	expression tag	UNP P02452
F	10	PRO	-	expression tag	UNP P02452



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
F	11	PRO	=	expression tag	UNP P02452
F	12	GLY	-	expression tag	UNP P02452
F	13	PRO	-	expression tag	UNP P02452
F	14	PRO	-	expression tag	UNP P02452
F	27	GLY	-	linker	UNP P02452
F	28	PRO	_	linker	UNP P02452
F	29	PRO	-	linker	UNP P02452
F	30	GLY	_	linker	UNP P02452
F	31	PRO	-	linker	UNP P02452
F	32	PRO	-	linker	UNP P02452
F	33	GLY	=	linker	UNP P02452
F	34	PRO	-	linker	UNP P02452
F	35	PRO	-	linker	UNP P02452

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	С	1	Total C O 6 3 3	0	0
4	F	1	Total C O 6 3 3	0	0
4	F	1	Total C O 6 3 3	0	0
4	F	1	Total C O 6 3 3	0	0



### • Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	13	Total O 13 13	0	0
5	В	15	Total O 15 15	0	0
5	С	25	Total O 25 25	0	0
5	D	16	Total O 16 16	0	0
5	E	8	Total O 8 8	0	0
5	F	19	Total O 19 19	0	0



# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Collagen alpha-1(I) chain, Collagen alpha-1(IX) chain



• Molecule 3: Collagen alpha-1(I) chain, Collagen alpha-3(IX) chain







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	51.77Å 64.46Å 65.39Å	Danagitan
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $112.72^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	47.75 - 2.25	Depositor
Resolution (A)	47.75 - 2.25	EDS
% Data completeness	88.2 (47.75-2.25)	Depositor
(in resolution range)	88.1 (47.75-2.25)	EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.73 (at 2.24Å)	Xtriage
Refinement program	PHENIX 1.9-1692	Depositor
D D.	0.227 , 0.278	Depositor
$R, R_{free}$	0.229 , $0.278$	DCC
$R_{free}$ test set	1728 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.6	Xtriage
Anisotropy	0.603	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36 , 50.6	EDS
L-test for twinning <sup>2</sup>	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.032 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	2786	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.70% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bo	nd angles
MIOI	Wioi Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.19	0/482	0.40	0/655
1	D	0.28	0/488	0.60	1/663 (0.2%)
2	В	0.20	0/415	0.40	0/574
2	E	0.33	0/409	0.64	0/567
3	С	0.20	0/482	0.38	0/657
3	F	0.25	0/490	0.50	0/667
All	All	0.25	0/2766	0.50	1/3783 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	D	6	GLY	C-N-CD	-5.13	109.31	120.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	465	0	454	7	0
1	D	471	0	459	16	0
2	В	399	0	385	4	0
2	Ε	393	0	372	7	0



~ · · 1	e	•	
Continued	trom	nremous	naae
-	110110	produdas	payc

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	С	462	0	464	4	0
3	F	470	0	476	5	0
4	A	6	0	8	0	0
4	С	6	0	8	0	0
4	F	18	0	24	0	0
5	A	13	0	0	2	0
5	В	15	0	0	0	0
5	С	25	0	0	0	0
5	D	16	0	0	3	0
5	E	8	0	0	0	0
5	F	19	0	0	0	0
All	All	2786	0	2650	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{ll}  ext{Interatomic} \  ext{distance} \ ( ext{\AA}) \end{array}$	$egin{array}{c}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{array}$
1:D:13:PRO:HG2	2:E:14:PRO:HA	1.56	0.88
1:D:17:ARG:NH1	5:D:101:HOH:O	2.29	0.64
1:D:17:ARG:NH1	3:F:18:GLY:HA2	2.11	0.64
1:D:10:PRO:O	2:E:11:PRO:HA	1.97	0.64
1:A:50:ARG:NH1	5:A:201:HOH:O	2.34	0.59

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$64/71 \; (90\%)$	63 (98%)	1 (2%)	0	100	100



Continued	trom	meaningile	naaa
-	110116	DICUIUU	Duuc

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	D	65/71 (92%)	60 (92%)	2 (3%)	3 (5%)	2 0
2	В	58/71 (82%)	58 (100%)	0	0	100 100
2	E	58/71 (82%)	52 (90%)	3 (5%)	3 (5%)	2 0
3	С	$66/72 \; (92\%)$	65 (98%)	1 (2%)	0	100 100
3	F	$67/72 \ (93\%)$	64 (96%)	1 (2%)	2 (3%)	4 1
All	All	378/428 (88%)	362 (96%)	8 (2%)	8 (2%)	7 2

#### 5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	F	16	ALA
1	D	12	GLY
2	Е	8	PRO
1	D	7	PRO
1	D	11	PRO

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	48/47~(102%)	48 (100%)	0	100	100	
1	D	49/47 (104%)	49 (100%)	0	100	100	
2	В	40/48 (83%)	40 (100%)	0	100	100	
2	E	39/48 (81%)	38 (97%)	1 (3%)	46	52	
3	С	46/49 (94%)	46 (100%)	0	100	100	
3	F	47/49 (96%)	46 (98%)	1 (2%)	53	60	
All	All	269/288 (93%)	267 (99%)	2 (1%)	84	88	

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type	
2	Ε	42	GLN	



Continued from previous page...

Mol	Chain	Res	Type
3	F	17	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Res		Link	Bond lengths			Bond angles			
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	101	-	5,5,5	0.36	0	5, 5, 5	0.29	0
4	GOL	F	102	_	5,5,5	0.36	0	5, 5, 5	0.27	0
4	GOL	F	101	-	5,5,5	0.36	0	5,5,5	0.30	0
4	GOL	С	101	-	5,5,5	0.35	0	5, 5, 5	0.32	0
4	GOL	F	103	_	5,5,5	0.35	0	5, 5, 5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



,_,	means	nο	outliers	$\alpha f$	that	kind	Word	identified.
_	шсань	110	Outhers	OI	u	MIIIU	MCIC	idenuned.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	101	_	-	2/4/4/4	_
4	GOL	F	102	_	-	4/4/4/4	_
4	GOL	F	101	-	-	4/4/4/4	-
4	GOL	С	101	-	-	2/4/4/4	-
4	GOL	F	103	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	102	GOL	O1-C1-C2-C3
4	F	102	GOL	C1-C2-C3-O3
4	F	101	GOL	O1-C1-C2-C3
4	С	101	GOL	O1-C1-C2-C3
4	F	103	GOL	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	63/71 (88%)	0.68	6 (9%) 8 7	18, 32, 65, 77	0
1	D	$64/71 \; (90\%)$	2.12	19 (29%) 0 0	17, 37, 111, 118	0
2	В	60/71 (84%)	0.91	10 (16%) 1 1	15, 34, 71, 86	0
2	E	60/71 (84%)	1.88	18 (30%) 0 0	21, 41, 108, 112	0
3	С	67/72 (93%)	0.75	7 (10%) 6 5	18, 33, 71, 85	0
3	F	68/72 (94%)	1.91	21 (30%) 0 0	18, 39, 107, 113	0
All	All	382/428 (89%)	1.38	81 (21%) 0 0	15, 36, 106, 118	0

The worst 5 of 81 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	10	PRO	14.0
2	Ε	2	SER	14.0
1	D	10	PRO	12.9
1	D	6	GLY	11.0
1	D	5	PRO	11.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.



### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q < 0.9
4	GOL	F	101	6/6	0.82	0.21	28,40,41,43	0
4	GOL	С	101	6/6	0.84	0.14	37,44,49,51	0
4	GOL	F	102	6/6	0.89	0.25	31,35,44,52	0
4	GOL	A	101	6/6	0.90	0.21	34,38,40,42	0
4	GOL	F	103	6/6	0.92	0.24	42,44,47,48	0

### 6.5 Other polymers (i)

There are no such residues in this entry.

