



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 05:23 am BST

PDB ID : 5DV7
Title : Crystal Structure of NF90 tandem dsRBDs with dsRNA
Authors : Jayachandran, U.; Grey, H.; Cook, A.G.
Deposited on : 2015-09-21
Resolution : 3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)
RNA backbone	3102	1002 (4.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	18	 11% 83% 6%
2	B	18	 28% 72%
3	C	700	 13% 7% 80%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1637 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(*CP*CP*AP*GP*CP*AP*UP*UP*AP*UP*GP*AP*AP*AP*GP*UP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	18	383	173	72	121	17	0	0	0

- Molecule 2 is a RNA chain called RNA (5'-R(*UP*CP*AP*CP*UP*UP*UP*CP*AP*UP*AP*AP*UP*GP*CP*UP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	18	374	169	61	127	17	0	0	0

- Molecule 3 is a protein called Interleukin enhancer-binding factor 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	137	879	548	156	171	4	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total O 1 1	0	0

F443	THR
E444	PRO
A445	GLY
T452	VAL
A453	ASN
T452	ALA
K523	GLY
H524	GLY
G525	PRO
K526	LYS
N527	PHE
P528	ALA
I544	ALA
L544	LYS
T547	PRO
G548	HIS
G549	ASN
K553	PRO
R554	GLY
F555	MET
V556	GLY
M557	GLY
E558	TYR
V559	PRO
E560	MET
V561	HIS
D562	ASN
G563	GLU
Q564	VAL
K565	PRO
F566	PRO
A569	PRO
G570	ASN
S571	ILE
M572	ARG
V575	GLY
V575	ALA
Y579	GLY
A584	ASN
F589	ILE
P590	ARG
ASP	GLY
THR	ARG
PRO	GLY
SER	ARG
SER	GLY
SER	ARG
VAL	ARG
PHE	GLY
PRO	PHE
PRO	GLY
ASP	GLY
ALA	ALA
THR	ASN
THR	HIS
GLU	GLY

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 1 2	Depositor
Cell constants a, b, c, α , β , γ	100.10Å 100.10Å 108.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.41 – 3.50 45.41 – 3.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.41-3.50) 100.0 (45.41-3.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 3.48Å)	Xtrriage
Refinement program	PHENIX 1.7_650	Depositor
R, R_{free}	0.239 , 0.273 0.239 , 0.264	Depositor DCC
R_{free} test set	368 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å ²)	127.0	Xtrriage
Anisotropy	0.037	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 113.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.057 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1637	wwPDB-VP
Average B, all atoms (Å ²)	132.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.77	1/429 (0.2%)	1.02	1/667 (0.1%)
2	B	0.76	0/416	1.01	0/645
3	C	0.52	0/893	0.68	1/1222 (0.1%)
All	All	0.65	1/1738 (0.1%)	0.87	2/2534 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	3	A	N9-C4	-5.09	1.34	1.37

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	519	PRO	N-CA-CB	5.80	110.26	103.30
1	A	3	A	C8-N9-C4	5.51	108.00	105.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	383	0	196	22	0
2	B	374	0	193	16	0
3	C	879	0	715	38	0
4	C	1	0	0	0	0
All	All	1637	0	1104	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:544:ILE:HD11	3:C:558:GLU:CB	1.92	1.00
2:B:18:G:O2'	3:C:549:GLY:O	1.89	0.90
3:C:561:VAL:N	3:C:564:GLN:O	2.06	0.87
3:C:561:VAL:O	3:C:563:GLY:N	2.09	0.85
3:C:560:GLU:HA	3:C:565:LYS:HA	1.60	0.82
3:C:555:PHE:HE1	3:C:571:SER:O	1.65	0.80
1:A:1:C:HO5'	1:A:1:C:H6	1.37	0.73
3:C:544:ILE:HB	3:C:556:VAL:O	1.88	0.73
3:C:416:LEU:HD23	3:C:416:LEU:C	2.09	0.72
3:C:416:LEU:HD23	3:C:417:GLN:N	2.07	0.69
2:B:15:C:O2'	2:B:16:U:H5'	1.92	0.69
1:A:1:C:C2'	1:A:2:C:H5'	2.24	0.67
3:C:445:ALA:HB3	3:C:457:VAL:CG2	2.26	0.66
3:C:572:ASN:HB2	3:C:575:VAL:HG23	1.76	0.66
3:C:527:ASN:OD1	3:C:528:PRO:HD2	1.97	0.65
3:C:520:ILE:HG13	3:C:579:TYR:CE1	2.32	0.65
2:B:16:U:H2'	2:B:17:G:C8	2.36	0.61
2:B:7:U:H2'	2:B:8:C:C6	2.36	0.61
2:B:16:U:H2'	2:B:17:G:H8	1.65	0.60
2:B:8:C:H2'	2:B:9:A:H8	1.66	0.60
2:B:8:C:H2'	2:B:9:A:C8	2.35	0.60
2:B:16:U:O2'	2:B:17:G:H5'	2.01	0.60
3:C:561:VAL:HG12	3:C:562:ASP:OD1	2.03	0.58
3:C:555:PHE:CE1	3:C:571:SER:O	2.53	0.58
3:C:443:PHE:N	3:C:443:PHE:CD1	2.72	0.57
1:A:1:C:H2'	1:A:2:C:H5'	1.87	0.56
2:B:5:U:H2'	2:B:6:U:H6	1.72	0.55
1:A:9:A:H2'	1:A:10:U:C6	2.41	0.54
1:A:1:C:O5'	1:A:1:C:H6	1.90	0.54
3:C:445:ALA:HB3	3:C:457:VAL:HG23	1.88	0.54
3:C:445:ALA:HB3	3:C:457:VAL:HG22	1.90	0.54
3:C:522:THR:O	3:C:525:GLY:N	2.42	0.52
2:B:7:U:H2'	2:B:8:C:H6	1.74	0.51
1:A:13:A:O5'	1:A:13:A:H8	1.92	0.51
1:A:17:G:H2'	1:A:18:A:C8	2.46	0.50
3:C:426:PRO:HB2	3:C:428:HIS:CE1	2.47	0.50
3:C:555:PHE:O	3:C:569:ALA:HA	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:416:LEU:HD22	3:C:418:TYR:CE1	2.46	0.50
3:C:523:LYS:C	3:C:525:GLY:H	2.14	0.50
1:A:5:C:O2'	1:A:6:A:H5'	2.12	0.49
2:B:11:A:O2'	2:B:12:A:H5'	2.12	0.49
2:B:8:C:O2'	2:B:9:A:H5'	2.13	0.48
3:C:589:PHE:HB2	3:C:590:PRO:HD3	1.94	0.47
3:C:418:TYR:HA	3:C:436:VAL:HG22	1.95	0.47
3:C:423:GLN:O	3:C:423:GLN:HG2	2.14	0.47
1:A:17:G:H2'	1:A:18:A:H8	1.80	0.47
3:C:452:THR:O	3:C:453:ALA:C	2.53	0.47
2:B:5:U:H2'	2:B:6:U:C6	2.50	0.47
3:C:416:LEU:CD2	3:C:416:LEU:C	2.83	0.46
1:A:12:A:H2'	1:A:13:A:C8	2.50	0.46
3:C:405:ALA:HB1	3:C:458:ALA:CB	2.46	0.45
3:C:559:VAL:HG13	3:C:559:VAL:O	2.15	0.45
1:A:4:G:N2	2:B:16:U:C2	2.86	0.43
1:A:4:G:H2'	1:A:5:C:C6	2.53	0.43
3:C:547:THR:O	3:C:553:LYS:HA	2.19	0.43
2:B:17:G:O2'	2:B:18:G:H5'	2.18	0.43
3:C:435:SER:HA	3:C:443:PHE:O	2.19	0.42
3:C:589:PHE:CB	3:C:590:PRO:HD3	2.49	0.42
1:A:11:G:H2'	1:A:12:A:C8	2.54	0.42
2:B:3:A:H2'	2:B:4:C:H6	1.85	0.42
1:A:9:A:H2'	1:A:10:U:H6	1.85	0.41
3:C:566:PHE:CB	3:C:584:ALA:HB2	2.51	0.41
1:A:13:A:H2'	1:A:14:A:C8	2.56	0.41
3:C:421:ILE:HD11	3:C:435:SER:OG	2.20	0.41
1:A:12:A:O5'	1:A:12:A:H8	2.03	0.41
3:C:445:ALA:CB	3:C:457:VAL:HG22	2.51	0.41
1:A:12:A:H2'	1:A:13:A:H8	1.86	0.41
1:A:6:A:H2'	1:A:7:U:C6	2.56	0.41
3:C:562:ASP:CG	3:C:563:GLY:N	2.74	0.40
1:A:4:G:C6	1:A:5:C:N4	2.90	0.40
1:A:13:A:H2'	1:A:14:A:H8	1.87	0.40
1:A:15:G:H8	1:A:15:G:O5'	2.04	0.40
1:A:10:U:H6	1:A:10:U:O5'	2.04	0.40
3:C:437:GLU:CB	3:C:442:ASN:OD1	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	C	133/700 (19%)	120 (90%)	12 (9%)	1 (1%)	19 58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	562	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	C	65/575 (11%)	65 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	527	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	17/18 (94%)	1 (5%)	0
2	B	17/18 (94%)	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
All	All	34/36 (94%)	1 (2%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	2	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	18/18 (100%)	-0.56	0 100 100	98, 123, 215, 218	0
2	B	18/18 (100%)	-0.51	0 100 100	84, 123, 212, 228	0
3	C	137/700 (19%)	-0.34	0 100 100	75, 126, 183, 200	0
All	All	173/736 (23%)	-0.38	0 100 100	75, 125, 193, 228	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.