

wwPDB X-ray Structure Validation Summary Report (i)

Sep 19, 2023 – 12:49 PM EDT

PDB ID : 5E29

Title : Crystal Structure of Deoxygenated Hemoglobin in Complex with an Allosteric

Effector and Nitric Oxide

Authors: Safo, M.K.; Deshpande, T.M.

Deposited on : 2015-09-30

Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	141	90%	10%
1	С	141	86%	14%
2	В	145	75%	24% •
2	D	145	86%	14%



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 5613 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	141	10001	С	- '	О	S	0	0	0
1	11	111	1069	685	187	194	3	Ü	0	
1	С	141	Total	С	N	Ο	S	0	0	0
	141	1069	685	187	194	3	0	U	. 0	

• Molecule 2 is a protein called Hemoglobin subunit beta.

Mol	Chain	Residues	\mathbf{Atoms}			ZeroOcc	AltConf	Trace		
9	D	145	Total	С	N	О	S	0	0	0
	140	1116	719	194	200	3	0	U	U	
9	D	1.45	Total	С	N	О	S	0	0	0
2	2 D	145	1116	719	194	200	3	U	U	U


• Molecule 3 is NITRIC OXIDE (three-letter code: NO) (formula: NO).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total N O 2 1 1	0	0
3	С	1	Total N O 2 1 1	0	0


• Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
4	4 A	1	Total	С	Fe	N	О	0	0	
4		1	43	34	1	4	4		U	
4	В	1	Total	С	Fe	N	О	0	0	
4	4 D	1	43	34	1	4	4			
4	C	1	Total	С	Fe	N	О	0	0	
4	• C	1	43	34	1	4	4	0	U	
1	D	1	Total	С	Fe	N	О	0	0	
4	$\mid D \mid$	1	43	34	1	4	4	U	0	


• Molecule 5 is 2-(nitrooxy)ethyl 2-(4- $\{2-[(3,5-dimethylphenyl)amino]-2-oxoethyl\}$ phenoxy)-2 -methylpropanoate (three-letter code: 5JN) (formula: $C_{22}H_{26}N_2O_7$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	Δ	1	Total	С	N	О	0	1
	Λ	1	31	22	2	7	U	1


 \bullet Molecule 6 is 2-{4-[(3,5-DIMETHYLANILINO)-CARBONYL-METHYL]-PHENOXY}-2-M ETHYLPROPIONIC ACID (three-letter code: RQ3) (formula: $C_{20}H_{23}NO_4).$



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total 25	_		_	0	0
6	В	1	Total 25	C 20		O 4	0	1




 \bullet Molecule 7 is NITRATE ION (three-letter code: NO3) (formula: NO3).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	В	1	Total N O 4 1 3	0	0
7	D	1	Total N O 4 1 3	0	0

 \bullet Molecule 8 is CACODYLATE ION (three-letter code: CAC) (formula: $\mathrm{C_2H_6AsO_2}).$



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	D	1	Total 5	As 1	C 2	O 2	0	0



• Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	221	Total O 221 221	0	0
9	В	220	Total O 220 220	0	0
9	С	262	Total O 262 262	0	0
9	D	270	Total O 270 270	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hemoglobin subunit alpha





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	95.88Å 98.42Å 65.44Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.54 - 1.85	Depositor
Resolution (A)	29.54 - 1.85	EDS
% Data completeness	98.1 (29.54-1.85)	Depositor
(in resolution range)	98.2 (29.54-1.85)	EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.96 (at 1.85Å)	Xtriage
Refinement program	CNS	Depositor
D D	0.198 , 0.253	Depositor
R, R_{free}	0.194 , 0.250	DCC
R_{free} test set	2669 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.30 , 60.4	EDS
L-test for twinning ²	$< L >=0.38, < L^2>=0.21$	Xtriage
Estimated twinning fraction	0.045 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5613	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.51% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: RQ3, NO3, HEM, NO, 5JN, CAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.39	0/1097	0.56	0/1491	
1	С	0.42	0/1097	0.59	0/1491	
2	В	0.40	0/1146	0.57	0/1556	
2	D	0.44	0/1146	0.59	0/1556	
All	All	0.41	0/4486	0.58	0/6094	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1073	11	0
1	С	1069	0	1073	18	0
2	В	1116	0	1106	36	0
2	D	1116	0	1106	18	0
3	A	2	0	0	0	0
3	С	2	0	0	0	0
4	A	43	0	30	0	0
4	В	43	0	30	3	0
4	С	43	0	30	0	0

Continued on next page...



I 'omtamalod	tmom	mmonia	maaa
Continued	110111	DIEUIUU	DUJUE
00,000,000	.,	p	p = 9 =

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	43	0	30	2	0
5	A	31	0	19	2	0
6	A	25	0	22	0	0
6	В	25	0	17	1	0
7	В	4	0	0	1	0
7	D	4	0	0	0	0
8	D	5	0	0	1	0
9	A	221	0	0	2	0
9	В	220	0	0	11	0
9	С	262	0	0	8	0
9	D	270	0	0	8	0
All	All	5613	0	4536	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 82 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
2:B:13:ALA:HB3	9:B:301:HOH:O	1.58	1.03
1:C:56:LYS:HE2	9:C:476:HOH:O	1.82	0.79
2:B:143:HIS:HD2	2:B:144:LYS:NZ	1.82	0.76
2:B:43:GLU:HG3	9:B:305:HOH:O	1.88	0.74
2:B:97:HIS:NE2	7:B:203:NO3:O2	2.21	0.74

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	lowed Outliers			
1	A	139/141 (99%)	134 (96%)	5 (4%)	0	100	100	

Continued on next page...



~ · · 1	c		
Continued	trom	nremous	naae.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	С	139/141 (99%)	135 (97%)	4 (3%)	0	100	100
2	В	143/145 (99%)	138 (96%)	5 (4%)	0	100	100
2	D	143/145 (99%)	140 (98%)	3 (2%)	0	100	100
All	All	564/572 (99%)	547 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	113/113 (100%)	113 (100%)	0	100	100	
1	\mathbf{C}	113/113 (100%)	113 (100%)	0	100	100	
2	В	117/117 (100%)	115 (98%)	2 (2%)	60	47	
2	D	117/117 (100%)	117 (100%)	0	100	100	
All	All	460/460 (100%)	458 (100%)	2 (0%)	91	89	

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	В	2	HIS
2	В	19	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
2	В	139	ASN
2	В	143	HIS
2	D	143	HIS
2	D	77	HIS
2	D	80	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Т	Clasia	Das	Link	В	ond leng	$_{ m gths}$	Bond angles		
Mol	Type	Chain	Res	ites Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HEM	A	202	3	41,50,50	1.43	7 (17%)	45,82,82	0.98	3 (6%)
6	RQ3	В	202[B]	-	26,26,26	2.28	12 (46%)	37,37,37	1.59	3 (8%)
8	CAC	D	202	-	0,4,4	-	-	0,6,6	-	-
4	HEM	D	201	2	41,50,50	1.52	8 (19%)	45,82,82	1.27	5 (11%)
4	HEM	В	201	2	41,50,50	1.21	5 (12%)	45,82,82	1.20	4 (8%)
6	RQ3	A	204	-	26,26,26	2.96	10 (38%)	37,37,37	1.59	6 (16%)
3	NO	С	201	4	0,1,1	-	-	-		
5	5JN	A	203[A]	-	30,32,32	1.99	13 (43%)	39,44,44	1.35	3 (7%)
7	NO3	В	203	-	1,3,3	3.61	1 (100%)	0,3,3	-	-
4	HEM	С	202	3	41,50,50	1.34	7 (17%)	45,82,82	1.15	3 (6%)
3	NO	A	201	4	0,1,1	-	-	-		
7	NO3	D	203	_	1,3,3	3.61	1 (100%)	0,3,3	_	_

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEM	A	202	3	-	4/12/54/54	-
6	RQ3	В	202[B]	-	-	4/19/19/19	0/2/2/2
4	HEM	D	201	2	-	5/12/54/54	-
4	HEM	В	201	2	-	6/12/54/54	-
6	RQ3	A	204	-	-	0/19/19/19	0/2/2/2
5	5JN	A	203[A]	_	-	11/24/26/26	0/2/2/2
4	HEM	С	202	3	-	3/12/54/54	-

The worst 5 of 64 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
6	A	204	RQ3	C13-C11	8.41	1.57	1.38
6	В	202[B]	RQ3	C10-C9	5.14	1.48	1.38
6	A	204	RQ3	C9-C8	4.98	1.48	1.38
6	A	204	RQ3	C22-C18	4.86	1.47	1.39
6	A	204	RQ3	C14-C15	-4.48	1.42	1.51

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
6	A	204	RQ3	C4-O7-C8	6.16	130.84	120.95
6	В	202[B]	RQ3	C4-O7-C8	5.62	129.97	120.95
6	В	202[B]	RQ3	O7-C4-C1	5.33	117.26	109.99
5	A	203[A]	5JN	C20-O19-C16	4.94	128.87	120.95
4	С	202	HEM	C4B-CHC-C1C	3.81	127.59	122.56

There are no chirality outliers.

5 of 33 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	В	201	HEM	C2B-C3B-CAB-CBB
5	A	203[A]	5JN	C21-C20-C23-O24
5	A	203[A]	5JN	C21-C20-C23-O25
5	A	203[A]	5JN	O19-C20-C23-O24
5	A	203[A]	5JN	O19-C20-C23-O25

There are no ring outliers.


6 monomers are involved in 10 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	В	202[B]	RQ3	1	0
8	D	202	CAC	1	0
4	D	201	HEM	2	0
4	В	201	HEM	3	0
5	A	203[A]	5JN	2	0
7	В	203	NO3	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






















5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2	$OWAB(A^2)$	Q < 0.9
1	A	141/141 (100%)	-0.52	0 100 100	13, 21, 32, 40	0
1	С	141/141 (100%)	-0.64	0 100 100	12, 19, 28, 39	0
2	В	145/145 (100%)	-0.33	1 (0%) 87 88	15, 25, 41, 60	0
2	D	145/145 (100%)	-0.50	1 (0%) 87 88	12, 19, 34, 60	0
All	All	572/572 (100%)	-0.50	2 (0%) 94 93	12, 21, 35, 60	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	2	HIS	3.9
2	В	2	HIS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ m B ext{-}factors(\AA^2)$	Q<0.9
5	5JN	A	203[A]	31/31	0.75	0.27	31,34,41,48	31
6	RQ3	В	202[B]	25/25	0.75	0.27	29,33,40,47	25
6	RQ3	A	204	25/25	0.80	0.24	31,40,47,50	25
8	CAC	D	202	5/5	0.86	0.14	93,93,95,100	0
7	NO3	В	203	4/4	0.89	0.19	45,46,53,59	0
7	NO3	D	203	4/4	0.90	0.15	25,41,49,49	0
4	HEM	В	201	43/43	0.97	0.08	15,25,42,44	0
4	HEM	A	202	43/43	0.98	0.07	7,18,29,39	0
4	HEM	С	202	43/43	0.98	0.07	10,16,37,44	0
4	HEM	D	201	43/43	0.98	0.08	10,15,22,25	0
3	NO	A	201	2/2	0.99	0.09	17,17,17,19	0
3	NO	С	201	2/2	1.00	0.04	17,17,17,19	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


























6.5 Other polymers (i)

There are no such residues in this entry.

