



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2020 – 02:42 PM BST

PDB ID : 5EHD
Title : Crystal structure of human nucleophosmin-core in complex with cytochrome c
Authors : Bernardo-Garcia, N.; Hermoso, J.A.; Gonzalez-Arzola, K.; Diaz-Moreno, I.;
De la Rosa, M.A.
Deposited on : 2015-10-28
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.13
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1284 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	116	 8% 86% 5% 9%
1	B	116	 12% 86% 5% 9%
1	C	116	 10% 89% 5% 8%
1	D	116	 7% 89% 5% 9%
1	E	116	 9% 88% 5% 9%
1	F	116	 5% 89% 5% 8%
1	G	116	 5% 88% 5% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	116	<p>3% 85% 7% 8%</p>
1	I	116	<p>4% 85% 7% 8%</p>
1	J	116	<p>6% 88% 7% 8%</p>
1	a	116	<p>6% 90% 7% 9%</p>
1	b	116	<p>8% 85% 7% 9%</p>
1	c	116	<p>6% 84% 8% 9%</p>
1	d	116	<p>7% 84% 8% 8%</p>
1	e	116	<p>2% 89% 7% 8%</p>
1	f	116	<p>6% 84% 7% 9%</p>
1	g	116	<p>7% 87% 7% 9%</p>
1	h	116	<p>3% 85% 5% 9%</p>
1	i	116	<p>7% 87% 5% 8%</p>
1	j	116	<p>6% 83% 9% 9%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 16452 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleophosmin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	106	807	511	137	155	4	0	0	0
1	B	105	800	506	136	154	4	0	0	0
1	C	107	818	517	141	156	4	0	0	0
1	D	105	800	506	136	154	4	0	0	0
1	E	106	807	511	137	155	4	0	0	0
1	F	107	818	517	141	156	4	0	0	0
1	G	106	807	511	137	155	4	0	0	0
1	H	107	818	517	141	156	4	0	0	0
1	I	107	818	517	141	156	4	0	0	0
1	J	107	816	516	138	158	4	0	0	0
1	a	106	807	511	137	155	4	0	0	0
1	b	105	800	506	136	154	4	0	0	0
1	c	106	807	511	137	155	4	0	0	0
1	d	107	818	517	141	156	4	0	0	0
1	e	107	818	517	141	156	4	0	0	0
1	f	106	807	511	137	155	4	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	g	106	Total	C	N	O	S	0	0	0
			807	511	137	155	4			
1	h	106	Total	C	N	O	S	0	0	0
			811	512	140	155	4			
1	i	107	Total	C	N	O	S	0	0	0
			818	517	141	156	4			
1	j	106	Total	C	N	O	S	0	0	0
			811	512	140	155	4			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Cl	0	0
			1	1		
2	b	1	Total	Cl	0	0
			1	1		
2	A	1	Total	Cl	0	0
			1	1		
2	f	1	Total	Cl	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	6	Total	O	0	0
			6	6		
3	B	7	Total	O	0	0
			7	7		
3	C	1	Total	O	0	0
			1	1		
3	D	6	Total	O	0	0
			6	6		
3	E	9	Total	O	0	0
			9	9		
3	F	20	Total	O	0	0
			20	20		
3	G	14	Total	O	0	0
			14	14		
3	H	15	Total	O	0	0
			15	15		
3	I	17	Total	O	0	0
			17	17		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	J	12	Total O 12 12	0	0
3	a	8	Total O 8 8	0	0
3	b	5	Total O 5 5	0	0
3	c	8	Total O 8 8	0	0
3	d	16	Total O 16 16	0	0
3	e	14	Total O 14 14	0	0
3	f	12	Total O 12 12	0	0
3	g	18	Total O 18 18	0	0
3	h	12	Total O 12 12	0	0
3	i	16	Total O 16 16	0	0
3	j	19	Total O 19 19	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin




- Molecule 1: Nucleophosmin




- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin




- Molecule 1: Nucleophosmin




- Molecule 1: Nucleophosmin




- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin





- Molecule 1: Nucleophosmin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.80Å 179.72Å 104.30Å 90.00° 94.32° 90.00°	Depositor
Resolution (Å)	45.53 – 2.55 45.53 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.5 (45.53-2.55) 99.6 (45.53-2.55)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.54Å)	Xtrriage
Refinement program	PHENIX (1.10_2155: 000)	Depositor
R, R_{free}	0.164 , 0.229 0.162 , 0.160	Depositor DCC
R_{free} test set	3428 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	44.5	Xtrriage
Anisotropy	0.396	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16452	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/823	0.66	0/1115
1	B	0.44	0/816	0.77	1/1105 (0.1%)
1	C	0.42	0/834	0.65	0/1130
1	D	0.41	0/816	0.66	0/1105
1	E	0.52	1/823 (0.1%)	0.76	1/1115 (0.1%)
1	F	0.47	0/834	0.70	0/1130
1	G	0.44	0/823	0.67	0/1115
1	H	0.46	0/834	0.71	0/1130
1	I	0.51	1/834 (0.1%)	0.71	0/1130
1	J	0.53	0/832	0.74	1/1127 (0.1%)
1	a	0.42	0/823	0.65	0/1115
1	b	0.44	0/816	0.72	1/1105 (0.1%)
1	c	0.46	0/823	0.71	0/1115
1	d	0.44	0/834	0.68	0/1130
1	e	0.46	0/834	0.72	1/1130 (0.1%)
1	f	0.44	0/823	0.70	0/1115
1	g	0.44	0/823	0.76	1/1115 (0.1%)
1	h	0.48	0/827	0.71	0/1120
1	i	0.48	0/834	0.76	1/1130 (0.1%)
1	j	0.51	0/827	0.80	2/1120 (0.2%)
All	All	0.46	2/16533 (0.0%)	0.71	9/22397 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	J	0	1
1	b	0	1
1	f	0	2

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
All	All	0	4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	69	GLY	C-N	-7.22	1.17	1.34
1	E	32	LYS	CD-CE	-6.57	1.34	1.51

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	j	93	GLU	CA-CB-CG	-7.47	96.97	113.40
1	g	32	LYS	C-N-CA	6.43	137.76	121.70
1	b	32	LYS	C-N-CA	5.95	136.57	121.70
1	e	116	LEU	CA-CB-CG	5.76	128.56	115.30
1	B	31	PHE	C-N-CA	5.64	135.80	121.70
1	E	93	GLU	CA-CB-CG	-5.62	101.03	113.40
1	i	45	ARG	NE-CZ-NH2	-5.27	117.66	120.30
1	J	103	LYS	CD-CE-NZ	5.21	123.67	111.70
1	j	45	ARG	NE-CZ-NH2	-5.11	117.74	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	J	35	ASN	Peptide
1	b	32	LYS	Peptide
1	f	35	ASN	Peptide
1	f	37	GLU	Peptide

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	104/116 (90%)	95 (91%)	7 (7%)	2 (2%)	8	9
1	B	103/116 (89%)	96 (93%)	4 (4%)	3 (3%)	4	4
1	C	105/116 (90%)	100 (95%)	4 (4%)	1 (1%)	15	22
1	D	103/116 (89%)	98 (95%)	4 (4%)	1 (1%)	15	22
1	E	104/116 (90%)	97 (93%)	7 (7%)	0	100	100
1	F	105/116 (90%)	97 (92%)	6 (6%)	2 (2%)	8	9
1	G	104/116 (90%)	97 (93%)	5 (5%)	2 (2%)	8	9
1	H	105/116 (90%)	96 (91%)	5 (5%)	4 (4%)	3	2
1	I	105/116 (90%)	99 (94%)	4 (4%)	2 (2%)	8	9
1	J	105/116 (90%)	98 (93%)	6 (6%)	1 (1%)	15	22
1	a	104/116 (90%)	100 (96%)	4 (4%)	0	100	100
1	b	103/116 (89%)	95 (92%)	5 (5%)	3 (3%)	4	4
1	c	104/116 (90%)	95 (91%)	5 (5%)	4 (4%)	3	2
1	d	105/116 (90%)	96 (91%)	5 (5%)	4 (4%)	3	2
1	e	105/116 (90%)	100 (95%)	5 (5%)	0	100	100
1	f	104/116 (90%)	95 (91%)	7 (7%)	2 (2%)	8	9
1	g	104/116 (90%)	99 (95%)	3 (3%)	2 (2%)	8	9
1	h	104/116 (90%)	98 (94%)	4 (4%)	2 (2%)	8	9
1	i	105/116 (90%)	94 (90%)	10 (10%)	1 (1%)	15	22
1	j	104/116 (90%)	96 (92%)	4 (4%)	4 (4%)	3	2
All	All	2085/2320 (90%)	1941 (93%)	104 (5%)	40 (2%)	8	9

All (40) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	32	LYS
1	B	33	VAL
1	B	35	ASN
1	F	36	ASP
1	H	35	ASN
1	I	36	ASP
1	I	37	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	b	35	ASN
1	c	32	LYS
1	c	35	ASN
1	d	25	ALA
1	d	33	VAL
1	d	34	ASP
1	f	33	VAL
1	g	33	VAL
1	g	35	ASN
1	h	35	ASN
1	h	36	ASP
1	j	14	PRO
1	j	32	LYS
1	A	118	ALA
1	C	35	ASN
1	D	35	ASN
1	H	34	ASP
1	b	33	VAL
1	c	34	ASP
1	d	35	ASN
1	j	35	ASN
1	A	33	VAL
1	H	32	LYS
1	J	36	ASP
1	c	28	ASP
1	i	33	VAL
1	b	36	ASP
1	F	33	VAL
1	G	36	ASP
1	G	118	ALA
1	f	34	ASP
1	j	33	VAL
1	H	33	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	90/99 (91%)	86 (96%)	4 (4%)	28	38
1	B	89/99 (90%)	88 (99%)	1 (1%)	73	83
1	C	91/99 (92%)	88 (97%)	3 (3%)	38	51
1	D	89/99 (90%)	88 (99%)	1 (1%)	73	83
1	E	90/99 (91%)	88 (98%)	2 (2%)	52	66
1	F	91/99 (92%)	89 (98%)	2 (2%)	52	66
1	G	90/99 (91%)	88 (98%)	2 (2%)	52	66
1	H	91/99 (92%)	87 (96%)	4 (4%)	28	38
1	I	91/99 (92%)	86 (94%)	5 (6%)	21	29
1	J	91/99 (92%)	89 (98%)	2 (2%)	52	66
1	a	90/99 (91%)	88 (98%)	2 (2%)	52	66
1	b	89/99 (90%)	86 (97%)	3 (3%)	37	50
1	c	90/99 (91%)	85 (94%)	5 (6%)	21	28
1	d	91/99 (92%)	86 (94%)	5 (6%)	21	29
1	e	91/99 (92%)	88 (97%)	3 (3%)	38	51
1	f	90/99 (91%)	86 (96%)	4 (4%)	28	38
1	g	90/99 (91%)	88 (98%)	2 (2%)	52	66
1	h	90/99 (91%)	84 (93%)	6 (7%)	16	21
1	i	91/99 (92%)	87 (96%)	4 (4%)	28	38
1	j	90/99 (91%)	86 (96%)	4 (4%)	28	38
All	All	1805/1980 (91%)	1741 (96%)	64 (4%)	36	49

All (64) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	LEU
1	A	26	ASP
1	A	35	ASN
1	A	92	PHE
1	B	92	PHE
1	C	16	ASN
1	C	32	LYS
1	C	92	PHE
1	D	92	PHE
1	E	67	TYR
1	E	92	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	65	MET
1	F	92	PHE
1	G	92	PHE
1	G	115	HIS
1	H	15	GLN
1	H	36	ASP
1	H	92	PHE
1	H	98	VAL
1	I	13	ARG
1	I	18	LEU
1	I	48	SER
1	I	65	MET
1	I	92	PHE
1	J	18	LEU
1	J	92	PHE
1	a	37	GLU
1	a	92	PHE
1	b	36	ASP
1	b	54	LYS
1	b	92	PHE
1	c	18	LEU
1	c	36	ASP
1	c	38	ASN
1	c	92	PHE
1	c	115	HIS
1	d	13	ARG
1	d	18	LEU
1	d	27	LYS
1	d	48	SER
1	d	92	PHE
1	e	13	ARG
1	e	32	LYS
1	e	92	PHE
1	f	18	LEU
1	f	92	PHE
1	f	114	GLN
1	f	115	HIS
1	g	34	ASP
1	g	92	PHE
1	h	13	ARG
1	h	18	LEU
1	h	35	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	h	45	ARG
1	h	92	PHE
1	h	102	LEU
1	i	16	ASN
1	i	18	LEU
1	i	38	ASN
1	i	92	PHE
1	j	18	LEU
1	j	30	HIS
1	j	34	ASP
1	j	92	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (64) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	16	ASN
1	A	38	ASN
1	A	84	GLN
1	A	110	HIS
1	A	114	GLN
1	C	30	HIS
1	C	38	ASN
1	C	84	GLN
1	D	15	GLN
1	D	16	ASN
1	D	84	GLN
1	E	30	HIS
1	E	84	GLN
1	F	40	HIS
1	F	58	HIS
1	F	114	GLN
1	G	16	ASN
1	G	41	GLN
1	G	84	GLN
1	G	114	GLN
1	G	115	HIS
1	H	114	GLN
1	I	16	ASN
1	I	114	GLN
1	J	15	GLN
1	J	38	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	J	114	GLN
1	a	16	ASN
1	a	114	GLN
1	b	30	HIS
1	b	84	GLN
1	c	16	ASN
1	c	38	ASN
1	c	58	HIS
1	c	115	HIS
1	d	16	ASN
1	d	30	HIS
1	d	40	HIS
1	d	41	GLN
1	d	58	HIS
1	e	38	ASN
1	e	41	GLN
1	e	84	GLN
1	e	114	GLN
1	f	16	ASN
1	f	41	GLN
1	f	114	GLN
1	f	115	HIS
1	g	16	ASN
1	g	58	HIS
1	g	84	GLN
1	g	114	GLN
1	h	41	GLN
1	h	58	HIS
1	h	84	GLN
1	h	114	GLN
1	i	16	ASN
1	i	38	ASN
1	i	114	GLN
1	j	15	GLN
1	j	40	HIS
1	j	58	HIS
1	j	114	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	I	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	I	69:GLY	C	70:SER	N	1.17

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	106/116 (91%)	0.24	9 (8%) 10 12	39, 53, 109, 142	0
1	B	105/116 (90%)	0.56	14 (13%) 3 4	36, 54, 99, 130	0
1	C	107/116 (92%)	0.65	12 (11%) 5 7	39, 60, 123, 165	0
1	D	105/116 (90%)	0.41	8 (7%) 13 17	39, 59, 100, 137	0
1	E	106/116 (91%)	0.32	10 (9%) 8 10	40, 57, 99, 117	0
1	F	107/116 (92%)	0.19	6 (5%) 24 29	31, 44, 89, 119	0
1	G	106/116 (91%)	0.10	6 (5%) 23 28	33, 43, 86, 121	0
1	H	107/116 (92%)	0.08	3 (2%) 53 60	31, 46, 94, 134	0
1	I	107/116 (92%)	0.14	5 (4%) 31 38	32, 47, 84, 102	0
1	J	107/116 (92%)	0.31	7 (6%) 18 22	32, 46, 91, 129	0
1	a	106/116 (91%)	0.32	7 (6%) 18 21	36, 52, 96, 129	0
1	b	105/116 (90%)	0.32	9 (8%) 10 12	35, 53, 93, 122	0
1	c	106/116 (91%)	0.30	7 (6%) 18 21	35, 52, 123, 141	0
1	d	107/116 (92%)	0.27	8 (7%) 14 17	34, 50, 104, 132	0
1	e	107/116 (92%)	0.05	2 (1%) 66 73	34, 46, 77, 98	0
1	f	106/116 (91%)	0.30	7 (6%) 18 21	29, 46, 92, 122	0
1	g	106/116 (91%)	0.27	8 (7%) 14 17	32, 45, 88, 140	0
1	h	106/116 (91%)	0.12	4 (3%) 40 47	28, 43, 87, 100	0
1	i	107/116 (92%)	0.18	8 (7%) 14 17	30, 44, 89, 117	0
1	j	106/116 (91%)	0.25	7 (6%) 18 21	30, 43, 84, 125	0
All	All	2125/2320 (91%)	0.27	147 (6%) 16 20	28, 50, 99, 165	0

All (147) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	34	ASP	9.3
1	j	35	ASN	9.0
1	C	119	VAL	7.8
1	B	35	ASN	7.1
1	C	35	ASN	6.9
1	g	35	ASN	6.8
1	f	35	ASN	5.6
1	d	36	ASP	5.5
1	a	119	VAL	5.0
1	b	36	ASP	4.9
1	D	37	GLU	4.8
1	j	37	GLU	4.7
1	d	34	ASP	4.7
1	B	33	VAL	4.6
1	c	36	ASP	4.5
1	f	67	TYR	4.4
1	c	119	VAL	4.4
1	I	119	VAL	4.4
1	E	119	VAL	4.3
1	G	37	GLU	4.3
1	c	37	GLU	4.2
1	F	37	GLU	4.2
1	J	120	GLU	4.2
1	C	33	VAL	4.2
1	i	119	VAL	4.1
1	A	35	ASN	4.1
1	j	13	ARG	4.1
1	g	36	ASP	4.1
1	B	37	GLU	4.1
1	a	37	GLU	4.1
1	H	37	GLU	4.1
1	J	33	VAL	3.9
1	B	67	TYR	3.8
1	g	119	VAL	3.8
1	F	34	ASP	3.8
1	A	38	ASN	3.7
1	G	36	ASP	3.7
1	j	36	ASP	3.7
1	A	119	VAL	3.6
1	d	38	ASN	3.6
1	J	34	ASP	3.6
1	E	117	VAL	3.5
1	B	34	ASP	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	c	34	ASP	3.5
1	E	33	VAL	3.4
1	C	37	GLU	3.4
1	J	35	ASN	3.4
1	E	67	TYR	3.3
1	B	118	ALA	3.3
1	D	36	ASP	3.3
1	b	34	ASP	3.3
1	B	32	LYS	3.3
1	c	68	GLU	3.3
1	C	34	ASP	3.3
1	a	36	ASP	3.3
1	f	34	ASP	3.3
1	D	35	ASN	3.2
1	D	118	ALA	3.2
1	c	38	ASN	3.2
1	C	38	ASN	3.2
1	f	119	VAL	3.2
1	C	68	GLU	3.2
1	e	37	GLU	3.2
1	G	35	ASN	3.2
1	F	36	ASP	3.1
1	h	37	GLU	3.1
1	C	67	TYR	3.1
1	e	119	VAL	3.1
1	g	38	ASN	3.1
1	B	38	ASN	3.0
1	g	34	ASP	3.0
1	g	37	GLU	2.9
1	B	26	ASP	2.9
1	J	36	ASP	2.9
1	g	39	GLU	2.9
1	a	35	ASN	2.9
1	f	68	GLU	2.9
1	i	13	ARG	2.9
1	I	67	TYR	2.9
1	b	38	ASN	2.8
1	b	39	GLU	2.8
1	h	118	ALA	2.8
1	b	35	ASN	2.7
1	C	13	ARG	2.7
1	B	39	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	f	37	GLU	2.7
1	A	39	GLU	2.6
1	C	14	PRO	2.6
1	C	36	ASP	2.6
1	h	13	ARG	2.6
1	H	39	GLU	2.6
1	F	13	ARG	2.6
1	H	34	ASP	2.6
1	d	118	ALA	2.6
1	d	37	GLU	2.6
1	F	70	SER	2.6
1	d	67	TYR	2.6
1	A	33	VAL	2.6
1	D	117	VAL	2.5
1	G	34	ASP	2.5
1	b	31	PHE	2.5
1	d	68	GLU	2.5
1	i	37	GLU	2.5
1	G	38	ASN	2.5
1	E	39	GLU	2.5
1	I	36	ASP	2.4
1	j	34	ASP	2.4
1	c	26	ASP	2.4
1	I	13	ARG	2.4
1	i	35	ASN	2.4
1	f	71	PRO	2.4
1	A	34	ASP	2.4
1	A	37	GLU	2.3
1	d	35	ASN	2.3
1	B	36	ASP	2.3
1	b	26	ASP	2.3
1	h	33	VAL	2.3
1	B	14	PRO	2.3
1	A	36	ASP	2.3
1	a	34	ASP	2.2
1	E	95	THR	2.2
1	E	38	ASN	2.2
1	a	38	ASN	2.2
1	i	38	ASN	2.2
1	a	31	PHE	2.2
1	i	39	GLU	2.2
1	F	67	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	33	VAL	2.2
1	J	38	ASN	2.2
1	D	33	VAL	2.1
1	b	33	VAL	2.1
1	D	38	ASN	2.1
1	C	70	SER	2.1
1	i	34	ASP	2.1
1	j	38	ASN	2.1
1	b	67	TYR	2.1
1	E	68	GLU	2.1
1	E	37	GLU	2.1
1	B	31	PHE	2.1
1	E	35	ASN	2.1
1	g	33	VAL	2.1
1	i	26	ASP	2.1
1	J	37	GLU	2.0
1	j	31	PHE	2.0
1	A	118	ALA	2.0
1	I	33	VAL	2.0
1	B	65	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	A	201	1/1	0.90	0.53	86,86,86,86	0
2	CL	f	201	1/1	0.93	0.44	90,90,90,90	0
2	CL	G	201	1/1	0.93	0.47	79,79,79,79	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	b	201	1/1	0.94	0.54	88,88,88,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.