



wwPDB X-ray Structure Validation Summary Report

Sep 20, 2023 – 01:47 PM EDT

PDB ID : 5HYS
Title : Structure of IgE complexed with omalizumab
Authors : Pennington, L.F.; Tarchevskaya, S.S.; Sathiyamoorthy, K.; Jardetzky, T.S.
Deposited on : 2016-02-01
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.














Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	 4% 96%
1	C	222	 3% 93% 7%
1	E	222	 7% 93%
1	H	222	 3% 95% 5%
2	B	218	 97%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	218	
2	F	218	
2	L	218	
3	G	230	
3	I	230	
3	J	230	
3	K	230	
4	M	5	
4	N	5	
4	O	5	
4	P	5	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	D	305	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 20626 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epididymis luminal protein 214.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	222	1667	1054	283	325	5	0	0	0
1	A	222	1667	1054	283	325	5	0	0	0
1	C	222	1667	1054	283	325	5	0	0	0
1	E	218	1639	1038	278	318	5	0	0	0

- Molecule 2 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	217	1675	1045	277	348	5	0	0	0
2	B	217	1675	1045	277	348	5	0	0	0
2	D	217	1675	1045	277	348	5	0	0	0
2	F	215	1662	1038	275	344	5	0	0	0

- Molecule 3 is a protein called Ig epsilon chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	G	214	1693	1058	314	314	7	0	0	0
3	I	213	1685	1054	312	312	7	0	0	0
3	J	214	1693	1058	314	314	7	0	0	0
3	K	213	1685	1054	312	312	7	0	0	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	325	ALA	-	expression tag	UNP P01854
G	326	ASP	-	expression tag	UNP P01854
G	327	PRO	-	expression tag	UNP P01854
G	328	ALA	CYS	engineered mutation	UNP P01854
G	335	CYS	GLY	engineered mutation	UNP P01854
G	548	ALA	-	expression tag	UNP P01854
G	549	ALA	-	expression tag	UNP P01854
G	550	ASP	-	expression tag	UNP P01854
G	551	ASP	-	expression tag	UNP P01854
G	552	ASP	-	expression tag	UNP P01854
G	553	ASP	-	expression tag	UNP P01854
G	554	LYS	-	expression tag	UNP P01854
I	325	ALA	-	expression tag	UNP P01854
I	326	ASP	-	expression tag	UNP P01854
I	327	PRO	-	expression tag	UNP P01854
I	328	ALA	CYS	engineered mutation	UNP P01854
I	335	CYS	GLY	engineered mutation	UNP P01854
I	548	ALA	-	expression tag	UNP P01854
I	549	ALA	-	expression tag	UNP P01854
I	550	ASP	-	expression tag	UNP P01854
I	551	ASP	-	expression tag	UNP P01854
I	552	ASP	-	expression tag	UNP P01854
I	553	ASP	-	expression tag	UNP P01854
I	554	LYS	-	expression tag	UNP P01854
J	325	ALA	-	expression tag	UNP P01854
J	326	ASP	-	expression tag	UNP P01854
J	327	PRO	-	expression tag	UNP P01854
J	328	ALA	CYS	engineered mutation	UNP P01854
J	335	CYS	GLY	engineered mutation	UNP P01854
J	548	ALA	-	expression tag	UNP P01854
J	549	ALA	-	expression tag	UNP P01854
J	550	ASP	-	expression tag	UNP P01854
J	551	ASP	-	expression tag	UNP P01854
J	552	ASP	-	expression tag	UNP P01854
J	553	ASP	-	expression tag	UNP P01854
J	554	LYS	-	expression tag	UNP P01854
K	325	ALA	-	expression tag	UNP P01854
K	326	ASP	-	expression tag	UNP P01854
K	327	PRO	-	expression tag	UNP P01854
K	328	ALA	CYS	engineered mutation	UNP P01854
K	335	CYS	GLY	engineered mutation	UNP P01854
K	548	ALA	-	expression tag	UNP P01854

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
K	549	ALA	-	expression tag	UNP P01854
K	550	ASP	-	expression tag	UNP P01854
K	551	ASP	-	expression tag	UNP P01854
K	552	ASP	-	expression tag	UNP P01854
K	553	ASP	-	expression tag	UNP P01854
K	554	LYS	-	expression tag	UNP P01854

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	M	5	61	34	2	25	0	0	0
4	N	5	61	34	2	25	0	0	0
4	O	5	61	34	2	25	0	0	0
4	P	5	61	34	2	25	0	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	H	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	E	1	Total	O	S	0	0
			5	4	1		
5	E	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	G	1	Total	O	S	0	0
			5	4	1		
5	G	1	Total	O	S	0	0
			5	4	1		
5	I	1	Total	O	S	0	0
			5	4	1		
5	I	1	Total	O	S	0	0
			5	4	1		
5	I	1	Total	O	S	0	0
			5	4	1		
5	I	1	Total	O	S	0	0
			5	4	1		
5	I	1	Total	O	S	0	0
			5	4	1		
5	J	1	Total	O	S	0	0
			5	4	1		
5	J	1	Total	O	S	0	0
			5	4	1		
5	J	1	Total	O	S	0	0
			5	4	1		
5	J	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	K	1	Total 5	O 4	S 1	0	0
5	K	1	Total 5	O 4	S 1	0	0
5	K	1	Total 5	O 4	S 1	0	0
5	K	1	Total 5	O 4	S 1	0	0
5	K	1	Total 5	O 4	S 1	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	6	Total 6	O 6	0	0
6	L	14	Total 14	O 14	0	0
6	A	4	Total 4	O 4	0	0
6	B	10	Total 10	O 10	0	0
6	C	3	Total 3	O 3	0	0
6	D	6	Total 6	O 6	0	0
6	E	3	Total 3	O 3	0	0
6	F	6	Total 6	O 6	0	0
6	G	1	Total 1	O 1	0	0
6	I	2	Total 2	O 2	0	0
6	J	4	Total 4	O 4	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Epididymis luminal protein 214

Chain H: 




- Molecule 1: Epididymis luminal protein 214

Chain A: 




- Molecule 1: Epididymis luminal protein 214

Chain C: 




- Molecule 1: Epididymis luminal protein 214

Chain E: 



- Molecule 2: Uncharacterized protein

Chain L: 




- Molecule 2: Uncharacterized protein





• Molecule 2: Uncharacterized protein





• Molecule 2: Uncharacterized protein



• Molecule 3: Ig epsilon chain C region





• Molecule 3: Ig epsilon chain C region





LYS

• Molecule 3: Ig epsilon chain C region



• Molecule 3: Ig epsilon chain C region





- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M: 20% 60% 20%



- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N: 60% 40%



- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O: 60% 40%



- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain P: 60% 40%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	100.10Å 107.14Å 151.04Å 90.00° 95.18° 90.00°	Depositor
Resolution (Å)	37.61 – 2.50 37.61 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.8 (37.61-2.50) 97.8 (37.61-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.72 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.221 , 0.239 0.222 , 0.240	Depositor DCC
R_{free} test set	1512 reflections (1.41%)	wwPDB-VP
Wilson B-factor (Å ²)	61.9	Xtrriage
Anisotropy	0.243	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20626	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, SO4, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/1712	0.42	0/2336
1	C	0.31	0/1712	0.47	0/2336
1	E	0.26	0/1683	0.43	0/2296
1	H	0.36	0/1712	0.48	0/2336
2	B	0.24	0/1712	0.43	0/2325
2	D	0.23	0/1712	0.41	0/2325
2	F	0.23	0/1699	0.41	0/2308
2	L	0.30	0/1712	0.43	0/2325
3	G	0.23	0/1737	0.43	0/2366
3	I	0.23	0/1729	0.45	0/2354
3	J	0.25	0/1737	0.45	0/2366
3	K	0.24	0/1729	0.44	0/2354
All	All	0.27	0/20586	0.44	0/28027

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1667	0	1618	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1667	0	1620	9	0
1	E	1639	0	1589	5	0
1	H	1667	0	1620	6	0
2	B	1675	0	1600	4	0
2	D	1675	0	1600	4	0
2	F	1662	0	1590	2	0
2	L	1675	0	1600	2	0
3	G	1693	0	1673	6	0
3	I	1685	0	1668	8	0
3	J	1693	0	1673	4	0
3	K	1685	0	1669	8	0
4	M	61	0	52	2	0
4	N	61	0	52	0	0
4	O	61	0	52	0	0
4	P	61	0	52	0	0
5	A	15	0	0	0	0
5	B	20	0	0	1	0
5	C	20	0	0	0	0
5	D	30	0	0	3	0
5	E	10	0	0	0	0
5	F	25	0	0	1	0
5	G	10	0	0	0	0
5	H	20	0	0	0	0
5	I	25	0	0	0	0
5	J	20	0	0	0	0
5	K	25	0	0	1	0
5	L	20	0	0	0	0
6	A	4	0	0	0	0
6	B	10	0	0	2	0
6	C	3	0	0	1	0
6	D	6	0	0	0	0
6	E	3	0	0	0	0
6	F	6	0	0	0	0
6	G	1	0	0	0	0
6	H	6	0	0	0	0
6	I	2	0	0	2	0
6	J	4	0	0	0	0
6	L	14	0	0	1	0
All	All	20626	0	19728	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 61 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:M:5:MAN:HO2	4:M:5:MAN:HO6	1.11	0.97
1:C:122:ALA:O	6:C:401:HOH:O	2.08	0.72
2:B:116:ALA:O	6:B:401:HOH:O	2.08	0.71
5:D:305:SO4:O2	3:G:408:ARG:NH2	2.24	0.70
2:L:98:ASP:OD1	6:L:401:HOH:O	2.10	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/222 (99%)	217 (99%)	3 (1%)	0	100	100
1	C	220/222 (99%)	216 (98%)	3 (1%)	1 (0%)	29	48
1	E	214/222 (96%)	209 (98%)	5 (2%)	0	100	100
1	H	220/222 (99%)	217 (99%)	3 (1%)	0	100	100
2	B	215/218 (99%)	209 (97%)	6 (3%)	0	100	100
2	D	215/218 (99%)	209 (97%)	6 (3%)	0	100	100
2	F	213/218 (98%)	207 (97%)	6 (3%)	0	100	100
2	L	215/218 (99%)	210 (98%)	5 (2%)	0	100	100
3	G	212/230 (92%)	205 (97%)	6 (3%)	1 (0%)	29	48
3	I	211/230 (92%)	205 (97%)	6 (3%)	0	100	100
3	J	212/230 (92%)	205 (97%)	6 (3%)	1 (0%)	29	48
3	K	211/230 (92%)	204 (97%)	7 (3%)	0	100	100
All	All	2578/2680 (96%)	2513 (98%)	62 (2%)	3 (0%)	51	73

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	G	456	SER
3	J	454	PRO
1	C	137	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/186 (100%)	184 (99%)	2 (1%)	73	89
1	C	186/186 (100%)	185 (100%)	1 (0%)	88	96
1	E	182/186 (98%)	179 (98%)	3 (2%)	62	84
1	H	186/186 (100%)	183 (98%)	3 (2%)	62	84
2	B	191/192 (100%)	189 (99%)	2 (1%)	76	90
2	D	191/192 (100%)	189 (99%)	2 (1%)	76	90
2	F	190/192 (99%)	189 (100%)	1 (0%)	88	96
2	L	191/192 (100%)	189 (99%)	2 (1%)	76	90
3	G	190/200 (95%)	190 (100%)	0	100	100
3	I	189/200 (94%)	188 (100%)	1 (0%)	88	96
3	J	190/200 (95%)	190 (100%)	0	100	100
3	K	189/200 (94%)	188 (100%)	1 (0%)	88	96
All	All	2261/2312 (98%)	2243 (99%)	18 (1%)	81	93

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	207	ASN
3	K	475	SER
3	I	430	MET
2	B	187	LYS
1	E	186	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	I	424	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	M	1	4,3	14,14,15	0.35	0	17,19,21	0.43	0
4	NAG	M	2	4	14,14,15	0.24	0	17,19,21	0.47	0
4	BMA	M	3	4	11,11,12	0.91	1 (9%)	15,15,17	1.26	1 (6%)
4	MAN	M	4	4	11,11,12	0.84	1 (9%)	15,15,17	0.86	1 (6%)
4	MAN	M	5	4	11,11,12	0.95	1 (9%)	15,15,17	1.68	2 (13%)
4	NAG	N	1	4,3	14,14,15	0.17	0	17,19,21	0.50	0
4	NAG	N	2	4	14,14,15	0.39	0	17,19,21	0.40	0
4	BMA	N	3	4	11,11,12	0.76	0	15,15,17	0.79	0
4	MAN	N	4	4	11,11,12	0.60	0	15,15,17	1.19	2 (13%)
4	MAN	N	5	4	11,11,12	0.95	0	15,15,17	2.05	4 (26%)
4	NAG	O	1	4,3	14,14,15	0.26	0	17,19,21	0.48	0
4	NAG	O	2	4	14,14,15	0.28	0	17,19,21	0.42	0
4	BMA	O	3	4	11,11,12	0.72	0	15,15,17	0.78	0
4	MAN	O	4	4	11,11,12	0.82	1 (9%)	15,15,17	0.89	1 (6%)
4	MAN	O	5	4	11,11,12	0.79	1 (9%)	15,15,17	0.94	1 (6%)
4	NAG	P	1	4,3	14,14,15	0.31	0	17,19,21	0.49	0
4	NAG	P	2	4	14,14,15	0.29	0	17,19,21	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BMA	P	3	4	11,11,12	0.73	0	15,15,17	0.93	0
4	MAN	P	4	4	11,11,12	0.81	1 (9%)	15,15,17	0.89	1 (6%)
4	MAN	P	5	4	11,11,12	0.89	1 (9%)	15,15,17	1.70	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	M	1	4,3	-	0/6/23/26	0/1/1/1
4	NAG	M	2	4	-	2/6/23/26	0/1/1/1
4	BMA	M	3	4	-	2/2/19/22	0/1/1/1
4	MAN	M	4	4	-	2/2/19/22	0/1/1/1
4	MAN	M	5	4	-	0/2/19/22	1/1/1/1
4	NAG	N	1	4,3	-	0/6/23/26	0/1/1/1
4	NAG	N	2	4	-	2/6/23/26	0/1/1/1
4	BMA	N	3	4	-	2/2/19/22	0/1/1/1
4	MAN	N	4	4	-	2/2/19/22	0/1/1/1
4	MAN	N	5	4	-	0/2/19/22	0/1/1/1
4	NAG	O	1	4,3	-	2/6/23/26	0/1/1/1
4	NAG	O	2	4	-	1/6/23/26	0/1/1/1
4	BMA	O	3	4	-	0/2/19/22	0/1/1/1
4	MAN	O	4	4	-	1/2/19/22	0/1/1/1
4	MAN	O	5	4	-	2/2/19/22	0/1/1/1
4	NAG	P	1	4,3	-	2/6/23/26	0/1/1/1
4	NAG	P	2	4	-	2/6/23/26	0/1/1/1
4	BMA	P	3	4	-	2/2/19/22	0/1/1/1
4	MAN	P	4	4	-	2/2/19/22	0/1/1/1
4	MAN	P	5	4	-	0/2/19/22	1/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	M	4	MAN	O5-C1	-2.37	1.39	1.43
4	O	4	MAN	O5-C1	-2.35	1.40	1.43
4	M	5	MAN	O5-C5	2.31	1.48	1.43
4	M	3	BMA	O5-C1	-2.30	1.40	1.43
4	P	4	MAN	O5-C1	-2.27	1.40	1.43

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	P	5	MAN	C1-O5-C5	5.44	119.57	112.19
4	M	5	MAN	C1-O5-C5	5.27	119.33	112.19
4	N	5	MAN	C1-O5-C5	4.94	118.88	112.19
4	N	5	MAN	O5-C1-C2	3.68	116.44	110.77
4	N	5	MAN	C1-C2-C3	3.29	113.72	109.67

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	N	2	NAG	O5-C5-C6-O6
4	P	2	NAG	O5-C5-C6-O6
4	M	3	BMA	O5-C5-C6-O6
4	P	4	MAN	O5-C5-C6-O6
4	N	2	NAG	C4-C5-C6-O6


All (2) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	M	5	MAN	C1-C2-C3-C4-C5-O5
4	P	5	MAN	C1-C2-C3-C4-C5-O5


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	M	5	MAN	1	0
4	M	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

48 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	SO4	F	304	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	B	304	-	4,4,4	0.13	0	6,6,6	0.07	0
5	SO4	I	606	-	4,4,4	0.14	0	6,6,6	0.07	0
5	SO4	D	302	-	4,4,4	0.15	0	6,6,6	0.04	0
5	SO4	A	301	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	C	301	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	G	606	-	4,4,4	0.14	0	6,6,6	0.05	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	A	302	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	D	301	-	4,4,4	0.13	0	6,6,6	0.07	0
5	SO4	F	305	2,3	4,4,4	0.28	0	6,6,6	0.35	0
5	SO4	D	304	-	4,4,4	0.14	0	6,6,6	0.04	0
5	SO4	D	306	-	4,4,4	0.14	0	6,6,6	0.07	0
5	SO4	K	607	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	H	304	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	J	607	-	4,4,4	0.11	0	6,6,6	0.26	0
5	SO4	E	301	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	A	303	1	4,4,4	0.17	0	6,6,6	0.49	0
5	SO4	I	609	-	4,4,4	0.15	0	6,6,6	0.05	0
5	SO4	J	606	-	4,4,4	0.14	0	6,6,6	0.11	0
5	SO4	F	301	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	K	608	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	L	304	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	G	607	-	4,4,4	0.12	0	6,6,6	0.06	0
5	SO4	K	609	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	H	302	-	4,4,4	0.13	0	6,6,6	0.08	0
5	SO4	I	607	-	4,4,4	0.14	0	6,6,6	0.07	0
5	SO4	B	301	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	J	609	-	4,4,4	0.17	0	6,6,6	0.22	0
5	SO4	C	303	-	4,4,4	0.14	0	6,6,6	0.04	0
5	SO4	F	303	-	4,4,4	0.14	0	6,6,6	0.08	0
5	SO4	B	303	-	4,4,4	0.13	0	6,6,6	0.08	0
5	SO4	B	302	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	C	302	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	C	304	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	I	610	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	I	608	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	K	606	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	K	610	-	4,4,4	0.33	0	6,6,6	0.30	0
5	SO4	L	301	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	D	303	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	E	302	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	F	302	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	J	608	-	4,4,4	0.14	0	6,6,6	0.06	0
5	SO4	D	305	-	4,4,4	0.13	0	6,6,6	0.10	0
5	SO4	H	301	-	4,4,4	0.13	0	6,6,6	0.05	0
5	SO4	L	302	-	4,4,4	0.14	0	6,6,6	0.07	0
5	SO4	L	303	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	H	303	-	4,4,4	0.15	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	304	SO4	1	0
5	F	303	SO4	1	0
5	B	303	SO4	1	0
5	K	610	SO4	1	0
5	D	305	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	222/222 (100%)	0.40	9 (4%) 37 40	49, 62, 83, 116	0
1	C	222/222 (100%)	0.35	7 (3%) 47 51	53, 70, 85, 106	0
1	E	218/222 (98%)	0.42	15 (6%) 16 17	48, 70, 104, 128	0
1	H	222/222 (100%)	0.17	7 (3%) 47 51	46, 58, 85, 96	0
2	B	217/218 (99%)	0.09	0 100 100	43, 57, 79, 100	0
2	D	217/218 (99%)	0.25	3 (1%) 75 77	48, 65, 94, 105	0
2	F	215/218 (98%)	0.37	8 (3%) 41 45	46, 63, 103, 110	0
2	L	217/218 (99%)	0.02	4 (1%) 68 71	43, 55, 79, 110	0
3	G	214/230 (93%)	0.51	13 (6%) 21 22	56, 71, 108, 127	0
3	I	213/230 (92%)	0.62	18 (8%) 10 10	56, 76, 112, 120	0
3	J	214/230 (93%)	0.65	18 (8%) 11 11	47, 75, 110, 154	0
3	K	213/230 (92%)	0.85	31 (14%) 2 2	62, 81, 123, 137	0
All	All	2604/2680 (97%)	0.39	133 (5%) 28 29	43, 68, 102, 154	0

The worst 5 of 133 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	141	GLY	9.4
3	J	455	GLY	9.0
3	K	365	PRO	6.9
1	C	1	GLU	5.9
3	I	363	LEU	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MAN	M	4	11/12	0.72	0.25	112,127,134,134	0
4	MAN	O	4	11/12	0.76	0.24	126,138,143,148	0
4	MAN	N	4	11/12	0.77	0.21	123,130,141,142	0
4	MAN	P	4	11/12	0.83	0.19	124,135,139,145	0
4	MAN	N	5	11/12	0.84	0.18	96,103,105,107	0
4	MAN	O	5	11/12	0.85	0.22	120,126,130,142	0
4	BMA	N	3	11/12	0.86	0.15	93,99,107,114	0
4	MAN	P	5	11/12	0.87	0.14	93,99,104,108	0
4	BMA	P	3	11/12	0.90	0.16	97,100,110,116	0
4	NAG	O	1	14/15	0.91	0.22	85,88,94,100	0
4	NAG	O	2	14/15	0.91	0.16	83,86,93,96	0
4	NAG	P	1	14/15	0.91	0.19	96,103,105,106	0
4	NAG	N	1	14/15	0.92	0.16	92,95,99,100	0
4	MAN	M	5	11/12	0.92	0.10	89,90,96,97	0
4	BMA	O	3	11/12	0.93	0.17	93,102,116,117	0
4	BMA	M	3	11/12	0.94	0.10	84,88,97,104	0
4	NAG	M	2	14/15	0.94	0.20	82,84,88,91	0
4	NAG	N	2	14/15	0.94	0.16	86,92,94,95	0
4	NAG	P	2	14/15	0.95	0.17	90,96,99,99	0
4	NAG	M	1	14/15	0.96	0.14	86,90,94,97	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain M:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around Chain N:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	F	305	5/5	0.41	0.40	82,83,93,103	5
5	SO4	D	305	5/5	0.74	0.22	95,102,104,111	0
5	SO4	A	303	5/5	0.75	0.26	62,69,71,72	5
5	SO4	D	306	5/5	0.81	0.28	60,60,64,65	5
5	SO4	G	607	5/5	0.81	0.23	62,63,66,67	5
5	SO4	I	608	5/5	0.81	0.22	98,98,113,113	0
5	SO4	B	304	5/5	0.85	0.16	97,99,106,112	0
5	SO4	L	304	5/5	0.85	0.24	113,113,122,123	0
5	SO4	J	608	5/5	0.85	0.16	91,98,102,103	0
5	SO4	I	610	5/5	0.86	0.24	92,94,105,114	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	SO4	K	610	5/5	0.86	0.24	59,61,68,89	5
5	SO4	C	303	5/5	0.87	0.19	95,101,102,102	0
5	SO4	J	607	5/5	0.88	0.15	98,98,101,105	0
5	SO4	C	304	5/5	0.89	0.19	108,115,118,118	0
5	SO4	J	609	5/5	0.89	0.15	91,92,101,103	0
5	SO4	A	302	5/5	0.89	0.16	87,89,96,111	0
5	SO4	F	303	5/5	0.90	0.16	101,106,111,115	0
5	SO4	L	303	5/5	0.90	0.14	115,117,124,133	0
5	SO4	K	609	5/5	0.90	0.20	94,100,105,109	0
5	SO4	D	303	5/5	0.90	0.15	95,101,102,104	0
5	SO4	E	301	5/5	0.91	0.14	74,79,87,88	0
5	SO4	F	301	5/5	0.91	0.12	83,84,89,94	0
5	SO4	C	301	5/5	0.91	0.16	93,97,103,106	0
5	SO4	D	304	5/5	0.92	0.20	96,98,106,113	0
5	SO4	D	302	5/5	0.92	0.14	72,76,94,96	0
5	SO4	K	608	5/5	0.92	0.15	93,93,98,101	0
5	SO4	F	304	5/5	0.92	0.25	89,91,106,109	0
5	SO4	E	302	5/5	0.92	0.17	70,73,79,81	5
5	SO4	B	303	5/5	0.93	0.24	100,104,106,116	0
5	SO4	L	302	5/5	0.93	0.18	83,83,93,109	0
5	SO4	K	607	5/5	0.93	0.15	79,87,95,98	0
5	SO4	A	301	5/5	0.94	0.18	96,96,99,101	0
5	SO4	F	302	5/5	0.94	0.17	100,107,111,121	0
5	SO4	I	606	5/5	0.94	0.11	84,90,95,99	0
5	SO4	B	302	5/5	0.95	0.10	90,96,103,104	0
5	SO4	H	302	5/5	0.95	0.12	74,74,81,82	0
5	SO4	D	301	5/5	0.95	0.09	85,91,92,92	0
5	SO4	H	304	5/5	0.95	0.15	97,98,105,106	0
5	SO4	L	301	5/5	0.95	0.11	71,75,77,78	0
5	SO4	I	607	5/5	0.95	0.17	79,80,83,85	0
5	SO4	C	302	5/5	0.95	0.18	86,89,94,102	0
5	SO4	I	609	5/5	0.95	0.10	83,92,96,97	0
5	SO4	J	606	5/5	0.96	0.11	67,81,82,89	0
5	SO4	H	301	5/5	0.96	0.09	77,77,82,87	0
5	SO4	G	606	5/5	0.97	0.11	76,80,82,84	0
5	SO4	K	606	5/5	0.97	0.12	76,77,83,85	0
5	SO4	H	303	5/5	0.98	0.10	62,72,78,83	0
5	SO4	B	301	5/5	0.98	0.10	63,73,79,84	0

6.5 Other polymers [i](#)

There are no such residues in this entry.