



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 11, 2023 – 08:06 am GMT

PDB ID : 5JLZ  
Title : Crystal structure of HLA-DRB1\*04:01 in complex with modified alpha-enolase peptide 26-40 with citrulline at the position 32  
Authors : Dubnovitsky, A.; Kozhukh, G.; Sandalova, T.; Achour, A.  
Deposited on : 2016-04-28  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.


---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36



*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	15	 87% 7% 7%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	180	1482	960	241	276	5	0	0	0
1	C	179	1487	965	243	273	6	0	2	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	SER	-	expression tag	UNP P01903
A	183	SER	-	expression tag	UNP P01903
A	184	ALA	-	expression tag	UNP P01903
A	185	ASP	-	expression tag	UNP P01903
A	186	LEU	-	expression tag	UNP P01903
A	187	VAL	-	expression tag	UNP P01903
A	188	PRO	-	expression tag	UNP P01903
A	189	ARG	-	expression tag	UNP P01903
C	182	SER	-	expression tag	UNP P01903
C	183	SER	-	expression tag	UNP P01903
C	184	ALA	-	expression tag	UNP P01903
C	185	ASP	-	expression tag	UNP P01903
C	186	LEU	-	expression tag	UNP P01903
C	187	VAL	-	expression tag	UNP P01903
C	188	PRO	-	expression tag	UNP P01903
C	189	ARG	-	expression tag	UNP P01903

- Molecule 2 is a protein called HLA class II histocompatibility antigen, DRB1-4 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	193	1590	1003	282	300	5	0	0	0
2	D	190	1575	995	278	297	5	0	1	0


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	191	SER	-	expression tag	UNP P13760
B	192	SER	-	expression tag	UNP P13760
B	193	ALA	-	expression tag	UNP P13760
B	194	ASP	-	expression tag	UNP P13760
B	195	LEU	-	expression tag	UNP P13760
B	196	VAL	-	expression tag	UNP P13760
B	197	PRO	-	expression tag	UNP P13760
B	198	ARG	-	expression tag	UNP P13760
D	191	SER	-	expression tag	UNP P13760
D	192	SER	-	expression tag	UNP P13760
D	193	ALA	-	expression tag	UNP P13760
D	194	ASP	-	expression tag	UNP P13760
D	195	LEU	-	expression tag	UNP P13760
D	196	VAL	-	expression tag	UNP P13760
D	197	PRO	-	expression tag	UNP P13760
D	198	ARG	-	expression tag	UNP P13760

- Molecule 3 is a protein called Alpha-enolase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	14	Total	C	N	O	0	0	0
			95	59	17	19			
3	F	14	Total	C	N	O	0	1	0
			103	64	19	20			

- Molecule 4 is MALONIC ACID (three-letter code: MLA) (formula: C<sub>3</sub>H<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			7	3	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	126	Total	O	0	0
			126	126		
5	B	61	Total	O	0	0
			61	61		
5	C	57	Total	O	0	0
			57	57		
5	D	109	Total	O	0	0
			109	109		
5	E	9	Total	O	0	0
			9	9		
5	F	4	Total	O	0	0
			4	4		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain A: 





- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain C: 





- Molecule 2: HLA class II histocompatibility antigen, DRB1-4 beta chain

Chain B: 




- Molecule 2: HLA class II histocompatibility antigen, DRB1-4 beta chain

Chain D: 




- Molecule 3: Alpha-enolase

Chain E: 



- Molecule 3: Alpha-enolase

Chain F:  87% 7% 7%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.44Å 73.78Å 145.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.76 – 1.99 47.76 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.1 (47.76-1.99) 99.1 (47.76-1.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.8.0131	Depositor
R, $R_{free}$	0.206 , 0.243 0.211 , 0.249	Depositor DCC
$R_{free}$ test set	2100 reflections (4.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.7	Xtrriage
Anisotropy	0.661	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 36.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6705	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CIR, MLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/1527	0.85	4/2081 (0.2%)
1	C	0.68	0/1538	0.80	0/2094
2	B	0.73	0/1634	0.85	2/2221 (0.1%)
2	D	0.79	0/1622	0.89	3/2207 (0.1%)
3	E	0.78	0/84	0.73	0/109
3	F	0.78	0/84	0.71	0/109
All	All	0.75	0/6489	0.85	9/8821 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	ARG	NE-CZ-NH1	7.27	123.93	120.30
2	D	41	ASP	CB-CG-OD1	6.67	124.30	118.30
2	B	41	ASP	CB-CG-OD1	6.48	124.14	118.30
1	A	146	ARG	NE-CZ-NH2	-5.75	117.42	120.30
2	D	158	LEU	CA-CB-CG	5.71	128.43	115.30
2	B	94	ARG	NE-CZ-NH1	5.31	122.96	120.30
1	A	140	ARG	NE-CZ-NH2	-5.20	117.70	120.30
2	D	130	ARG	NE-CZ-NH1	5.17	122.88	120.30
1	A	50	ARG	NE-CZ-NH1	5.14	122.87	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	130	THR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1482	0	1420	4	0
1	C	1487	0	1438	13	0
2	B	1590	0	1499	6	0
2	D	1575	0	1479	8	0
3	E	95	0	91	0	0
3	F	103	0	98	0	0
4	D	7	0	2	0	0
5	A	126	0	0	0	0
5	B	61	0	0	1	0
5	C	57	0	0	0	0
5	D	109	0	0	1	0
5	E	9	0	0	0	0
5	F	4	0	0	0	0
All	All	6705	0	6027	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:13:HIS:HD2	2:D:28:ASP:OD1	1.68	0.76
1:C:160:VAL:HG23	1:C:179:GLU:HB3	1.70	0.71
2:B:55:ARG:NH1	2:B:59:GLU:OE2	2.23	0.71
1:C:130:THR:HG23	1:C:132:VAL:HG22	1.81	0.63
2:B:19:ASN:HB2	2:B:22:GLU:HG2	1.79	0.62
1:C:130:THR:HA	1:C:132:VAL:HG22	1.80	0.62
1:C:130:THR:HG23	1:C:132:VAL:CG2	2.28	0.62

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:13:HIS:CD2	2:D:28:ASP:OD1	2.56	0.57
1:A:3:GLU:HB3	2:B:18:PHE:CE2	2.41	0.56
2:B:52:GLU:OE1	2:D:52:GLU:OE1	2.24	0.56
1:C:160:VAL:CG2	1:C:177:HIS:CE1	2.91	0.54
1:C:160:VAL:HG21	1:C:177:HIS:CE1	2.43	0.53
1:C:11:GLU:OE2	2:D:13:HIS:HE1	1.92	0.53
1:C:36[A]:MET:CE	1:C:63:ILE:CD1	2.88	0.52
1:C:128:VAL:HG12	1:C:129:THR:O	2.09	0.52
2:D:33:HIS:HB2	5:D:313:HOH:O	2.11	0.50
1:C:35:ASP:OD2	1:C:38:LYS:N	2.43	0.50
1:A:46:GLU:OE1	1:A:50:ARG:NH2	2.47	0.48
2:B:134:ASN:OD1	2:B:170:VAL:HG12	2.13	0.48
2:D:96:TYR:HE1	2:D:179:SER:O	1.97	0.47
1:C:34:VAL:O	1:C:36[A]:MET:HE2	2.16	0.46
1:A:162:ASP:OD2	2:D:162:GLU:OE2	2.33	0.46
2:B:33:HIS:HB2	5:B:222:HOH:O	2.17	0.44
1:A:39:LYS:CD	1:A:60:LEU:HD21	2.47	0.44
1:C:39:LYS:CD	1:C:60:LEU:HD21	2.49	0.43
2:D:18:PHE:CD1	2:D:23:ARG:HG3	2.54	0.42
1:C:129:THR:OG1	1:C:130:THR:N	2.53	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/189 (94%)	175 (98%)	3 (2%)	0	100	100
1	C	179/189 (95%)	174 (97%)	5 (3%)	0	100	100
2	B	191/198 (96%)	182 (95%)	9 (5%)	0	100	100
2	D	189/198 (96%)	186 (98%)	3 (2%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	11/15 (73%)	11 (100%)	0	0	100	100
3	F	11/15 (73%)	11 (100%)	0	0	100	100
All	All	759/804 (94%)	739 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/173 (95%)	161 (98%)	4 (2%)	49	51
1	C	166/173 (96%)	159 (96%)	7 (4%)	30	27
2	B	173/178 (97%)	170 (98%)	3 (2%)	60	65
2	D	172/178 (97%)	167 (97%)	5 (3%)	42	43
3	E	8/9 (89%)	8 (100%)	0	100	100
3	F	8/9 (89%)	8 (100%)	0	100	100
All	All	692/720 (96%)	673 (97%)	19 (3%)	43	46

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	SER
1	A	92	LEU
1	A	100	ARG
1	A	181	ASP
2	B	23	ARG
2	B	185	THR
2	B	192	SER
1	C	19	SER
1	C	47	GLU
1	C	75	LYS
1	C	92	LEU
1	C	118	ASN

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	C	172	GLU
1	C	176	LYS
2	D	35	GLU
2	D	128	GLU
2	D	167	SER
2	D	182	SER
2	D	185	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	ASN
1	A	149	HIS
1	C	15	ASN
1	C	84	ASN
2	D	13	HIS
2	D	16	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	CIR	F	32[A]	-	9,10,11	0.61	0	6,11,13	1.50	1 (16%)
3	CIR	E	32	3	9,10,11	0.77	0	6,11,13	3.45	4 (66%)
3	CIR	F	32[B]	-	9,10,11	0.67	0	6,11,13	3.32	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CIR	F	32[A]	-	-	4/8/9/11	-
3	CIR	E	32	3	-	2/8/9/11	-
3	CIR	F	32[B]	-	-	5/8/9/11	-

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	32[B]	CIR	N8-C7-N6	5.76	122.93	116.85
3	E	32	CIR	C5-N6-C7	5.38	129.13	122.73
3	F	32[B]	CIR	O7-C7-N6	-4.95	118.07	121.74
3	E	32	CIR	N8-C7-N6	4.80	121.91	116.85
3	E	32	CIR	O7-C7-N6	-3.14	119.41	121.74
3	F	32[A]	CIR	C5-N6-C7	2.70	125.94	122.73
3	E	32	CIR	O7-C7-N8	-2.68	118.61	123.22
3	F	32[B]	CIR	O7-C7-N8	-2.46	119.00	123.22

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	32[B]	CIR	C4-C3-CA-C
3	F	32[B]	CIR	C4-C3-CA-N
3	F	32[A]	CIR	C3-C4-C5-N6
3	E	32	CIR	O7-C7-N6-C5
3	E	32	CIR	N8-C7-N6-C5
3	F	32[A]	CIR	O7-C7-N6-C5
3	F	32[A]	CIR	N8-C7-N6-C5
3	F	32[B]	CIR	O7-C7-N6-C5
3	F	32[B]	CIR	N8-C7-N6-C5
3	F	32[B]	CIR	C3-C4-C5-N6
3	F	32[A]	CIR	C4-C5-N6-C7

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	MLA	D	201	-	6,6,6	1.16	0	7,7,7	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MLA	D	201	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	201	MLA	C1-C2-C3-O3A
4	D	201	MLA	C1-C2-C3-O3B

There are no ring outliers.

No monomer is involved in short contacts.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	180/189 (95%)	0.12	1 (0%) 89 88	22, 36, 62, 80	0
1	C	179/189 (94%)	0.66	15 (8%) 11 10	25, 47, 78, 98	0
2	B	193/198 (97%)	0.82	30 (15%) 2 1	27, 54, 106, 116	0
2	D	190/198 (95%)	0.10	5 (2%) 56 54	22, 34, 60, 86	0
3	E	13/15 (86%)	0.35	0 100 100	33, 40, 54, 55	0
3	F	13/15 (86%)	-0.06	0 100 100	32, 37, 51, 63	0
All	All	768/804 (95%)	0.42	51 (6%) 18 17	22, 41, 86, 116	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	131	GLY	6.5
2	B	110	GLN	5.6
2	B	132	PHE	5.4
2	B	137	GLU	5.4
1	C	130	THR	5.3
2	B	111	HIS	4.6
2	B	131	TRP	4.6
2	B	167	SER	4.5
2	D	108	PRO	4.2
2	D	106	THR	3.7
2	B	108	PRO	3.6
2	B	166	ARG	3.6
2	B	139	LYS	3.6
1	C	154	LEU	3.5
2	B	187	GLU	3.5
1	C	153	PHE	3.3
2	B	1	GLY	3.3
1	A	50	ARG	3.3
2	B	96	TYR	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	100	ARG	3.1
1	C	160	VAL	3.1
1	C	79	TYR	3.1
2	D	2	ASP	3.1
2	B	164	VAL	3.1
2	B	184	LEU	3.0
2	D	191	SER	3.0
1	C	132	VAL	2.9
1	C	37	ALA	2.9
2	B	134	ASN	2.9
2	B	162	GLU	2.8
2	B	163	THR	2.8
2	B	109	LEU	2.7
2	B	112	HIS	2.7
2	B	138	GLU	2.6
2	B	165	PRO	2.5
2	D	107	GLN	2.5
2	B	193	ALA	2.5
1	C	36[A]	MET	2.5
2	B	141	GLY	2.4
1	C	129	THR	2.4
1	C	60	LEU	2.3
2	B	189	ARG	2.3
2	B	173	CYS	2.3
2	B	169	GLU	2.2
2	B	136	GLN	2.2
2	B	171	TYR	2.2
1	C	38	LYS	2.2
1	C	180	PHE	2.2
2	B	170	VAL	2.0
1	C	124	ASN	2.0
2	B	21	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CIR	E	32	11/12	0.86	0.17	35,38,56,61	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CIR	F	32[A]	11/12	0.94	0.21	32,35,40,40	8
3	CIR	F	32[B]	11/12	0.94	0.21	32,35,38,40	8

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MLA	D	201	7/7	0.90	0.12	53,54,56,58	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.