



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 02:53 am BST

PDB ID : 5MJV  
Title : Rebuild and re-refined model for Human Parechovirus 1  
Authors : Shakeel, S.; Dykeman, E.C.; White, S.J.; Ora, A.; Cockburn, J.J.B.; Butcher, S.J.; Stockley, P.G.; Twarock, R.  
Deposited on : 2016-12-01  
Resolution : 3.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RNA backbone	3102	1116 (3.40-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	234	 34% 46% 18%
2	B	253	 47% 46% 6%
3	C	289	 47% 41% 11%
4	D	6	 50% 17% 33%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5486 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CAPSID SUBUNIT VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	192	1507	971	250	279	7	0	0	0

- Molecule 2 is a protein called CAPSID SUBUNIT VP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	239	1864	1185	320	349	10	0	0	0

- Molecule 3 is a protein called CAPSID SUBUNIT VP0.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	257	1995	1265	332	392	6	0	0	0


- Molecule 4 is a RNA chain called RNA (5'-R(\*GP\*UP\*UP\*UP\*UP\*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	6	120	55	15	45	5	0	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CAPSID SUBUNIT VP1




- Molecule 2: CAPSID SUBUNIT VP3



- Molecule 3: CAPSID SUBUNIT VP0





- Molecule 4: RNA (5'-R(\*GP\*UP\*UP\*UP\*UP\*U)-3')

Chain D:  50% 17% 33%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	399.50Å 399.50Å 332.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	345.98 – 3.09 63.93 – 3.09	Depositor EDS
% Data completeness (in resolution range)	82.7 (345.98-3.09) 82.8 (63.93-3.09)	Depositor EDS
$R_{merge}$	0.35	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.32 (at 3.07Å)	Xtrriage
Refinement program	CNS 1.3, REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.258 , 0.261 0.517 , 0.517	Depositor DCC
$R_{free}$ test set	11563 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.8	Xtrriage
Anisotropy	0.012	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.20	EDS
Total number of atoms	5486	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.45	0/1551	0.72	5/2113 (0.2%)
2	B	0.53	2/1909 (0.1%)	0.64	2/2592 (0.1%)
3	C	0.37	0/2048	0.61	0/2810
4	D	1.85	3/132 (2.3%)	1.28	2/203 (1.0%)
All	All	0.53	5/5640 (0.1%)	0.68	9/7718 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	5	U	O3'-P	-9.23	1.50	1.61
4	D	2	U	O3'-P	-8.16	1.51	1.61
4	D	3	U	O3'-P	-7.73	1.51	1.61
2	B	99	TYR	CE1-CZ	-6.88	1.29	1.38
2	B	99	TYR	CE2-CZ	-5.88	1.30	1.38

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	D	5	U	C2'-C3'-O3'	-7.70	92.57	109.50
1	A	203	CYS	N-CA-C	-6.69	92.94	111.00
1	A	173	ARG	NE-CZ-NH1	-6.28	117.16	120.30
1	A	202	ARG	N-CA-C	5.89	126.91	111.00
1	A	184	LYS	C-N-CD	5.53	140.01	128.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1507	0	1444	217	0
2	B	1864	0	1814	162	0
3	C	1995	0	1901	135	0
4	D	120	0	63	10	0
All	All	5486	0	5222	457	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 457 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:GLU:HG2	1:A:185:PRO:O	1.29	1.26
1:A:158:THR:OG1	2:B:35:MET:CE	1.85	1.24
1:A:78:PHE:HB2	1:A:199:LEU:CD1	1.72	1.19
1:A:158:THR:OG1	2:B:35:MET:HE2	1.41	1.17
2:B:28:ILE:HD11	2:B:32:PRO:CB	1.78	1.13

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	190/234 (81%)	176 (93%)	12 (6%)	2 (1%)	14 46
2	B	237/253 (94%)	226 (95%)	11 (5%)	0	100 100
3	C	255/289 (88%)	231 (91%)	22 (9%)	2 (1%)	19 54
All	All	682/776 (88%)	633 (93%)	45 (7%)	4 (1%)	25 59



All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	37	GLY
1	A	185	PRO
3	C	210	VAL
1	A	204	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/209 (79%)	164 (99%)	2 (1%)	71	88
2	B	204/220 (93%)	200 (98%)	4 (2%)	55	80
3	C	222/252 (88%)	221 (100%)	1 (0%)	88	94
All	All	592/681 (87%)	585 (99%)	7 (1%)	71	88

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	196	SER
3	C	35	GLU
2	B	198	TRP
1	A	144	SER
2	B	202	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	216	ASN
3	C	39	ASN
3	C	206	ASN
2	B	151	ASN
3	C	213	ASN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	D	6/6 (100%)	4 (66%)	2 (33%)

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	D	3	U
4	D	4	U
4	D	5	U
4	D	6	U

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
4	D	1	G
4	D	5	U

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.