



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 7, 2024 – 01:27 pm GMT

PDB ID : 5MOK  
Title : Crystal structure of human IgE-Fc epsilon 3-4  
Authors : Dore, K.A.; Davies, A.M.; Drinkwater, N.; Beavil, A.J.; McDonnell, J.M.; Sutton, B.J.  
Deposited on : 2016-12-14  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	 2% 90% 7%
1	B	223	 2% 88% 8%
1	C	223	 4% 80% 9% 11%
1	D	223	 5% 81% 9% 9%
2	E	4	 75% 25%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	7	
3	H	7	
4	G	4	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MAN	F	5	-	-	-	X
3	MAN	F	6	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ig epsilon chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	208	Total 1601	C 1007	N 280	O 307	S 7	0	3	0
1	B	205	Total 1594	C 1002	N 285	O 301	S 6	0	5	0
1	C	198	Total 1536	C 964	N 272	O 294	S 6	0	3	0
1	D	202	Total 1627	C 1026	N 291	O 304	S 6	0	10	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	325	ALA	-	expression tag	UNP P01854
A	326	ASP	-	expression tag	UNP P01854
A	327	PRO	-	expression tag	UNP P01854
A	371	GLN	ASN	conflict	UNP P01854
A	383	GLN	ASN	conflict	UNP P01854
B	325	ALA	-	expression tag	UNP P01854
B	326	ASP	-	expression tag	UNP P01854
B	327	PRO	-	expression tag	UNP P01854
B	371	GLN	ASN	conflict	UNP P01854
B	383	GLN	ASN	conflict	UNP P01854
C	325	ALA	-	expression tag	UNP P01854
C	326	ASP	-	expression tag	UNP P01854
C	327	PRO	-	expression tag	UNP P01854
C	371	GLN	ASN	conflict	UNP P01854
C	383	GLN	ASN	conflict	UNP P01854
D	325	ALA	-	expression tag	UNP P01854
D	326	ASP	-	expression tag	UNP P01854
D	327	PRO	-	expression tag	UNP P01854
D	371	GLN	ASN	conflict	UNP P01854
D	383	GLN	ASN	conflict	UNP P01854

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	4	50	28	2	20	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	7	83	46	2	35	0	0	0
3	H	7	83	46	2	35	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	4	47	26	1	20	0	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 4 2	0	0
5	D	1	Total C O 7 4 3	0	0

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0



- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	132	Total O 132 132	0	0
7	B	141	Total O 141 141	0	0
7	C	133	Total O 133 133	0	0
7	D	113	Total O 113 113	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Ig epsilon chain C region





- Molecule 1: Ig epsilon chain C region



- Molecule 1: Ig epsilon chain C region



- Molecule 1: Ig epsilon chain C region



- Molecule 2: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



Chain E:  75% 25%

MAG1  
MAG2  
BMA3  
MAN4

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  43% 57%

MAG1  
MAG2  
BMA3  
MAN4  
MAN5  
MAN6  
MAN7

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  43% 57%

MAG1  
MAG2  
BMA3  
MAN4  
MAN5  
MAN6  
MAN7

- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  50% 25% 25%

MAG1  
BMA2  
MAN3  
MAN4

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.48Å 100.36Å 77.86Å 90.00° 97.35° 90.00°	Depositor
Resolution (Å)	77.22 – 2.00 77.22 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (77.22-2.00) 99.9 (77.22-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.73 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, $R_{free}$	0.202 , 0.226 0.206 , 0.228	Depositor DCC
$R_{free}$ test set	3444 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.9	Xtrriage
Anisotropy	0.234	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 49.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7173	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.3957e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, EDO, NAG, PEG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/1649	0.46	0/2252
1	B	0.25	0/1648	0.46	0/2253
1	C	0.24	0/1580	0.46	0/2155
1	D	0.25	0/1693	0.45	0/2304
All	All	0.25	0/6570	0.46	0/8964

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1601	0	1526	6	0
1	B	1594	0	1541	5	0
1	C	1536	0	1466	10	0
1	D	1627	0	1607	13	0
2	E	50	0	43	0	0
3	F	83	0	70	0	0
3	H	83	0	70	0	0
4	G	47	0	40	1	0
5	A	6	0	7	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	7	0	10	1	0
6	A	8	0	12	1	0
6	C	8	0	12	1	0
6	D	4	0	6	1	0
7	A	132	0	0	0	0
7	B	141	0	0	0	0
7	C	133	0	0	0	0
7	D	113	0	0	2	0
All	All	7173	0	6410	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 35 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:351[B]:ARG:NH1	7:D:701:HOH:O	2.22	0.71
1:C:370:VAL:O	1:C:387:ARG:NH2	2.24	0.68
1:C:371:GLN:HB2	1:C:421:THR:HB	1.80	0.63
1:A:351:ARG:NH1	6:A:606:EDO:O1	2.33	0.62
1:D:408:ARG:NH2	1:D:412:GLU:OE2	2.35	0.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/223 (93%)	203 (98%)	4 (2%)	0	100	100
1	B	206/223 (92%)	201 (98%)	5 (2%)	0	100	100
1	C	195/223 (87%)	192 (98%)	3 (2%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	206/223 (92%)	203 (98%)	3 (2%)	0	100	100
All	All	814/892 (91%)	799 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/195 (88%)	171 (100%)	0	100	100
1	B	174/195 (89%)	173 (99%)	1 (1%)	86	90
1	C	165/195 (85%)	164 (99%)	1 (1%)	86	90
1	D	179/195 (92%)	178 (99%)	1 (1%)	86	90
All	All	689/780 (88%)	686 (100%)	3 (0%)	91	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	477	GLN
1	C	337	SER
1	D	344	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates i

22 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	E	1	2,1	14,14,15	0.32	0	17,19,21	0.42	0
2	NAG	E	2	2	14,14,15	0.20	0	17,19,21	0.45	0
2	BMA	E	3	2	11,11,12	0.59	0	15,15,17	0.74	0
2	MAN	E	4	2	11,11,12	0.97	1 (9%)	15,15,17	1.23	3 (20%)
3	NAG	F	1	3,1	14,14,15	0.24	0	17,19,21	0.56	0
3	NAG	F	2	3	14,14,15	0.20	0	17,19,21	0.49	0
3	BMA	F	3	3	11,11,12	0.55	0	15,15,17	0.88	0
3	MAN	F	4	3	11,11,12	0.69	0	15,15,17	0.98	2 (13%)
3	MAN	F	5	3	11,11,12	0.66	0	15,15,17	1.04	2 (13%)
3	MAN	F	6	3	11,11,12	0.64	0	15,15,17	0.96	2 (13%)
3	MAN	F	7	3	11,11,12	0.68	0	15,15,17	1.14	2 (13%)
4	NAG	G	1	4	14,14,15	0.22	0	17,19,21	0.40	0
4	BMA	G	2	4	11,11,12	0.57	0	15,15,17	0.92	0
4	MAN	G	3	4	11,11,12	0.83	0	15,15,17	0.84	1 (6%)
4	MAN	G	4	4	11,11,12	0.88	1 (9%)	15,15,17	1.33	3 (20%)
3	NAG	H	1	3,1	14,14,15	0.23	0	17,19,21	0.39	0
3	NAG	H	2	3	14,14,15	0.33	0	17,19,21	0.49	0
3	BMA	H	3	3	11,11,12	0.59	0	15,15,17	0.92	1 (6%)
3	MAN	H	4	3	11,11,12	0.71	0	15,15,17	1.14	2 (13%)
3	MAN	H	5	3	11,11,12	0.89	0	15,15,17	0.88	0
3	MAN	H	6	3	11,11,12	0.76	0	15,15,17	1.15	2 (13%)
3	MAN	H	7	3	11,11,12	0.62	0	15,15,17	0.92	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1
2	BMA	E	3	2	-	2/2/19/22	0/1/1/1
2	MAN	E	4	2	-	2/2/19/22	0/1/1/1
3	NAG	F	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	F	2	3	-	1/6/23/26	0/1/1/1
3	BMA	F	3	3	-	2/2/19/22	0/1/1/1
3	MAN	F	4	3	-	0/2/19/22	0/1/1/1
3	MAN	F	5	3	-	0/2/19/22	0/1/1/1
3	MAN	F	6	3	-	2/2/19/22	0/1/1/1
3	MAN	F	7	3	-	2/2/19/22	0/1/1/1
4	NAG	G	1	4	-	4/6/23/26	0/1/1/1
4	BMA	G	2	4	-	0/2/19/22	0/1/1/1
4	MAN	G	3	4	-	0/2/19/22	0/1/1/1
4	MAN	G	4	4	-	1/2/19/22	0/1/1/1
3	NAG	H	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	H	2	3	-	1/6/23/26	0/1/1/1
3	BMA	H	3	3	-	0/2/19/22	0/1/1/1
3	MAN	H	4	3	-	0/2/19/22	0/1/1/1
3	MAN	H	5	3	-	1/2/19/22	0/1/1/1
3	MAN	H	6	3	-	0/2/19/22	0/1/1/1
3	MAN	H	7	3	-	1/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	4	MAN	C1-C2	2.72	1.58	1.52
4	G	4	MAN	C1-C2	2.14	1.57	1.52

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	7	MAN	C1-O5-C5	3.23	116.56	112.19
3	H	4	MAN	C1-O5-C5	2.94	116.18	112.19
2	E	4	MAN	C1-O5-C5	2.88	116.09	112.19
3	H	6	MAN	C1-O5-C5	2.86	116.06	112.19
3	F	5	MAN	C1-O5-C5	2.66	115.80	112.19

There are no chirality outliers.

5 of 26 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	2	NAG	O5-C5-C6-O6
2	E	4	MAN	O5-C5-C6-O6
4	G	1	NAG	O5-C5-C6-O6
3	F	7	MAN	O5-C5-C6-O6
4	G	1	NAG	C4-C5-C6-O6

There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	4	MAN	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	PEG	D	608	-	6,6,6	0.83	0	5,5,5	0.24	0
6	EDO	A	606	-	3,3,3	0.45	0	2,2,2	0.38	0
6	EDO	D	609	-	3,3,3	0.47	0	2,2,2	0.31	0
5	PEG	A	605	-	5,5,6	0.85	0	4,4,5	0.25	0
6	EDO	A	607	-	3,3,3	0.47	0	2,2,2	0.31	0
6	EDO	C	606	-	3,3,3	0.43	0	2,2,2	0.35	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	EDO	C	605	-	3,3,3	0.49	0	2,2,2	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	D	608	-	-	2/4/4/4	-
6	EDO	A	606	-	-	0/1/1/1	-
6	EDO	D	609	-	-	0/1/1/1	-
5	PEG	A	605	-	-	3/3/3/4	-
6	EDO	A	607	-	-	0/1/1/1	-
6	EDO	C	606	-	-	1/1/1/1	-
6	EDO	C	605	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	605	PEG	C1-C2-O2-C3
5	A	605	PEG	O2-C3-C4-O4
5	A	605	PEG	C4-C3-O2-C2
5	D	608	PEG	C1-C2-O2-C3
5	D	608	PEG	O2-C3-C4-O4

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	608	PEG	1	0
6	A	606	EDO	1	0
6	D	609	EDO	1	0
5	A	605	PEG	3	0
6	C	606	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	208/223 (93%)	0.07	4 (1%) 66 65	24, 40, 90, 143	0
1	B	205/223 (91%)	0.02	5 (2%) 59 57	24, 37, 91, 121	0
1	C	198/223 (88%)	0.05	9 (4%) 33 32	20, 37, 92, 118	0
1	D	202/223 (90%)	0.13	11 (5%) 25 24	24, 40, 85, 123	0
All	All	813/892 (91%)	0.06	29 (3%) 42 42	20, 39, 92, 143	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	370	VAL	5.7
1	D	396	THR	5.0
1	D	364	ALA	4.2
1	D	422	HIS	4.1
1	A	363	LEU	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MAN	F	5	11/12	0.56	0.45	86,89,93,93	0
4	MAN	G	4	11/12	0.60	0.30	77,89,94,100	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MAN	F	6	11/12	0.63	0.44	103,106,108,108	0
2	NAG	E	1	14/15	0.64	0.37	79,86,96,97	0
4	NAG	G	1	14/15	0.67	0.16	76,81,83,83	0
2	BMA	E	3	11/12	0.69	0.16	77,80,82,84	0
2	MAN	E	4	11/12	0.70	0.17	82,84,85,85	0
3	NAG	H	1	14/15	0.70	0.23	75,80,85,87	0
3	MAN	F	4	11/12	0.75	0.19	80,83,93,98	0
3	MAN	F	7	11/12	0.76	0.29	77,83,84,85	0
3	MAN	H	6	11/12	0.77	0.22	38,45,50,50	0
4	BMA	G	2	11/12	0.79	0.20	86,88,90,90	0
3	NAG	F	1	14/15	0.81	0.19	69,78,83,84	0
4	MAN	G	3	11/12	0.81	0.19	90,91,91,91	0
3	MAN	H	5	11/12	0.81	0.25	56,62,65,68	0
3	MAN	H	7	11/12	0.82	0.26	77,81,82,83	0
3	MAN	H	4	11/12	0.83	0.15	48,51,54,54	0
3	BMA	H	3	11/12	0.83	0.12	51,55,62,70	0
2	NAG	E	2	14/15	0.84	0.18	70,78,82,82	0
3	NAG	F	2	14/15	0.88	0.16	60,62,65,66	0
3	NAG	H	2	14/15	0.91	0.18	54,69,73,75	0
3	BMA	F	3	11/12	0.92	0.12	61,69,74,77	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain E:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	EDO	D	609	4/4	0.69	0.29	68,69,69,70	0
6	EDO	C	605	4/4	0.76	0.15	36,38,40,42	0
5	PEG	A	605	6/7	0.79	0.28	51,57,61,61	0
6	EDO	A	607	4/4	0.84	0.23	55,57,58,59	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	EDO	A	606	4/4	0.86	0.35	58,59,59,60	0
5	PEG	D	608	7/7	0.88	0.30	60,61,63,64	0
6	EDO	C	606	4/4	0.92	0.32	49,49,49,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.