



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 10:02 am BST

PDB ID : 5OFV
Title : Crystal structure of human 3-phosphoglycerate dehydrogenase in complex with 5-fluoro-2-methylbenzoic acid
Authors : Unterlass, J.E.; Basle, A.; Blackburn, T.J.; Tucker, J.; Cano, C.; Noble, M.E.M.; Curtin, N.J.
Deposited on : 2017-07-11
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	 5% 71% 12% • 13%
1	B	223	 5% 72% 13% • 13%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3386 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called D-3-phosphoglycerate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	195	Total 1539	C 957	N 275	O 292	S 15	0	9	0
1	A	195	Total 1550	C 963	N 280	O 292	S 15	0	10	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	93	MET	-	initiating methionine	UNP O43175
A	93	MET	-	initiating methionine	UNP O43175

- Molecule 2 is 5-fluoranyl-2-methyl-benzoic acid (three-letter code: 9TZ) (formula: C₈H₇FO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	F	O		
2	B	1	Total 11	C 8	F 1	O 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	F	O		
2	A	1	11	8	1	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	142	Total	O	0	0
			142	142		
3	A	133	Total	O	0	0
			133	133		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-3-phosphoglycerate dehydrogenase



- Molecule 1: D-3-phosphoglycerate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	43.34Å 45.61Å 56.27Å 98.02° 111.14° 105.85°	Depositor
Resolution (Å)	50.60 – 1.50 39.19 – 1.49	Depositor EDS
% Data completeness (in resolution range)	89.5 (50.60-1.50) 89.5 (39.19-1.49)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.224 , 0.273 0.226 , 0.271	Depositor DCC
R_{free} test set	2775 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	11.8	Xtrriage
Anisotropy	0.391	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3386	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9TZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.23	3/1573 (0.2%)	1.42	16/2120 (0.8%)
1	B	1.22	1/1562 (0.1%)	1.31	7/2106 (0.3%)
All	All	1.23	4/3135 (0.1%)	1.36	23/4226 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	194	GLU	CG-CD	5.46	1.60	1.51
1	A	238	GLY	N-CA	5.46	1.54	1.46
1	B	190	GLN	CD-OE1	5.21	1.35	1.24
1	A	194	GLU	CD-OE2	5.01	1.31	1.25

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	119[A]	ARG	NE-CZ-NH2	-16.56	112.02	120.30
1	B	119[B]	ARG	NE-CZ-NH2	-16.56	112.02	120.30
1	A	119[A]	ARG	NE-CZ-NH1	16.40	128.50	120.30
1	A	119[B]	ARG	NE-CZ-NH1	16.40	128.50	120.30
1	A	119[A]	ARG	NE-CZ-NH2	-13.17	113.72	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	119[B]	ARG	NE-CZ-NH2	-13.17	113.72	120.30
1	B	119[A]	ARG	NE-CZ-NH1	12.97	126.78	120.30
1	B	119[B]	ARG	NE-CZ-NH1	12.97	126.78	120.30
1	A	236	ARG	NE-CZ-NH1	12.63	126.62	120.30
1	A	236	ARG	NE-CZ-NH2	-11.86	114.37	120.30
1	A	294	ARG	NE-CZ-NH1	8.65	124.62	120.30
1	A	236	ARG	CD-NE-CZ	8.23	135.12	123.60
1	B	294	ARG	NE-CZ-NH1	6.78	123.69	120.30
1	B	236	ARG	NE-CZ-NH2	-6.27	117.17	120.30
1	A	236	ARG	CG-CD-NE	6.26	124.95	111.80
1	A	175	ASP	CB-CG-OD1	-5.54	113.31	118.30
1	A	175	ASP	CB-CG-OD2	5.51	123.26	118.30
1	B	175	ASP	CB-CG-OD2	5.42	123.18	118.30
1	A	158[A]	ARG	NE-CZ-NH2	5.39	123.00	120.30
1	A	158[B]	ARG	NE-CZ-NH2	5.39	123.00	120.30
1	A	115[A]	MET	CG-SD-CE	5.28	108.64	100.20
1	A	115[B]	MET	CG-SD-CE	5.28	108.64	100.20
1	A	268	ARG	CG-CD-NE	-5.01	101.28	111.80

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	100	ASN	Peptide
1	A	172	ILE	Mainchain
1	B	100	ASN	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1550	0	1566	18	3
1	B	1539	0	1551	24	1
2	A	11	0	0	1	0
2	B	11	0	0	1	0
3	A	133	0	0	4	0
3	B	142	0	0	6	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3386	0	3117	38	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:284:LEU:O	1:A:119[A]:ARG:NH2	2.08	0.87
1:B:119[A]:ARG:NH2	1:A:284:LEU:O	2.13	0.82
1:B:158:ARG:HG2	1:B:186:PHE:CZ	2.19	0.77
1:B:181[B]:GLU:HA	1:B:181[B]:GLU:OE1	1.82	0.76
1:B:194:GLU:HB3	3:B:600:HOH:O	1.86	0.74
1:B:113[B]:MET:SD	1:B:234:CYS:SG	2.85	0.74
1:B:283:HIS:O	1:A:135:ARG:NH1	2.28	0.67
1:B:175:ASP:OD1	1:B:177:ILE:HG22	1.96	0.66
1:B:115[B]:MET:HE3	1:B:115[B]:MET:HA	1.80	0.63
1:B:102:ASN:HB2	3:B:587:HOH:O	1.98	0.62
1:B:273:VAL:HG23	3:B:502:HOH:O	2.02	0.59
1:A:113[B]:MET:SD	1:A:234:CYS:SG	3.01	0.59
1:B:111:CYS:O	1:B:115[B]:MET:HG2	2.03	0.58
1:A:220[A]:ASN:ND2	3:A:503:HOH:O	2.32	0.58
1:B:265:GLU:OE1	1:B:283:HIS:HD2	1.88	0.56
1:A:146[B]:LYS:NZ	3:A:502:HOH:O	2.31	0.56
1:B:282:PRO:O	1:A:135:ARG:HG2	2.08	0.54
1:A:137:LYS:HB2	3:A:609:HOH:O	2.06	0.54
1:A:181[A]:GLU:CD	1:A:181[A]:GLU:H	2.13	0.52
1:A:100:ASN:N	1:A:103:SER:HG	2.07	0.52
1:A:265:GLU:OE1	1:A:283:HIS:HD2	1.92	0.52
1:A:175:ASP:OD1	1:A:177:ILE:HG13	2.11	0.51
1:B:154:GLY:O	1:B:158:ARG:HG3	2.11	0.50
1:B:100:ASN:N	1:B:103:SER:HG	2.10	0.50
1:A:154:GLY:O	1:A:158[A]:ARG:HG3	2.12	0.49
1:B:197:TRP:HB2	1:B:198:PRO:HD3	1.93	0.49
1:A:102:ASN:HA	3:A:515:HOH:O	2.12	0.49
1:A:267:PRO:O	1:A:270:ARG:NH1	2.47	0.48
3:B:522:HOH:O	1:A:129:LYS:HE3	2.15	0.46
1:B:158:ARG:HG2	1:B:186:PHE:CE2	2.52	0.45
1:B:183:SER:HA	1:B:186:PHE:CZ	2.51	0.44
1:B:135:ARG:HG2	1:A:282:PRO:O	2.18	0.44
1:A:217[B]:LEU:HA	1:A:217[B]:LEU:HD23	1.84	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:267:PRO:O	1:B:270:ARG:NH1	2.52	0.42
1:B:194:GLU:HB2	3:B:568:HOH:O	2.20	0.41
1:B:208:PRO:HD2	2:B:401:9TZ:O1	2.20	0.41
1:B:102:ASN:CB	3:B:587:HOH:O	2.63	0.40
2:A:401:9TZ:C7	2:A:401:9TZ:O2	2.70	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:GLU:OE2	1:A:268:ARG:NH2[1_455]	2.12	0.08
1:A:220[A]:ASN:ND2	3:B:603:HOH:O[1_444]	2.13	0.07
1:B:155:ARG:NH1	1:A:100:ASN:OD1[1_565]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/223 (91%)	199 (98%)	3 (2%)	1 (0%)	29	9
1	B	202/223 (91%)	198 (98%)	3 (2%)	1 (0%)	29	9
All	All	405/446 (91%)	397 (98%)	6 (2%)	2 (0%)	29	9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	101	GLY
1	A	101	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/182 (92%)	158 (94%)	10 (6%)	19	2
1	B	167/182 (92%)	160 (96%)	7 (4%)	30	6
All	All	335/364 (92%)	318 (95%)	17 (5%)	24	4

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	102	ASN
1	B	155	ARG
1	B	194	GLU
1	B	270	ARG
1	B	272	LEU
1	B	292	GLN
1	B	293	SER
1	A	102	ASN
1	A	155	ARG
1	A	181[A]	GLU
1	A	181[B]	GLU
1	A	194	GLU
1	A	236	ARG
1	A	270	ARG
1	A	272	LEU
1	A	292	GLN
1	A	293	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	190	GLN
1	B	283	HIS
1	A	190	GLN
1	A	283	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	9TZ	A	401	-	9,11,11	1.35	1 (11%)	10,15,15	0.89	0
2	9TZ	B	401	-	9,11,11	1.52	1 (11%)	10,15,15	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9TZ	A	401	-	-	0/0/4/4	0/1/1/1
2	9TZ	B	401	-	-	0/0/4/4	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	9TZ	C6-C8	4.46	1.51	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	9TZ	C6-C8	3.90	1.51	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	9TZ	1	0
2	B	401	9TZ	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	195/223 (87%)	0.55	11 (5%) 24 26	8, 16, 39, 51	0
1	B	195/223 (87%)	0.57	12 (6%) 20 22	9, 17, 34, 48	0
All	All	390/446 (87%)	0.56	23 (5%) 22 24	8, 17, 36, 51	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	100	ASN	7.2
1	B	100	ASN	4.7
1	A	101	GLY	4.5
1	A	293	SER	4.4
1	B	293	SER	4.4
1	B	294	ARG	3.5
1	B	182	VAL	3.3
1	B	102	ASN	3.2
1	B	177	ILE	3.2
1	B	186	PHE	3.0
1	A	294	ARG	2.9
1	A	102	ASN	2.9
1	A	177	ILE	2.7
1	B	191	LEU	2.7
1	B	178	ILE	2.7
1	A	268	ARG	2.7
1	B	101	GLY	2.6
1	A	138	PHE	2.5
1	B	268	ARG	2.4
1	A	136	LYS	2.4
1	B	181[A]	GLU	2.3
1	A	269	ASP	2.3
1	A	270	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	9TZ	B	401	11/11	0.86	0.16	29,31,34,35	0
2	9TZ	A	401	11/11	0.92	0.12	19,20,32,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.