



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 8, 2025 – 10:03 PM JST

PDB ID : 5X07 / pdb_00005x07
Title : Crystal structure of FOXA2 DNA binding domain bound to a full consensus DNA site
Authors : Li, J.; Guo, M.; Zhou, Z.; Jiang, L.; Chen, X.; Qu, L.; Wu, D.; Chen, Z.; Chen, L.; Chen, Y.
Deposited on : 2017-01-20
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Xtrriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3657 (2.80-2.80)
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)
RSRZ outliers	164620	3659 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	16	56% (green), 44% (yellow)
1	D	16	6% (red), 50% (green), 50% (yellow)
1	G	16	6% (red), 31% (green), 69% (yellow)
1	J	16	19% (red), 38% (green), 62% (yellow)
2	B	16	56% (green), 44% (yellow)
2	E	16	6% (red), 38% (green), 62% (yellow)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	16	<p>6% 31% 69%</p>
2	K	16	<p>50% 50%</p>
3	C	104	<p>3% 54% 15% 5% 7% 19%</p>
3	F	104	<p>7% 59% 13% 6% 18%</p>
3	I	104	<p>10% 56% 17% 18%</p>
3	L	104	<p>6% 61% 14% 6% 18%</p>

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 5405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*CP*TP*TP*GP*TP*TP*TP*AP*CP*AP*TP*TP*TP*TP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	16	Total	C	N	O	P	0	0	0
			321	158	46	102	15			
1	A	16	Total	C	N	O	P	0	0	0
			321	158	46	102	15			
1	G	16	Total	C	N	O	P	0	0	0
			321	158	46	102	15			
1	J	16	Total	C	N	O	P	0	0	0
			321	158	46	102	15			

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*AP*AP*AP*AP*TP*GP*TP*AP*AP*AP*CP*AP*AP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	16	Total	C	N	O	P	0	0	0
			329	158	70	86	15			
2	B	16	Total	C	N	O	P	0	0	0
			329	158	70	86	15			
2	H	16	Total	C	N	O	P	0	0	0
			329	158	70	86	15			
2	K	16	Total	C	N	O	P	0	0	0
			329	158	70	86	15			

- Molecule 3 is a protein called Hepatocyte nuclear factor 3-beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	85	Total	C	N	O	S	0	0	0
			707	458	122	123	4			
3	C	84	Total	C	N	O	S	0	0	0
			701	455	121	121	4			
3	I	82	Total	C	N	O	S	0	0	0
			690	448	119	119	4			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	85	707	458	122	123	4	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	155	GLY	-	expression tag	UNP Q9Y261
F	156	PRO	-	expression tag	UNP Q9Y261
C	155	GLY	-	expression tag	UNP Q9Y261
C	156	PRO	-	expression tag	UNP Q9Y261
I	155	GLY	-	expression tag	UNP Q9Y261
I	156	PRO	-	expression tag	UNP Q9Y261
L	155	GLY	-	expression tag	UNP Q9Y261
L	156	PRO	-	expression tag	UNP Q9Y261

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*TP*CP*TP*TP*GP*TP*TP*TP*AP*CP*AP*TP*TP*TP*TP*G)-3')



- Molecule 1: DNA (5'-D(*TP*CP*TP*TP*GP*TP*TP*TP*AP*CP*AP*TP*TP*TP*TP*G)-3')



- Molecule 1: DNA (5'-D(*TP*CP*TP*TP*GP*TP*TP*TP*AP*CP*AP*TP*TP*TP*TP*G)-3')



- Molecule 1: DNA (5'-D(*TP*CP*TP*TP*GP*TP*TP*TP*AP*CP*AP*TP*TP*TP*TP*G)-3')




- Molecule 2: DNA (5'-D(*CP*AP*AP*AP*AP*TP*GP*TP*AP*AP*AP*CP*AP*AP*GP*A)-3')





- Molecule 2: DNA (5'-D(*CP*AP*AP*AP*AP*TP*GP*TP*AP*AP*AP*CP*AP*AP*GP*A)-3')



- Molecule 2: DNA (5'-D(*CP*AP*AP*AP*AP*TP*GP*TP*AP*AP*AP*CP*AP*AP*GP*A)-3')



- Molecule 2: DNA (5'-D(*CP*AP*AP*AP*AP*TP*GP*TP*AP*AP*AP*CP*AP*AP*GP*A)-3')



- Molecule 3: Hepatocyte nuclear factor 3-beta



- Molecule 3: Hepatocyte nuclear factor 3-beta



- Molecule 3: Hepatocyte nuclear factor 3-beta





- Molecule 3: Hepatocyte nuclear factor 3-beta



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	195.13Å 71.93Å 72.16Å 90.00° 103.14° 90.00°	Depositor
Resolution (Å)	33.64 – 2.80 33.64 – 2.80	Depositor EDS
% Data completeness (in resolution range)	96.6 (33.64-2.80) 96.5 (33.64-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.240 , 0.279 0.250 , 0.278	Depositor DCC
R_{free} test set	2000 reflections (8.22%)	wwPDB-VP
Wilson B-factor (Å ²)	68.7	Xtrriage
Anisotropy	0.216	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5405	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/356	0.60	0/548
1	D	0.36	0/356	0.63	0/548
1	G	0.33	0/356	0.63	0/548
1	J	0.26	0/356	0.62	0/548
2	B	0.20	0/372	0.45	0/572
2	E	0.29	0/372	0.45	0/572
2	H	0.30	0/372	0.42	0/572
2	K	0.23	0/372	0.50	0/572
3	C	0.63	1/727 (0.1%)	1.11	12/988 (1.2%)
3	F	0.53	3/733 (0.4%)	1.03	8/996 (0.8%)
3	I	0.40	0/715	0.99	4/971 (0.4%)
3	L	0.30	0/733	0.95	6/996 (0.6%)
All	All	0.39	4/5820 (0.1%)	0.80	30/8431 (0.4%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	224	PRO	N-CD	5.50	1.55	1.47
3	F	223	SER	C-N	5.28	1.41	1.33
3	F	227	PRO	N-CD	5.11	1.54	1.47
3	C	176	PRO	N-CD	5.07	1.54	1.47

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	159	LYS	CA-C-N	6.47	127.04	120.38
3	I	159	LYS	C-N-CA	6.47	127.04	120.38
3	L	218	LEU	CA-C-N	6.42	133.26	121.70
3	L	218	LEU	C-N-CA	6.42	133.26	121.70
3	C	218	LEU	CA-C-N	6.41	133.24	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	321	0	188	5	0
1	D	321	0	188	9	44
1	G	321	0	188	18	44
1	J	321	0	188	8	0
2	B	329	0	180	4	0
2	E	329	0	180	6	46
2	H	329	0	180	9	46
2	K	329	0	180	7	0
3	C	701	0	682	39	0
3	F	707	0	687	37	0
3	I	690	0	672	36	0
3	L	707	0	687	18	0
All	All	5405	0	4200	177	90

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 177 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:222:ARG:HG3	3:C:232:PHE:CE2	1.65	1.30
3:I:237:PRO:C	3:I:238:ASP:OD1	1.76	1.27
3:F:222:ARG:CZ	3:F:229:LYS:H	1.60	1.13
3:I:223:SER:H	3:I:224:PRO:HD3	1.02	1.09
3:I:223:SER:H	3:I:224:PRO:CD	1.69	1.05

The worst 5 of 90 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:1:DT:C2	2:H:16:DA:C2[4_7410]	0.32	1.88

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:16:DA:C6	1:G:1:DT:O4[4_7410]	0.35	1.85
1:D:1:DT:O4	2:H:16:DA:N7[4_7410]	0.45	1.75
2:E:16:DA:C8	1:G:1:DT:C7[4_7410]	0.47	1.73
2:E:16:DA:O3'	1:G:1:DT:O5'[4_7410]	0.49	1.71

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	82/104 (79%)	68 (83%)	5 (6%)	9 (11%)	0	1
3	F	83/104 (80%)	73 (88%)	6 (7%)	4 (5%)	2	6
3	I	80/104 (77%)	72 (90%)	6 (8%)	2 (2%)	4	17
3	L	83/104 (80%)	76 (92%)	5 (6%)	2 (2%)	5	18
All	All	328/416 (79%)	289 (88%)	22 (7%)	17 (5%)	1	5

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	F	223	SER
3	C	223	SER
3	C	224	PRO
3	C	225	ASP
3	C	226	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	79/97 (81%)	74 (94%)	5 (6%)	15	42
3	F	80/97 (82%)	74 (92%)	6 (8%)	11	33
3	I	78/97 (80%)	72 (92%)	6 (8%)	10	31
3	L	80/97 (82%)	75 (94%)	5 (6%)	15	42
All	All	317/388 (82%)	295 (93%)	22 (7%)	13	37

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	I	225	ASP
3	L	182	LEU
3	I	238	ASP
3	L	198	GLN
3	C	182	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
3	F	187	GLN
3	C	187	GLN
3	C	236	HIS
3	I	187	GLN
3	L	187	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	16/16 (100%)	-0.04	0 100 100	48, 63, 76, 81	0
1	D	16/16 (100%)	-0.06	1 (6%) 27 21	39, 51, 80, 188	0
1	G	16/16 (100%)	0.29	1 (6%) 27 21	47, 74, 99, 249	0
1	J	16/16 (100%)	1.03	3 (18%) 4 4	61, 87, 124, 145	0
2	B	16/16 (100%)	0.14	0 100 100	44, 70, 91, 94	0
2	E	16/16 (100%)	0.28	1 (6%) 27 21	41, 57, 88, 90	0
2	H	16/16 (100%)	0.39	1 (6%) 27 21	54, 74, 96, 129	0
2	K	16/16 (100%)	0.30	0 100 100	66, 84, 101, 104	0
3	C	84/104 (80%)	0.67	3 (3%) 46 38	39, 57, 106, 140	0
3	F	85/104 (81%)	0.62	7 (8%) 19 14	38, 51, 130, 165	0
3	I	82/104 (78%)	1.02	10 (12%) 10 8	44, 68, 131, 189	0
3	L	85/104 (81%)	0.94	6 (7%) 23 18	48, 74, 116, 141	0
All	All	464/544 (85%)	0.67	33 (7%) 23 18	38, 67, 124, 249	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	229	LYS	5.2
3	I	157	HIS	4.8
3	F	155	GLY	4.8
3	I	195	PHE	4.5
3	I	229	LYS	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.