



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 14, 2023 – 01:34 AM JST

PDB ID : 5YTC
Title : Large fragment of DNA Polymerase I from *Thermus aquaticus* in a closed ternary complex with the unnatural base M-fC pair with dATP in the active site
Authors : Zeng, H.; Mondal, M.; Song, R.Y.; Zhang, J.; Xia, B.; Yi, C.Q.
Deposited on : 2017-11-17
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	539	
2	B	13	
3	C	16	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5147 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase I, thermostable.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	539	4280	2722	775	770	13	0	3	0


- Molecule 2 is a DNA chain called DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(DOC)P*(DOC))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	12	240	114	48	67	11	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(*AP*AP*AP*(92F)P*GP*GP*CP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	16	333	159	66	93	15	0	0	0

- Molecule 4 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: C₁₀H₁₆N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	30	10	5	12	3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	1	1	1	0	0
5	B	1	1	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	207	207	207	0	0
6	B	25	25	25	0	0
6	C	30	30	30	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase I, thermostable



- Molecule 2: DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(DOC)P*(DOC))-3')



- Molecule 3: DNA (5'-D(*AP*AP*AP*(92F)P*GP*GP*CP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3')



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	108.89Å 108.89Å 90.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	94.30 – 2.28 46.66 – 2.28	Depositor EDS
% Data completeness (in resolution range)	95.1 (94.30-2.28) 95.2 (46.66-2.28)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.194 , 0.258 0.199 , 0.260	Depositor DCC
R_{free} test set	1453 reflections (5.33%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtrriage
Anisotropy	0.125	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 31.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.032 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5147	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 92F, DOC, MG, DTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/4380	0.78	5/5935 (0.1%)
2	B	0.39	0/249	0.81	0/382
3	C	0.53	0/345	0.77	0/529
All	All	0.54	0/4974	0.78	5/6846 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	578	ASP	C-N-CD	-7.15	104.88	120.60
1	A	431	ARG	NE-CZ-NH1	5.73	123.17	120.30
1	A	559	ARG	NE-CZ-NH1	5.35	122.97	120.30
1	A	741	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	A	450	ARG	NE-CZ-NH1	5.07	122.84	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	299	TRP	Peptide
1	A	578	ASP	Peptide,Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4280	0	4330	26	0
2	B	240	0	134	1	0
3	C	333	0	169	2	0
4	A	30	0	12	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	207	0	0	3	1
6	B	25	0	0	0	0
6	C	30	0	0	0	0
All	All	5147	0	4645	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:691:ALA:HA	1:A:694:GLU:HG3	1.25	1.12
1:A:694:GLU:O	1:A:698:GLN:HB2	1.62	0.98
1:A:691:ALA:HA	1:A:694:GLU:CG	1.97	0.94
1:A:691:ALA:CA	1:A:694:GLU:HG3	2.02	0.88
1:A:586:VAL:HG23	1:A:595:ARG:HD2	1.70	0.72
1:A:696:TYR:CE1	6:A:1144:HOH:O	2.42	0.72
1:A:666:ASN:O	1:A:696:TYR:OH	2.10	0.67
1:A:660:ARG:NH2	2:B:113:DOC:OP2	2.24	0.65
1:A:783:VAL:HB	1:A:786:GLU:HG2	1.82	0.61
1:A:696:TYR:CD1	6:A:1144:HOH:O	2.52	0.61
1:A:691:ALA:HA	1:A:694:GLU:OE2	2.01	0.60
1:A:556[A]:ARG:HG3	1:A:556[A]:ARG:HH11	1.67	0.60
1:A:691:ALA:C	1:A:694:GLU:HG3	2.28	0.54
1:A:303:GLU:O	1:A:328:ARG:NH1	2.42	0.52
3:C:210:DC:H2'	3:C:211:DG:C8	2.44	0.52
1:A:691:ALA:HA	1:A:694:GLU:CD	2.29	0.51
1:A:654:VAL:HG22	1:A:658:MET:HE2	1.94	0.50
1:A:640:THR:HG23	1:A:654:VAL:HG11	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:668:GLY:HA2	3:C:204:92F:C2	2.43	0.48
1:A:586:VAL:HG23	1:A:595:ARG:CD	2.42	0.47
1:A:654:VAL:HG22	1:A:658:MET:CE	2.46	0.46
1:A:640:THR:HG22	1:A:651:ARG:HD3	1.99	0.45
1:A:541:LEU:HD22	1:A:585:PRO:HG3	1.99	0.44
1:A:306:PHE:O	1:A:326:ALA:HA	2.18	0.43
1:A:785:ASP:OD1	6:A:1001:HOH:O	2.21	0.43
1:A:627:ASN:O	1:A:631:VAL:HG23	2.19	0.43
1:A:690:GLN:HG2	1:A:691:ALA:N	2.35	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1172:HOH:O	6:A:1172:HOH:O[4_555]	1.51	0.69

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	540/539 (100%)	516 (96%)	19 (4%)	5 (1%)	17 18

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	300	PRO
1	A	578	ASP
1	A	579	PRO
1	A	814	ALA
1	A	586	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	438/441 (99%)	417 (95%)	21 (5%)	25 34

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	294	LEU
1	A	328	ARG
1	A	349	ARG
1	A	401	GLU
1	A	487	ARG
1	A	522	LEU
1	A	538	LEU
1	A	583	ASN
1	A	627	ASN
1	A	644	SER
1	A	649	VAL
1	A	651	ARG
1	A	679	SER
1	A	690	GLN
1	A	694	GLU
1	A	695	ARG
1	A	704	ARG
1	A	708	GLU
1	A	736	ARG
1	A	774	GLU
1	A	794	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	583	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DOC	B	113	3	16,19,20	0.82	0	20,26,29	1.04	0
3	92F	C	204	3	23,27,28	2.02	3 (13%)	27,39,42	1.42	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DOC	B	113	3	-	0/7/18/19	0/2/2/2
3	92F	C	204	3	-	0/7/23/24	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	204	92F	C24-C21	-7.24	1.29	1.43
3	C	204	92F	C21-C22	4.14	1.50	1.44
3	C	204	92F	C20-C5	-2.28	1.39	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	204	92F	C5-C20-C21	-3.15	119.74	121.61
3	C	204	92F	C21-C22-N23	-2.94	119.77	121.72
3	C	204	92F	O4'-C1'-N1	2.64	112.57	107.86
3	C	204	92F	C2'-C1'-N1	-2.48	108.05	113.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	204	92F	C22-N23-C4	2.45	121.59	115.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	113	DOC	1	0
3	C	204	92F	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DTP	A	902	5	26,32,32	0.90	2 (7%)	30,50,50	1.68	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTP	A	902	5	-	4/18/34/34	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	902	DTP	C2-N3	2.22	1.35	1.32
4	A	902	DTP	C5-C4	2.20	1.46	1.40

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	902	DTP	N3-C2-N1	-4.41	121.78	128.68
4	A	902	DTP	C2'-C1'-N9	-4.08	104.86	114.27
4	A	902	DTP	C2-N1-C6	3.14	124.12	118.75
4	A	902	DTP	PA-O3A-PB	-2.91	122.83	132.83
4	A	902	DTP	PB-O3B-PG	-2.29	124.96	132.83

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	902	DTP	PB-O3B-PG-O3G
4	A	902	DTP	PB-O3B-PG-O1G
4	A	902	DTP	PB-O3B-PG-O2G
4	A	902	DTP	PA-O3A-PB-O2B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	539/539 (100%)	0.20	47 (8%) 10 13	17, 34, 81, 98	0
2	B	11/13 (84%)	-0.81	0 100 100	18, 20, 49, 53	0
3	C	15/16 (93%)	0.35	3 (20%) 1 1	19, 27, 93, 97	1 (6%)
All	All	565/568 (99%)	0.18	50 (8%) 10 12	17, 34, 81, 98	1 (0%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	645	TRP	9.0
1	A	692	PHE	8.9
1	A	648	GLY	8.2
1	A	693	ILE	8.1
1	A	649	VAL	7.6
1	A	650	PRO	7.3
3	C	201	DA	6.7
1	A	653	ALA	6.7
1	A	651	ARG	6.3
1	A	697	PHE	6.1
1	A	654	VAL	6.1
1	A	647	PHE	6.0
1	A	696	TYR	5.7
1	A	656	PRO	5.4
1	A	662	ALA	5.3
1	A	644	SER	5.1
1	A	682	LEU	5.1
1	A	689	ALA	4.8
1	A	646	MET	4.7
1	A	683	ALA	4.7
1	A	695	ARG	4.5
1	A	655	ASP	4.4
1	A	652	GLU	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	203	DA	4.1
1	A	669	VAL	3.9
3	C	202	DA	3.8
1	A	687	GLU	3.8
1	A	699	SER	3.6
1	A	676	HIS	3.5
1	A	657	LEU	3.4
1	A	688	GLU	3.4
1	A	639	HIS	3.4
1	A	686	TYR	3.2
1	A	685	PRO	3.2
1	A	684	ILE	3.0
1	A	661	ALA	3.0
1	A	698	GLN	2.7
1	A	680	GLN	2.7
1	A	668	GLY	2.6
1	A	642	THR	2.6
1	A	643	ALA	2.5
1	A	691	ALA	2.5
1	A	658	MET	2.5
1	A	674	SER	2.5
1	A	681	GLU	2.4
1	A	667	PHE	2.4
1	A	524	GLU	2.3
1	A	637	ASP	2.2
1	A	636	ARG	2.1
1	A	673	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	92F	C	204	25/26	0.83	0.24	55,71,76,83	0
2	DOC	B	113	18/19	0.97	0.10	19,20,22,23	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	DTP	A	902	30/30	0.88	0.15	37,42,49,50	0
5	MG	B	201	1/1	0.91	0.10	37,37,37,37	0
5	MG	A	903	1/1	0.92	0.10	52,52,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.