

wwPDB NMR Structure Validation Summary Report (i)

Jun 3, 2023 – 11:14 AM EDT

PDB ID : 6BGG BMRB ID : 30367

Title : Solution NMR structures of the BRD3 ET domain in complex with a CHD4

peptide

Authors: Wai, D.C.C.; Szyszka, T.N.; Campbell, A.E.; Kwong, C.; Wilkinson-White,

L.; Silva, A.P.G.; Low, J.K.K.; Kwan, A.H.; Gamsjaeger, R.; Lu, B.; Vakoc,

C.R.; Blobel, G.A.; Mackay, J.P.

Deposited on : 2017-10-28

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-ShiftChecker} &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 83%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	NMR archive	
Metric	$(\# \mathrm{Entries})$	$(\# \mathrm{Entries})$	
Clashscore	158937	12864	
Ramachandran outliers	154571	11451	
Sidechain outliers	154315	11428	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	A	12	100%				
2	В	93	62%	12%	20%	5%	



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1	B:569-B:637 (69)	0.65	19		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 6 single-model clusters were found.

Cluster number	Models
1	1, 2, 8, 15, 18
2	5, 6, 17, 19
3	3, 9, 20
4	4, 16
Single-model clusters	7; 10; 11; 12; 13; 14



3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1636 atoms, of which 827 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called CHD4.

Mol	Chain	Residues	Atoms				Trace	
1	Λ	10	Total	С	Н	N	О	0
1	А	12	199	62	110	15	12	

• Molecule 2 is a protein called Bromodomain-containing protein 3.

Mol	Chain	Residues		\mathbf{Atoms}					Trace
2	D	00	Total	С	Н	N	О	S	0
	Б	88	1437	443	717	127	148	2	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	552	GLY	-	expression tag	UNP Q15059
В	553	PRO	-	expression tag	UNP Q15059
В	554	LEU	-	expression tag	UNP Q15059
В	555	GLY	-	expression tag	UNP Q15059
В	556	SER	-	expression tag	UNP Q15059




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: CHD4



4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 19. Colouring as in section 4.1 above.

• Molecule 1: CHD4

Chain A:

K290 V291 A292 P293 L294 K295 I296 K297 L298 G299 G300 F301

• Molecule 2: Bromodomain-containing protein 3





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: simulated annealing.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	3.97
CNS	refinement	1.2

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1039
Number of shifts mapped to atoms	1039
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	83%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	В	0.0 ± 0.0	0.5 ± 0.5
All	All	0	10

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	В	602	ARG	Sidechain	6
2	В	627	ARG	Sidechain	1
2	В	607	ARG	Sidechain	1
2	В	631	ARG	Sidechain	1
2	В	594	ARG	Sidechain	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	0	0	0	0±0
2	В	563	574	574	6±2
All	All	11260	11480	11480	115

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



5 of 34 unique clashes are listed below, so	sorted by their clash magnitude
---	---------------------------------

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Clash(A) Distance(A)		Total
2:B:600:GLN:HG2	2:B:606:LEU:HD23	0.61	1.72	8	1
2:B:625:THR:O	2:B:629:LEU:HG	0.61	1.95	13	15
2:B:578:ARG:O	2:B:582:LEU:HG	0.60	1.97	19	9
2:B:600:GLN:HG2	2:B:606:LEU:HD12	0.59	1.74	10	5
2:B:592:LEU:O	2:B:596:VAL:HG23	0.56	2.01	2	18

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	0	-	-	-	-
2	В	69/93~(74%)	68±1 (99±1%)	1±1 (1±1%)	0±0 (0±0%)	54 85
All	All	1380/2100 (66%)	1366 (99%)	13 (1%)	1 (0%)	54 85

All 1 unique Ramachandran outliers are listed below.

M	lol	Chain	Res	Type	Models (Total)
	2	В	609	SER	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	0	-	-	_
2	В	66/86 (77%)	63±2 (96±3%)	3±2 (4±3%)	36 84
All	All	1320/1900 (69%)	1269 (96%)	51 (4%)	36 84

5 of 16 unique residues with a non-rotameric sidechain are listed below. They are sorted by the



frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	В	579	GLN	14
2	В	620	THR	8
2	В	624	THR	6
2	В	634	LYS	4
2	В	614	ILE	3

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 83% for the well-defined parts and 74% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: BRD3-CHD4_NMRSTAR_3L.txt

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1039
Number of shifts mapped to atoms	1039
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	84	-0.24 ± 0.07	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	80	0.15 ± 0.09	None needed ($< 0.5 \text{ ppm}$)
¹³ C′	84	-0.22 ± 0.10	None needed ($< 0.5 \text{ ppm}$)
^{15}N	76	0.04 ± 0.21	None needed ($< 0.5 \text{ ppm}$)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 83%, i.e. 820 atoms were assigned a chemical shift out of a possible 992. 0 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	329/338 (97%)	132/136 (97%)	136/138 (99%)	61/64 (95%)
Sidechain	480/619 (78%)	319/398 (80%)	157/192 (82%)	4/29 (14%)

Continued on next page...



Continued from previous page...

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Aromatic	11/35 (31%)	7/17 (41%)	4/17 (24%)	0/1 (0%)
Overall	820/992 (83%)	458/551 (83%)	297/347 (86%)	65/94 (69%)


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)


The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:







8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1255
Intra-residue ($ i-j =0$)	321
Sequential ($ i-j =1$)	333
Medium range ($ i-j >1$ and $ i-j <5$)	336
Long range (i-j ≥5)	222
Inter-chain	43
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	12.0
Number of long range restraints per residue ¹	2.1

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	6.6	0.2
0.2-0.5 (Medium)	0.2	0.24
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Donton into topo o	Commit	07 1	Vic	olated ⁵	3	Consis	tentl	y Violated ⁴
Restraints type	Count	$\%^1$	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$
Intra-residue (i-j =0)	321	25.6	13	4.0	1.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	263	21.0	13	4.9	1.0	0	0.0	0.0
Sidechain-Sidechain	58	4.6	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	333	26.5	14	4.2	1.1	0	0.0	0.0
Backbone-Backbone	92	7.3	3	3.3	0.2	0	0.0	0.0
Backbone-Sidechain	209	16.7	9	4.3	0.7	0	0.0	0.0
Sidechain-Sidechain	32	2.5	2	6.2	0.2	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	336	26.8	10	3.0	0.8	0	0.0	0.0
Backbone-Backbone	107	8.5	2	1.9	0.2	0	0.0	0.0
Backbone-Sidechain	165	13.1	5	3.0	0.4	0	0.0	0.0
Sidechain-Sidechain	64	5.1	3	4.7	0.2	0	0.0	0.0
Long range ($ i-j \ge 5$)	222	17.7	7	3.2	0.6	0	0.0	0.0
Backbone-Backbone	3	0.2	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	90	7.2	5	5.6	0.4	0	0.0	0.0
Sidechain-Sidechain	129	10.3	2	1.6	0.2	0	0.0	0.0
Inter-chain	43	3.4	5	11.6	0.4	0	0.0	0.0
Backbone-Backbone	2	0.2	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	10	0.8	1	10.0	0.1	0	0.0	0.0
Sidechain-Sidechain	31	2.5	4	12.9	0.3	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1255	100.0	49	3.9	3.9	0	0.0	0.0
Backbone-Backbone	204	16.3	5	2.5	0.4	0	0.0	0.0
Backbone-Sidechain	737	58.7	33	4.5	2.6	0	0.0	0.0
Sidechain-Sidechain	314	25.0	11	3.5	0.9	0	0.0	0.0

 $^{^1}$ percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID		Nun	nber o	f viola	ations	5	Mean (Å)	M (Å)	${ m SD}^6$ (Å)	Madian (8)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (Å)	$SD^*(A)$	Median (Å)
1	2	0	1	1	0	4	0.12	0.14	0.01	0.12
2	1	2	2	1	0	6	0.14	0.19	0.03	0.12
3	2	5	1	0	1	9	0.14	0.24	0.04	0.14
4	2	1	0	0	1	4	0.15	0.23	0.05	0.13
5	3	2	0	1	0	6	0.14	0.23	0.04	0.12
6	3	2	1	0	2	8	0.12	0.15	0.02	0.12
7	4	0	1	0	0	5	0.13	0.15	0.01	0.13
8	2	2	1	0	1	6	0.14	0.17	0.02	0.13
9	1	2	0	0	0	3	0.14	0.15	0.01	0.15
10	0	3	1	2	0	6	0.13	0.17	0.02	0.12
11	2	2	0	1	1	6	0.14	0.18	0.03	0.12

Continued on next page...




Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	${ m SD}^6$ (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	tal Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	3	3	1	1	0	8	0.13	0.18	0.02	0.12
13	7	5	1	1	1	15	0.15	0.22	0.03	0.15
14	0	3	2	1	1	7	0.14	0.19	0.03	0.12
15	1	4	0	1	0	6	0.16	0.2	0.03	0.16
16	3	1	0	0	0	4	0.14	0.18	0.02	0.14
17	1	7	0	1	1	10	0.15	0.21	0.03	0.15
18	2	4	1	2	0	9	0.13	0.16	0.02	0.12
19	3	1	2	0	2	8	0.12	0.14	0.01	0.12
20	1	3	1	1	1	7	0.12	0.19	0.03	0.11

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1206(IR:308, SQ:319, MR:326, LR:215, IC:38) restraints are not violated in the ensemble.

Nu	Number of violated restraints						n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
3	3	7	5	4	22	1	5.0
3	4	2	1	0	10	2	10.0
4	3	0	0	0	7	3	15.0
0	0	0	0	0	0	4	20.0
1	0	1	0	0	2	5	25.0
0	0	0	0	0	0	6	30.0
1	2	0	1	0	4	7	35.0
0	1	0	0	1	2	8	40.0
0	0	0	0	0	0	9	45.0
1	1	0	0	0	2	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,1202)	2:B:619:GLU:HG2	2:B:620:THR:HG21	10	0.17	0.03	0.18
(1,1202)	2:B:619:GLU:HG2	2:B:620:THR:HG22	10	0.17	0.03	0.18
(1,1202)	2:B:619:GLU:HG2	2:B:620:THR:HG23	10	0.17	0.03	0.18
(1,1202)	2:B:619:GLU:HG3	2:B:620:THR:HG21	10	0.17	0.03	0.18
(1,1202)	2:B:619:GLU:HG3	2:B:620:THR:HG22	10	0.17	0.03	0.18
(1,1202)	2:B:619:GLU:HG3	2:B:620:THR:HG23	10	0.17	0.03	0.18
(1,266)	2:B:638:GLN:HA	2:B:638:GLN:HB2	10	0.11	0.0	0.11
(1,11)	1:A:298:LEU:HG	2:B:596:VAL:HG11	8	0.15	0.05	0.13
(1,11)	1:A:298:LEU:HG	2:B:596:VAL:HG12	8	0.15	0.05	0.13
(1,11)	1:A:298:LEU:HG	2:B:596:VAL:HG13	8	0.15	0.05	0.13
(1,1163)	2:B:606:LEU:HD11	2:B:607:ARG:HG3	8	0.14	0.02	0.14
(1,1163)	2:B:606:LEU:HD12	2:B:607:ARG:HG3	8	0.14	0.02	0.14
(1,1163)	2:B:606:LEU:HD13	2:B:607:ARG:HG3	8	0.14	0.02	0.14
(1,1163)	2:B:606:LEU:HD21	2:B:607:ARG:HG3	8	0.14	0.02	0.14
(1,1163)	2:B:606:LEU:HD22	2:B:607:ARG:HG3	8	0.14	0.02	0.14
(1,1163)	2:B:606:LEU:HD23	2:B:607:ARG:HG3	8	0.14	0.02	0.14

Continued on next page...



Continued from previous page...


Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	${ m SD}^1\ (m \AA)$	Median (Å)
(1,908)	2:B:564:GLU:HB2	2:B:565:GLU:H	7	0.15	0.03	0.14
(1,908)	2:B:564:GLU:HB3	2:B:565:GLU:H	7	0.15	0.03	0.14
(1,276)	2:B:584:ILE:HD11	2:B:592:LEU:HG	7	0.14	0.02	0.14
(1,276)	2:B:584:ILE:HD12	2:B:592:LEU:HG	7	0.14	0.02	0.14
(1,276)	2:B:584:ILE:HD13	2:B:592:LEU:HG	7	0.14	0.02	0.14
(1,981)	2:B:579:GLN:H	2:B:579:GLN:HG2	7	0.13	0.01	0.13
(1,981)	2:B:579:GLN:H	2:B:579:GLN:HG3	7	0.13	0.01	0.13
(1,605)	2:B:603:GLU:H	2:B:604:PRO:HB3	7	0.13	0.02	0.11
(1,173)	2:B:606:LEU:HA	2:B:606:LEU:HD11	5	0.13	0.01	0.13
(1,173)	2:B:606:LEU:HA	2:B:606:LEU:HD12	5	0.13	0.01	0.13
(1,173)	2:B:606:LEU:HA	2:B:606:LEU:HD13	5	0.13	0.01	0.13
(1,7)	2:B:595:VAL:HB	2:B:598:ILE:HG21	5	0.12	0.02	0.11

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table: All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,11)	1:A:298:LEU:HG	2:B:596:VAL:HG11	3	0.24
(1,11)	1:A:298:LEU:HG	2:B:596:VAL:HG12	3	0.24
(1,11)	1:A:298:LEU:HG	2:B:596:VAL:HG13	3	0.24
(1,905)	2:B:564:GLU:H	2:B:564:GLU:HG2	5	0.23
(1,905)	2:B:564:GLU:H	2:B:564:GLU:HG3	5	0.23
(1,1202)	2:B:619:GLU:HG2	2:B:620:THR:HG21	4	0.23
(1,1202)	2:B:619:GLU:HG2	2:B:620:THR:HG22	4	0.23
(1,1202)	2:B:619:GLU:HG2	2:B:620:THR:HG23	4	0.23
(1,1202)	2:B:619:GLU:HG3	2:B:620:THR:HG21	4	0.23
(1,1202)	2:B:619:GLU:HG3	2:B:620:THR:HG22	4	0.23
(1,1202)	2:B:619:GLU:HG3	2:B:620:THR:HG23	4	0.23
(1,11)	1:A:298:LEU:HG	2:B:596:VAL:HG11	13	0.22
(1,11)	1:A:298:LEU:HG	2:B:596:VAL:HG12	13	0.22
(1,11)	1:A:298:LEU:HG	2:B:596:VAL:HG13	13	0.22
(1,610)	2:B:568:GLU:HA	2:B:569:GLY:H	17	0.21
(1,908)	2:B:564:GLU:HB2	2:B:565:GLU:H	13	0.2
(1,908)	2:B:564:GLU:HB3	2:B:565:GLU:H	13	0.2
(1,904)	2:B:563:SER:HB2	2:B:564:GLU:H	3	0.2
(1,904)	2:B:563:SER:HB3	2:B:564:GLU:H	3	0.2
(1,610)	2:B:568:GLU:HA	2:B:569:GLY:H	15	0.2
(1,453)	2:B:559:ALA:HA	2:B:560:SER:H	15	0.19
(1,1202)	2:B:619:GLU:HG2	2:B:620:THR:HG21	2	0.19



10 Dihedral-angle violation analysis (i)

Dihedral angle analysis failed due to data error in the dihedral angle restraints, possibly missing target value

