



wwPDB EM Validation Summary Report ⓘ

Nov 19, 2022 – 08:14 pm GMT

PDB ID : 6EL1
EMDB ID : EMD-3885
Title : YaxAB pore complex
Authors : Braeuning, B.; Bertosin, E.; Dietz, H.; Groll, M.
Deposited on : 2017-09-27
Resolution : 6.10 Å(reported)
Based on initial models : 6EK7, 6EK8

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 6.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.















Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	
1	B	410	
1	C	410	
1	D	410	
1	E	410	
1	F	410	
1	G	410	
1	H	410	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	I	410	 67% 19% 14%
1	J	410	 68% 18% 14%
2	K	344	 80% 17% .
2	L	344	 80% 16% .
2	M	344	 81% 16% .
2	N	344	 80% 16% .
2	O	344	 77% 19% .
2	P	344	 79% 17% .
2	Q	344	 80% 16% .
2	R	344	 79% 17% .
2	S	344	 80% 16% .
2	T	344	 79% 17% .

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 54500 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YaxA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	E	351	2776	1757	474	537	8	0	0
1	A	351	2776	1757	474	537	8	0	0
1	B	351	2776	1757	474	537	8	0	0
1	C	351	2776	1757	474	537	8	0	0
1	D	351	2776	1757	474	537	8	0	0
1	F	351	2776	1757	474	537	8	0	0
1	G	351	2776	1757	474	537	8	0	0
1	H	351	2776	1757	474	537	8	0	0
1	I	351	2776	1757	474	537	8	0	0
1	J	351	2776	1757	474	537	8	0	0

- Molecule 2 is a protein called YaxB.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	O	332	2674	1691	450	532	1	0	0
2	K	332	2674	1691	450	532	1	0	0
2	L	332	2674	1691	450	532	1	0	0
2	M	332	2674	1691	450	532	1	0	0
2	N	332	2674	1691	450	532	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	P	332	2674	1691	450	532	1	0	0
2	Q	332	2674	1691	450	532	1	0	0
2	R	332	2674	1691	450	532	1	0	0
2	S	332	2674	1691	450	532	1	0	0
2	T	332	2674	1691	450	532	1	0	0


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	1	GLY	-	expression tag	UNP A1JM52
K	1	GLY	-	expression tag	UNP A1JM52
L	1	GLY	-	expression tag	UNP A1JM52
M	1	GLY	-	expression tag	UNP A1JM52
N	1	GLY	-	expression tag	UNP A1JM52
P	1	GLY	-	expression tag	UNP A1JM52
Q	1	GLY	-	expression tag	UNP A1JM52
R	1	GLY	-	expression tag	UNP A1JM52
S	1	GLY	-	expression tag	UNP A1JM52
T	1	GLY	-	expression tag	UNP A1JM52


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: YaxA




- Molecule 1: YaxA




- Molecule 1: YaxA






• Molecule 1: YaxA




• Molecule 1: YaxA




• Molecule 1: YaxA






• Molecule 1: YaxA



• Molecule 1: YaxA




• Molecule 1: YaxA







• Molecule 1: YaxA




• Molecule 2: YaxB





• Molecule 2: YaxB









• Molecule 2: YaxB




• Molecule 2: YaxB




• Molecule 2: YaxB





• Molecule 2: YaxB









• Molecule 2: YaxB



• Molecule 2: YaxB




• Molecule 2: YaxB



L343
ILE

• Molecule 2: YaxB

Chain T:  79% 17%

GLY
ALA
GLU
ILE
SER
THR
PHE
PRO
HIS
SER
GLY
L12
P15
D16
I17
N18
F19
F22
S23
Q24
K27
N28
H31
T38
F41
W42
L57
I60
L68
K71
T75
Y80
A90
M104
D118
I119
L122
L125
F126
S137
H141

I146
E147
G148
T149
Q160
R174
R192
M195
L196
V197
D198
K201
D202
L203
I204
N209
E221
L227
K232
L233
L234
F237
L271
E275
I287
K291
E299
K300
R303
A304
W305
H306
I307
F308
I309
I312
L315
E324
Y335

L336
I340
L343
ILE

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C10	Depositor
Number of particles used	25000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.145	Depositor
Minimum map value	-0.027	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.007	Depositor
Recommended contour level	0.045	Depositor
Map size (\AA)	442.39996, 442.39996, 442.39996	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.106, 1.106, 1.106	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/2808	0.51	0/3778
1	B	0.33	0/2808	0.50	0/3778
1	C	0.33	0/2808	0.51	0/3778
1	D	0.33	0/2808	0.50	0/3778
1	E	0.33	0/2808	0.50	0/3778
1	F	0.33	0/2808	0.51	0/3778
1	G	0.33	0/2808	0.51	0/3778
1	H	0.33	0/2808	0.50	0/3778
1	I	0.33	0/2808	0.51	0/3778
1	J	0.33	0/2808	0.50	0/3778
2	K	0.31	0/2697	0.54	0/3630
2	L	0.31	0/2697	0.54	0/3630
2	M	0.31	0/2697	0.54	0/3630
2	N	0.31	0/2697	0.54	0/3630
2	O	0.31	0/2697	0.54	0/3630
2	P	0.31	0/2697	0.54	0/3630
2	Q	0.31	0/2697	0.54	0/3630
2	R	0.31	0/2697	0.54	0/3630
2	S	0.31	0/2697	0.54	0/3630
2	T	0.31	0/2697	0.54	0/3630
All	All	0.32	0/55050	0.52	0/74080

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	K	0	2
2	L	0	2
2	M	0	2
2	N	0	2
2	O	0	2

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
2	P	0	2
2	Q	0	2
2	R	0	2
2	S	0	2
2	T	0	2
All	All	0	20

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 20 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	K	146	ILE	Peptide
2	K	324	GLU	Peptide
2	L	146	ILE	Peptide
2	O	146	ILE	Peptide
2	O	324	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2776	0	2857	49	0
1	B	2776	0	2857	44	0
1	C	2776	0	2857	47	0
1	D	2776	0	2857	46	0
1	E	2776	0	2857	52	0
1	F	2776	0	2857	48	0
1	G	2776	0	2857	45	0
1	H	2776	0	2857	49	0
1	I	2776	0	2857	53	0
1	J	2776	0	2857	47	0
2	K	2674	0	2789	36	0
2	L	2674	0	2789	34	0
2	M	2674	0	2789	33	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	N	2674	0	2789	35	0
2	O	2674	0	2789	45	0
2	P	2674	0	2789	39	0
2	Q	2674	0	2789	33	0
2	R	2674	0	2789	39	0
2	S	2674	0	2789	39	0
2	T	2674	0	2789	39	0
All	All	54500	0	56460	802	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 802 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:N:160:GLN:HE21	2:N:271:LEU:HG	1.59	0.67
2:P:160:GLN:HE21	2:P:271:LEU:HG	1.59	0.67
2:M:160:GLN:HE21	2:M:271:LEU:HG	1.59	0.67
2:S:160:GLN:HE21	2:S:271:LEU:HG	1.59	0.67
2:O:160:GLN:HE21	2:O:271:LEU:HG	1.59	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/410 (85%)	328 (94%)	19 (6%)	0	100	100
1	B	347/410 (85%)	328 (94%)	19 (6%)	0	100	100
1	C	347/410 (85%)	328 (94%)	19 (6%)	0	100	100
1	D	347/410 (85%)	328 (94%)	19 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	347/410 (85%)	328 (94%)	19 (6%)	0	100	100
1	F	347/410 (85%)	328 (94%)	19 (6%)	0	100	100
1	G	347/410 (85%)	328 (94%)	19 (6%)	0	100	100
1	H	347/410 (85%)	328 (94%)	19 (6%)	0	100	100
1	I	347/410 (85%)	328 (94%)	19 (6%)	0	100	100
1	J	347/410 (85%)	328 (94%)	19 (6%)	0	100	100
2	K	330/344 (96%)	299 (91%)	31 (9%)	0	100	100
2	L	330/344 (96%)	299 (91%)	31 (9%)	0	100	100
2	M	330/344 (96%)	299 (91%)	31 (9%)	0	100	100
2	N	330/344 (96%)	298 (90%)	32 (10%)	0	100	100
2	O	330/344 (96%)	299 (91%)	31 (9%)	0	100	100
2	P	330/344 (96%)	298 (90%)	32 (10%)	0	100	100
2	Q	330/344 (96%)	298 (90%)	32 (10%)	0	100	100
2	R	330/344 (96%)	298 (90%)	32 (10%)	0	100	100
2	S	330/344 (96%)	299 (91%)	31 (9%)	0	100	100
2	T	330/344 (96%)	298 (90%)	32 (10%)	0	100	100
All	All	6770/7540 (90%)	6265 (92%)	505 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/357 (86%)	305 (99%)	2 (1%)	84	90
1	B	307/357 (86%)	305 (99%)	2 (1%)	84	90
1	C	307/357 (86%)	305 (99%)	2 (1%)	84	90
1	D	307/357 (86%)	305 (99%)	2 (1%)	84	90
1	E	307/357 (86%)	305 (99%)	2 (1%)	84	90

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	307/357 (86%)	305 (99%)	2 (1%)	84	90
1	G	307/357 (86%)	305 (99%)	2 (1%)	84	90
1	H	307/357 (86%)	305 (99%)	2 (1%)	84	90
1	I	307/357 (86%)	305 (99%)	2 (1%)	84	90
1	J	307/357 (86%)	305 (99%)	2 (1%)	84	90
2	K	307/316 (97%)	306 (100%)	1 (0%)	92	94
2	L	307/316 (97%)	306 (100%)	1 (0%)	92	94
2	M	307/316 (97%)	306 (100%)	1 (0%)	92	94
2	N	307/316 (97%)	306 (100%)	1 (0%)	92	94
2	O	307/316 (97%)	306 (100%)	1 (0%)	92	94
2	P	307/316 (97%)	306 (100%)	1 (0%)	92	94
2	Q	307/316 (97%)	306 (100%)	1 (0%)	92	94
2	R	307/316 (97%)	306 (100%)	1 (0%)	92	94
2	S	307/316 (97%)	306 (100%)	1 (0%)	92	94
2	T	307/316 (97%)	306 (100%)	1 (0%)	92	94
All	All	6140/6730 (91%)	6110 (100%)	30 (0%)	89	93

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	N	174	ARG
1	J	124	ASN
2	P	174	ARG
2	T	174	ARG
1	I	124	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 172 such sidechains are listed below:

Mol	Chain	Res	Type
2	Q	24	GLN
1	I	130	ASN
2	Q	117	ASN
1	H	353	ASN
2	S	28	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-3885. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)

6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 247




Y Index: 233




Z Index: 190


6.3.2 Raw map



X Index: 151



Y Index: 232




Z Index: 165

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.045. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 621 nm^3 ; this corresponds to an approximate mass of 561 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.164 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.164 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	6.10	-	-
Author-provided FSC curve	5.91	7.03	6.16
Unmasked-calculated*	6.60	7.55	6.94

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-3885 and PDB model 6EL1. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.045 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.045).











































9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.045) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7296	 0.1780
A	 0.7600	 0.1750
B	 0.7596	 0.1760
C	 0.7611	 0.1740
D	 0.7592	 0.1750
E	 0.7614	 0.1750
F	 0.7603	 0.1740
G	 0.7578	 0.1750
H	 0.7589	 0.1760
I	 0.7581	 0.1750
J	 0.7581	 0.1760
K	 0.6976	 0.1820
L	 0.6987	 0.1810
M	 0.6964	 0.1820
N	 0.6976	 0.1810
O	 0.7010	 0.1810
P	 0.6983	 0.1800
Q	 0.7014	 0.1810
R	 0.6976	 0.1800
S	 0.6976	 0.1800
T	 0.7002	 0.1810

