



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 05:06 pm BST

PDB ID : 6FFH
Title : Crystal Structure of mGluR5 in complex with Fenobam at 2.65 Å
Authors : Christopher, J.A.; Orgovan, Z.; Congreve, M.; Dore, A.S.; Errey, J.C.; Marshall, F.H.; Mason, J.S.; Okrasa, K.; Rucktooa, P.; Serrano-Vega, M.J.; Ferenczy, G.G.; Keseru, G.M.
Deposited on : 2018-01-08
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	OLA	A	4004	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3371 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Metabotropic glutamate receptor 5, Endolysin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	414	3233	2109	538	562	24	0	0	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	566	ALA	-	expression tag	UNP P41594
A	567	ALA	-	expression tag	UNP P41594
A	568	SER	-	expression tag	UNP P41594
A	579	ALA	GLU	engineered mutation	UNP P41594
A	667	TYR	ASN	engineered mutation	UNP P41594
A	669	ALA	ILE	engineered mutation	UNP P41594
A	675	MET	GLY	engineered mutation	UNP P41594
A	1012	GLY	ARG	conflict	UNP P00720
A	1054	THR	CYS	engineered mutation	UNP P00720
A	1097	ALA	CYS	engineered mutation	UNP P00720
A	1137	ARG	ILE	conflict	UNP P00720
A	1742	ALA	THR	engineered mutation	UNP P41594
A	1753	ALA	SER	engineered mutation	UNP P41594
A	1837	ALA	-	expression tag	UNP P41594
A	1838	ALA	-	expression tag	UNP P41594
A	1839	ALA	-	expression tag	UNP P41594
A	1840	HIS	-	expression tag	UNP P41594
A	1841	HIS	-	expression tag	UNP P41594
A	1842	HIS	-	expression tag	UNP P41594
A	1843	HIS	-	expression tag	UNP P41594
A	1844	HIS	-	expression tag	UNP P41594
A	1845	HIS	-	expression tag	UNP P41594
A	1846	HIS	-	expression tag	UNP P41594
A	1847	HIS	-	expression tag	UNP P41594
A	1848	HIS	-	expression tag	UNP P41594
A	1849	HIS	-	expression tag	UNP P41594

- Molecule 2 is OLEIC ACID (three-letter code: OLA) (formula: $C_{18}H_{34}O_2$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	A	1	Total	C	O	0	0
			20	18	2		
2	A	1	Total	C	O	0	0
			15	13	2		
2	A	1	Total	C	O	0	0
			14	12	2		
2	A	1	Total	C	O	0	0
			13	11	2		
2	A	1	Total	C	O	0	0
			20	18	2		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	12	6	1	4	1	0	0

- Molecule 4 is 1-(3-chlorophenyl)-3-(3-methyl-5-oxidanylidene-4 {H}-imidazol-2-yl)urea (three-letter code: D7W) (formula: $C_{11}H_{11}ClN_4O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
4	A	1	18	11	1	4	2	0	0

- Molecule 5 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4$).



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	143.22Å 43.48Å 82.38Å 90.00° 99.22° 90.00°	Depositor
Resolution (Å)	34.50 – 2.65 34.50 – 2.65	Depositor EDS
% Data completeness (in resolution range)	87.7 (34.50-2.65) 87.8 (34.50-2.65)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.65Å)	Xtrriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.243 , 0.267 0.243 , 0.267	Depositor DCC
R_{free} test set	702 reflections (5.37%)	wwPDB-VP
Wilson B-factor (Å ²)	42.6	Xtrriage
Anisotropy	0.649	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3371	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OLA, OLC, MES, YCM, D7W

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.24	0/3286	0.39	1/4456 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1724	PRO	N-CA-CB	5.57	109.99	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3233	0	3347	21	0
2	A	82	0	120	7	0
3	A	12	0	12	0	0
4	A	18	0	0	0	0
5	A	13	0	15	0	0
6	A	13	0	0	0	0
All	All	3371	0	3494	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1011:GLU:OE1	1:A:1145:ARG:NH1	2.31	0.64
1:A:1106:MET:HE1	1:A:1114:PHE:HE2	1.62	0.64
1:A:1790:PRO:HA	2:A:4007:OLA:H22	1.80	0.63
1:A:616:GLU:HB3	1:A:1694:LEU:HD21	1.84	0.59
1:A:613:SER:OG	1:A:665:LYS:NZ	2.40	0.55
1:A:625:ILE:HG23	1:A:1810:ALA:HB1	1.87	0.55
1:A:639:PRO:HA	1:A:1731:LEU:HD13	1.91	0.52
1:A:1712:PHE:HB3	2:A:4003:OLA:H22	1.94	0.50
1:A:645:TYR:OH	2:A:4003:OLA:O1	2.25	0.49
1:A:634:YCM:HD3	1:A:646:LEU:HB3	1.94	0.48
1:A:1786:LEU:HD21	2:A:4001:OLA:H162	1.97	0.46
1:A:600:VAL:HG11	1:A:1822:VAL:HG13	1.98	0.46
1:A:1008:ARG:HB3	1:A:1008:ARG:HH11	1.81	0.45
1:A:1790:PRO:HD3	2:A:4007:OLA:H112	2.00	0.44
1:A:653:LEU:HA	1:A:653:LEU:HD23	1.89	0.43
1:A:677:LYS:HA	1:A:1003:ILE:HG22	1.99	0.43
1:A:1005:GLU:OE1	1:A:1008:ARG:NH1	2.51	0.42
1:A:1730:TYR:HE2	1:A:1800:ILE:HG12	1.84	0.42
1:A:1078:ILE:HD11	1:A:1103:VAL:HG21	2.02	0.42
2:A:4003:OLA:H82	2:A:4003:OLA:H111	1.64	0.42
1:A:1718:ASP:N	1:A:1734:ASN:OD1	2.52	0.42
1:A:1752:LEU:HD11	2:A:4001:OLA:H161	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	405/444 (91%)	394 (97%)	10 (2%)	1 (0%)	47 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1723	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	343/372 (92%)	340 (99%)	3 (1%)	78 87

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	586	PHE
1	A	649	ILE
1	A	1080	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	YCM	A	634	1	7,9,10	1.07	0	4,10,12	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	A	634	1	-	2/6/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	634	YCM	N-CA-CB-SG
1	A	634	YCM	C-CA-CB-SG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	634	YCM	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	OLA	A	4003	-	10,13,19	0.28	0	8,13,19	0.58	0
2	OLA	A	4002	-	11,14,19	0.24	0	10,14,19	0.55	0
2	OLA	A	4007	-	16,19,19	0.22	0	15,19,19	0.65	0
2	OLA	A	4004	-	9,12,19	0.29	0	8,12,19	0.82	0
5	OLC	A	4008	-	12,12,24	1.30	1 (8%)	13,13,25	1.06	1 (7%)
4	D7W	A	4006	-	18,19,19	2.57	8 (44%)	20,26,26	2.34	8 (40%)
2	OLA	A	4001	-	16,19,19	0.23	0	15,19,19	0.64	0
3	MES	A	4005	-	12,12,12	2.30	1 (8%)	14,16,16	2.05	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OLA	A	4003	-	-	3/9/11/17	-
2	OLA	A	4002	-	-	2/10/12/17	-
2	OLA	A	4007	-	-	5/15/17/17	-
2	OLA	A	4004	-	-	1/8/10/17	-
5	OLC	A	4008	-	-	2/12/12/24	-
4	D7W	A	4006	-	-	0/8/20/20	0/2/2/2
2	OLA	A	4001	-	-	2/15/17/17	-
3	MES	A	4005	-	-	4/6/14/14	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	4005	MES	C8-S	-7.72	1.66	1.77
4	A	4006	D7W	C12-N11	4.60	1.51	1.41
4	A	4006	D7W	C09-N08	4.54	1.49	1.39
4	A	4006	D7W	C09-N11	4.38	1.46	1.37
5	A	4008	OLC	O20-C1	4.32	1.46	1.33
4	A	4006	D7W	C04-N06	4.28	1.46	1.37
4	A	4006	D7W	C14-CL1	3.25	1.81	1.74
4	A	4006	D7W	C07-N08	2.81	1.47	1.37
4	A	4006	D7W	C07-N06	2.18	1.38	1.32
4	A	4006	D7W	C13-C12	2.00	1.42	1.39

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	4006	D7W	N11-C09-N08	5.70	127.06	114.23
3	A	4005	MES	C5-N4-C3	4.91	119.89	108.83
4	A	4006	D7W	O10-C09-N11	-4.10	116.68	123.62
4	A	4006	D7W	O05-C04-N06	3.43	128.53	124.42
4	A	4006	D7W	C03-C04-N06	-3.34	106.12	113.44
4	A	4006	D7W	C04-C03-N02	2.80	105.11	101.02
3	A	4005	MES	O3S-S-C8	2.76	110.24	105.77
5	A	4008	OLC	O20-C1-C2	2.72	120.45	111.91
4	A	4006	D7W	C12-C13-C14	2.68	120.81	118.69
3	A	4005	MES	C7-N4-C3	2.46	117.53	111.23
4	A	4006	D7W	C17-C12-C13	-2.45	116.75	119.65
3	A	4005	MES	O2S-S-C8	2.18	109.55	106.92
3	A	4005	MES	C7-N4-C5	2.17	116.77	111.23
3	A	4005	MES	C6-C5-N4	-2.14	106.85	110.10
4	A	4006	D7W	O05-C04-C03	2.04	125.35	122.06

There are no chirality outliers.

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	4003	OLA	C1-C2-C3-C4
2	A	4004	OLA	C1-C2-C3-C4
3	A	4005	MES	C7-C8-S-O1S
3	A	4005	MES	C7-C8-S-O3S
5	A	4008	OLC	C1-C2-C3-C4
2	A	4003	OLA	C3-C4-C5-C6
2	A	4003	OLA	C6-C7-C8-C9
2	A	4007	OLA	C5-C6-C7-C8
2	A	4007	OLA	C10-C11-C12-C13
2	A	4002	OLA	C6-C7-C8-C9
3	A	4005	MES	C8-C7-N4-C5
2	A	4007	OLA	C3-C4-C5-C6
2	A	4001	OLA	C10-C11-C12-C13
3	A	4005	MES	C7-C8-S-O2S
2	A	4007	OLA	C14-C15-C16-C17
5	A	4008	OLC	O20-C1-C2-C3
2	A	4007	OLA	C2-C3-C4-C5
2	A	4001	OLA	C9-C10-C11-C12
2	A	4002	OLA	C4-C5-C6-C7




There are no ring outliers.


3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	4003	OLA	3	0
2	A	4007	OLA	2	0
2	A	4001	OLA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	678:LYS	C	1002:ASN	N	3.05

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	413/444 (93%)	0.23	11 (2%) 54 50	27, 46, 86, 138	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	612	SER	5.0
1	A	613	SER	4.5
1	A	1036	SER	2.8
1	A	638	LYS	2.7
1	A	1729	VAL	2.5
1	A	1721	HIS	2.5
1	A	1728	GLU	2.4
1	A	609	VAL	2.3
1	A	1832	ASN	2.3
1	A	1022	GLU	2.1
1	A	606	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	YCM	A	634	10/11	0.88	0.20	44,47,55,56	0

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OLA	A	4004	13/20	0.72	0.49	76,77,79,79	0
2	OLA	A	4001	20/20	0.75	0.35	48,52,57,58	0
2	OLA	A	4007	20/20	0.76	0.36	54,59,66,66	0
2	OLA	A	4002	15/20	0.78	0.33	70,72,79,81	0
5	OLC	A	4008	13/25	0.80	0.36	68,69,71,72	0
2	OLA	A	4003	14/20	0.82	0.35	64,66,71,72	0
4	D7W	A	4006	18/18	0.89	0.24	46,50,53,54	0
3	MES	A	4005	12/12	0.91	0.23	67,72,75,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around OLA A 4001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLA A 4007:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLA A 4002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 4008:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLA A 4003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around D7W A 4006:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.