



Full wwPDB X-ray Structure Validation Report ⓘ

May 18, 2020 – 07:19 am BST

PDB ID : 6HR2
Title : Crystal structure of PROTAC 2 in complex with the bromodomain of human SMARCA4 and pVHL:ElonginC:ElonginB
Authors : Roy, M.; Bader, G.; Diers, E.; Trainor, N.; Farnaby, W.; Ciulli, A.
Deposited on : 2018-09-26
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	120	18% (Poor fit) 95% (0 outliers), 5% (1 outlier), 10% (Not modelled)
1	E	120	15% (Poor fit) 95% (0 outliers), 5% (1 outlier), 5% (Not modelled)
2	B	149	9% (Poor fit) 95% (0 outliers), 5% (1 outlier), 5% (Not modelled)
2	F	149	17% (Poor fit) 97% (0 outliers), 10% (Not modelled)
3	C	97	5% (Poor fit) 90% (0 outliers), 10% (Not modelled)
3	G	97	9% (Poor fit) 90% (0 outliers), 10% (Not modelled)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	104	 7% 98% .
4	H	104	 9% 98% ..

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 15746 atoms, of which 7698 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription activator BRG1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	117	1972	615	1005	167	182	3	1005	0	0
1	E	120	2046	637	1045	172	188	4	1045	2	0

- Molecule 2 is a protein called von Hippel-Lindau disease tumor suppressor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	149	2525	799	1264	234	226	2	1264	7	0
2	F	149	2482	786	1241	230	223	2	1241	4	0

- Molecule 3 is a protein called Elongin-C.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	C	87	1393	448	698	111	129	7	698	1	0
3	G	87	1392	448	697	111	129	7	697	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	16	MET	-	initiating methionine	UNP Q15369
G	16	MET	-	initiating methionine	UNP Q15369

- Molecule 4 is a protein called Elongin-B.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
4	D	104	1644	520	821	138	160	5	821	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
4	H	103	1624	514	811	136	158	5	811	0	0

- Molecule 5 is (2 {S},4 {R})- {N}-[[2-[2-[4-[[4-[3-azanyl-6-(2-hydroxyphenyl)pyridazin-4-yl]piperazin-1-yl]methyl]phenyl]ethoxy]-4-(4-methyl-1,3-thiazol-5-yl)phenyl]methyl]-1-[(2 {S})-2-[(1-fluoranylpropyl)carbonylamino]-3,3-dimethyl-butanoyl]-4-oxidanyl-pyrrolidine-2-carboxamide (three-letter code: FWZ) (formula: C₄₉H₅₈FN₉O₆S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
			Total	C	F	H	N	O			S
5	B	1	124	49	1	58	9	6	1	58	0
5	F	1	124	49	1	58	9	6	1	58	0

- Molecule 6 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	O	S	0	0
			4	2	1	1		
6	E	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	F	1	Total	C	O	0	0
			4	2	2		
7	H	1	Total	C	O	0	0
			4	2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	28	Total 28	O 28	0	0
8	B	78	Total 78	O 78	0	0
8	C	37	Total 37	O 37	0	0
8	D	71	Total 71	O 71	0	0
8	E	47	Total 47	O 47	0	0
8	F	71	Total 71	O 71	0	0
8	G	32	Total 32	O 32	0	0
8	H	40	Total 40	O 40	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcription activator BRG1




- Molecule 1: Transcription activator BRG1




- Molecule 2: von Hippel-Lindau disease tumor suppressor





- Molecule 2: von Hippel-Lindau disease tumor suppressor




- Molecule 3: Elongin-C




- Molecule 3: Elongin-C



- Molecule 4: Elongin-B



- Molecule 4: Elongin-B



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.33Å 61.68Å 81.60Å 69.43° 83.38° 86.42°	Depositor
Resolution (Å)	39.80 – 1.76 39.80 – 1.76	Depositor EDS
% Data completeness (in resolution range)	63.2 (39.80-1.76) 63.3 (39.80-1.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 1.76Å)	Xtrriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.216 , 0.236 0.211 , 0.229	Depositor DCC
R_{free} test set	2903 reflections (4.27%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtrriage
Anisotropy	0.017	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15746	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, EDO, FWZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/982	0.53	0/1317
1	E	0.42	0/1024	0.53	0/1372
2	B	0.41	0/1309	0.62	0/1785
2	F	0.41	0/1285	0.62	0/1751
3	C	0.43	0/718	0.55	0/967
3	G	0.43	0/709	0.54	0/955
4	D	0.44	0/839	0.61	0/1132
4	H	0.40	0/829	0.61	0/1121
All	All	0.42	0/7695	0.58	0/10400

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	967	1005	1005	0	0
1	E	1001	1045	1033	0	0
2	B	1261	1264	1242	4	0
2	F	1241	1241	1225	3	0
3	C	695	698	691	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	695	697	697	0	0
4	D	823	821	824	0	0
4	H	813	811	811	0	0
5	B	66	58	0	0	0
5	F	66	58	0	0	0
6	B	4	0	6	0	0
6	E	4	0	6	0	0
7	F	4	0	6	0	0
7	H	4	0	6	0	0
8	A	28	0	0	0	0
8	B	78	0	0	0	0
8	C	37	0	0	0	0
8	D	71	0	0	0	0
8	E	47	0	0	0	0
8	F	71	0	0	0	0
8	G	32	0	0	0	0
8	H	40	0	0	0	0
All	All	8048	7698	7552	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:139[A]:SER:HB2	2:F:139:SER:OG	2.12	0.49
2:B:115:HIS:O	2:B:138[A]:PRO:HD2	2.15	0.47
2:B:140[A]:LEU:HB2	2:F:140:LEU:HB2	1.98	0.45
2:B:133:THR:HG21	2:F:145:GLN:HG2	2.00	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/120 (96%)	114 (99%)	1 (1%)	0	100	100
1	E	120/120 (100%)	118 (98%)	2 (2%)	0	100	100
2	B	154/149 (103%)	149 (97%)	5 (3%)	0	100	100
2	F	151/149 (101%)	146 (97%)	5 (3%)	0	100	100
3	C	84/97 (87%)	84 (100%)	0	0	100	100
3	G	83/97 (86%)	82 (99%)	1 (1%)	0	100	100
4	D	102/104 (98%)	99 (97%)	3 (3%)	0	100	100
4	H	101/104 (97%)	96 (95%)	5 (5%)	0	100	100
All	All	910/940 (97%)	888 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/116 (97%)	110 (97%)	3 (3%)	44	22
1	E	118/116 (102%)	112 (95%)	6 (5%)	24	6
2	B	146/139 (105%)	140 (96%)	6 (4%)	30	10
2	F	142/139 (102%)	141 (99%)	1 (1%)	84	75
3	C	80/86 (93%)	80 (100%)	0	100	100
3	G	79/86 (92%)	79 (100%)	0	100	100
4	D	92/92 (100%)	90 (98%)	2 (2%)	52	29
4	H	91/92 (99%)	90 (99%)	1 (1%)	73	60
All	All	861/866 (99%)	842 (98%)	19 (2%)	55	29

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1488	LEU
1	A	1528	ASP
1	A	1568	ASP
2	B	140[A]	LEU
2	B	140[B]	LEU
2	B	141[A]	ASN
2	B	141[B]	ASN
2	B	182	ARG
2	B	196	LYS
4	D	37	ARG
4	D	97	PRO
1	E	1451	LEU
1	E	1488	LEU
1	E	1490	SER
1	E	1522	LEU
1	E	1528	ASP
1	E	1568	ASP
2	F	196	LYS
4	H	46	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	F	302	-	3,3,3	0.55	0	2,2,2	0.37	0
6	DMS	E	1601	-	3,3,3	0.29	0	3,3,3	0.36	0
7	EDO	H	201	-	3,3,3	0.57	0	2,2,2	0.25	0
5	FWZ	F	301	-	65,73,73	0.43	1 (1%)	85,107,107	0.86	1 (1%)
5	FWZ	B	301	-	65,73,73	0.29	0	85,107,107	0.80	1 (1%)
6	DMS	B	302	-	3,3,3	0.36	0	3,3,3	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	F	302	-	-	0/1/1/1	-
5	FWZ	B	301	-	-	2/53/81/81	0/8/8/8
7	EDO	H	201	-	-	0/1/1/1	-
5	FWZ	F	301	-	-	2/53/81/81	0/8/8/8

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	301	FWZ	C47-N44	2.09	1.45	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	301	FWZ	C36-C32-C31	4.90	126.22	116.20
5	B	301	FWZ	C36-C32-C31	3.97	124.31	116.20

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	301	FWZ	C64-C40-N41-C46
5	F	301	FWZ	C64-C40-N41-C42

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	B	301	FWZ	C64-C40-N41-C46
5	B	301	FWZ	C64-C40-N41-C42

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	117/120 (97%)	1.01	22 (18%) 1 1	31, 60, 119, 175	0
1	E	120/120 (100%)	0.73	18 (15%) 2 3	27, 49, 121, 167	0
2	B	149/149 (100%)	0.58	14 (9%) 8 11	24, 40, 118, 162	0
2	F	149/149 (100%)	1.15	25 (16%) 1 2	24, 41, 169, 219	0
3	C	87/97 (89%)	0.40	5 (5%) 23 29	29, 47, 83, 146	0
3	G	87/97 (89%)	0.83	9 (10%) 6 9	33, 49, 91, 155	0
4	D	104/104 (100%)	0.41	7 (6%) 17 23	25, 38, 124, 195	0
4	H	103/104 (99%)	0.73	9 (8%) 10 13	33, 49, 102, 202	0
All	All	916/940 (97%)	0.75	109 (11%) 4 6	24, 46, 126, 219	0

All (109) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	209	GLN	8.5
1	A	1568	ASP	8.4
4	H	101	ASP	8.0
2	F	208	HIS	7.4
2	F	181	VAL	6.3
2	F	206	ILE	6.0
2	F	180	ILE	5.7
1	A	1564	ILE	5.5
1	E	1564	ILE	5.4
2	F	182	ARG	5.4
4	H	99	LEU	5.3
1	A	1452	SER	5.3
3	C	47	SER	5.2
2	B	209	GLN	4.9
1	A	1456	PRO	4.9
1	A	1476	SER	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	102	VAL	4.8
2	B	182	ARG	4.7
2	F	173	GLU	4.7
3	G	46	LEU	4.7
3	G	47	SER	4.6
4	D	104	LYS	4.5
2	F	207	ALA	4.4
2	F	183	SER	4.3
2	F	140	LEU	4.3
2	F	169	LEU	4.3
3	G	44	ALA	4.3
2	F	184	LEU	4.2
4	D	103	MET	4.2
2	F	175	TYR	4.1
1	E	1568	ASP	4.0
2	B	180	ILE	3.9
1	A	1467	ASP	3.9
2	F	176	ARG	3.9
1	E	1450	LYS	3.7
2	B	208	HIS	3.5
2	F	172	PRO	3.5
2	B	206	ILE	3.5
2	F	178	LEU	3.5
2	B	181	VAL	3.5
1	E	1460	LYS	3.4
2	F	62	VAL	3.4
2	B	178	LEU	3.4
1	E	1456	PRO	3.4
1	A	1516	ASN	3.3
2	B	177	ARG	3.3
1	E	1458	LEU	3.3
4	D	1	MET	3.3
1	A	1566	LYS	3.2
4	H	102	VAL	3.1
1	A	1475	SER	3.1
1	E	1467	ASP	3.1
2	B	175	TYR	3.1
2	F	185	TYR	3.1
2	F	177	ARG	3.1
2	B	140[A]	LEU	3.1
1	E	1465	ILE	3.1
1	E	1469	VAL	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	185	TYR	3.1
2	F	179	ASP	3.1
4	H	103	MET	3.0
1	A	1477	SER	3.0
3	C	44	ALA	2.9
1	E	1449	GLU	2.9
2	B	188	LEU	2.9
3	C	88	THR	2.9
3	G	34	GLU	2.8
4	D	99	LEU	2.8
1	A	1454	ASN	2.8
1	E	1522	LEU	2.8
1	A	1567	GLU	2.8
1	A	1551	ILE	2.7
4	H	1	MET	2.7
1	E	1476	SER	2.7
2	F	170	VAL	2.7
2	F	188	LEU	2.7
1	E	1457	ASN	2.7
4	H	98	GLU	2.6
2	B	169	LEU	2.5
1	E	1565	GLU	2.5
3	C	17	MET	2.5
4	H	62	PHE	2.5
1	A	1545	LEU	2.5
4	D	101	ASP	2.5
1	E	1567	GLU	2.4
1	A	1502	ARG	2.4
4	D	100	PRO	2.4
1	E	1451	LEU	2.4
2	B	176	ARG	2.4
1	A	1517	HIS	2.4
3	G	22	ILE	2.3
3	G	45	MET	2.3
4	H	77	LEU	2.3
1	A	1469	VAL	2.3
1	A	1522	LEU	2.3
4	H	100	PRO	2.2
1	A	1472	TYR	2.2
2	F	186	GLU	2.2
3	C	64[A]	GLU	2.2
3	G	30	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	F	187	ASP	2.2
1	A	1455	PRO	2.1
1	E	1471	LYS	2.1
1	E	1475	SER	2.1
3	G	111	ASP	2.1
2	F	205	ARG	2.1
1	A	1471	LYS	2.1
3	G	16	MET	2.0
1	A	1499	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	EDO	F	302	4/4	0.84	0.16	42,44,47,48	0
7	EDO	H	201	4/4	0.84	0.14	65,66,66,66	0
6	DMS	E	1601	4/4	0.86	0.18	72,73,74,74	0
6	DMS	B	302	4/4	0.91	0.15	81,82,82,83	0
5	FWZ	F	301	66/66	0.94	0.12	22,31,37,38	58
5	FWZ	B	301	66/66	0.95	0.08	29,34,42,45	58

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers

There are no such residues in this entry.