



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 14, 2023 – 11:33 PM JST

PDB ID : 6IFP  
Title : X-ray Crystal Structure of Pseudoazurin Met16Ile Variant  
Authors : Sakai, C.; Yamaguchi, T.; Kohzuma, T.  
Deposited on : 2018-09-20  
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1050 (1.06-0.94)
Clashscore	141614	1117 (1.06-0.94)
Ramachandran outliers	138981	1043 (1.06-0.94)
Sidechain outliers	138945	1045 (1.06-0.94)
RSRZ outliers	127900	1023 (1.06-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	124	 23% 90% 9% .
1	C	124	 24% 85% 12% ..

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4474 atoms, of which 1986 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pseudoazurin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	124	1952	614	988	164	180	6	0	11	0
1	C	124	1974	621	998	165	184	6	0	13	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	ILE	MET	engineered mutation	UNP P19567
C	16	ILE	MET	engineered mutation	UNP P19567

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cu	0	0
			1	1		
2	C	1	Total	Cu	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	251	Total	O	0	68
			251	251		
3	C	295	Total	O	0	97
			295	295		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pseudoazurin



- Molecule 1: Pseudoazurin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	33.96Å 60.18Å 47.99Å 90.00° 96.16° 90.00°	Depositor
Resolution (Å)	30.00 – 1.00 20.55 – 1.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-1.00) 99.8 (20.55-1.00)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.40 (at 1.00Å)	Xtrriage
Refinement program	SHELX	Depositor
R, $R_{free}$	0.110 , 0.144 0.119 , 0.147	Depositor DCC
$R_{free}$ test set	5287 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	7.2	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 82.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.99	EDS
Total number of atoms	4474	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.77	2/982 (0.2%)	1.08	3/1323 (0.2%)
1	C	0.73	0/994	1.03	3/1341 (0.2%)
All	All	0.75	2/1976 (0.1%)	1.05	6/2664 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	124	ASN	C-O	-6.09	1.11	1.23
1	A	114	ARG	CG-CD	-5.22	1.39	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	94	ASP	CB-CG-OD1	8.88	126.29	118.30
1	A	46[B]	LYS	C-N-CA	6.85	136.69	122.30
1	A	46[C]	LYS	C-N-CA	6.85	136.69	122.30
1	C	103	LYS	CD-CE-NZ	5.34	123.98	111.70
1	C	100	GLU	OE1-CD-OE2	-5.23	117.02	123.30
1	C	51	ASP	CB-CG-OD2	5.16	122.95	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	45	ILE	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	964	988	970	17	5
1	C	976	998	968	25	1
2	A	1	0	0	0	0
2	C	1	0	0	0	0
3	A	251	0	0	10	3
3	C	295	0	0	16	3
All	All	2488	1986	1938	42	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:113:GLU:OE2	3:C:301[B]:HOH:O	1.63	1.15
1:C:113:GLU:HG2	3:C:301[B]:HOH:O	1.50	1.09
1:C:113:GLU:CG	3:C:301[B]:HOH:O	2.03	1.03
1:C:52:GLY:HA3	3:C:460[A]:HOH:O	1.56	1.02
1:A:84[B]:MET:HG3	3:A:481[B]:HOH:O	1.59	0.99
1:A:46[C]:LYS:HG3	3:A:489:HOH:O	1.64	0.95
1:C:60[B]:ILE:HD11	3:C:577[B]:HOH:O	1.70	0.90
1:C:113:GLU:CD	3:C:301[B]:HOH:O	2.03	0.88
1:A:46[B]:LYS:NZ	3:A:301:HOH:O	2.05	0.87
1:C:51:ASP:CG	3:C:302[A]:HOH:O	2.12	0.87
1:A:84[A]:MET:HE3	1:A:110:LYS:HZ3	1.38	0.87
1:A:84[A]:MET:HE3	1:A:110:LYS:NZ	1.89	0.86
1:C:16[B]:ILE:HG22	1:C:86:MET:HB2	1.57	0.84
1:C:60[B]:ILE:CD1	3:C:577[B]:HOH:O	2.26	0.81
1:C:81:HIS:CD2	1:C:84[B]:MET:HE2	2.16	0.80
1:C:81:HIS:CD2	1:C:84[B]:MET:CE	2.64	0.80
1:C:106:LYS:CE	3:C:479:HOH:O	2.30	0.79

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:106:LYS:HE2	3:C:479:HOH:O	1.82	0.79
1:C:81:HIS:HD2	1:C:84[B]:MET:HE2	1.47	0.78
1:A:46[C]:LYS:HE3	3:A:489:HOH:O	1.81	0.78
1:A:59[B]:LYS:HD2	3:A:483[B]:HOH:O	1.86	0.75
1:A:84[A]:MET:CE	1:A:110:LYS:NZ	2.49	0.75
1:C:51:ASP:HB3	3:C:453:HOH:O	1.93	0.68
1:C:16[B]:ILE:HD13	1:C:84[B]:MET:HE3	1.74	0.68
1:A:10:LYS:HD3	3:A:304[B]:HOH:O	1.95	0.66
1:C:16[B]:ILE:HD13	1:C:84[B]:MET:CE	2.27	0.64
1:A:84[A]:MET:HE1	1:A:110:LYS:HZ2	1.68	0.58
1:C:106:LYS:HE3	3:C:479:HOH:O	1.97	0.58
1:A:84[A]:MET:CE	1:A:110:LYS:HZ2	2.17	0.56
1:C:23[B]:LEU:HD21	3:C:329[B]:HOH:O	2.06	0.56
1:A:46[C]:LYS:CE	3:A:489:HOH:O	2.50	0.54
1:A:84[A]:MET:CE	1:A:110:LYS:HZ3	2.14	0.53
1:C:65:LYS:NZ	3:C:305:HOH:O	2.33	0.50
1:C:52:GLY:HA3	3:C:497[B]:HOH:O	2.10	0.50
1:C:22[B]:SER:OG	1:C:122:LEU:HG	2.12	0.49
1:A:46[C]:LYS:CD	3:A:489:HOH:O	2.60	0.48
1:C:69[B]:THR:HG22	3:C:464:HOH:O	2.15	0.44
1:C:16[B]:ILE:HG21	1:C:81:HIS:HB3	2.00	0.43
1:A:69:THR:HG23	3:A:313:HOH:O	2.18	0.43
1:A:84[A]:MET:HE3	1:A:110:LYS:CE	2.47	0.43
1:C:59[B]:LYS:HB2	1:C:59[B]:LYS:HE2	1.86	0.42
1:A:46[C]:LYS:CG	3:A:489:HOH:O	2.39	0.42

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12[B]:LYS:HZ1	3:C:525:HOH:O[2_545]	1.08	0.52
3:C:484:HOH:O	3:C:485:HOH:O[2_456]	1.83	0.37
1:A:13[A]:ASP:HB2	3:A:431:HOH:O[2_545]	1.26	0.34
1:A:12[B]:LYS:NZ	3:C:525:HOH:O[2_545]	1.93	0.27
1:A:13[A]:ASP:CB	3:A:431:HOH:O[2_545]	1.98	0.22
1:A:84[B]:MET:SD	3:A:439:HOH:O[1_455]	1.98	0.22
1:C:10:LYS:HZ2	1:C:100:GLU:OE2[2_546]	1.55	0.05



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/124 (103%)	126 (98%)	1 (1%)	1 (1%)	19	3
1	C	130/124 (105%)	125 (96%)	3 (2%)	2 (2%)	10	1
All	All	258/248 (104%)	251 (97%)	4 (2%)	3 (1%)	19	2

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	ILE
1	C	16[A]	ILE
1	C	16[B]	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	100/94 (106%)	99 (99%)	1 (1%)	76	47
1	C	102/94 (108%)	100 (98%)	2 (2%)	55	21
All	All	202/188 (107%)	199 (98%)	3 (2%)	73	31

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LYS
1	C	16[A]	ILE
1	C	16[B]	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	124/124 (100%)	1.52	29 (23%) 0   4	7, 10, 18, 22	0
1	C	124/124 (100%)	1.56	30 (24%) 0   3	8, 10, 17, 27	0
All	All	248/248 (100%)	1.54	59 (23%) 0   4	7, 10, 19, 27	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	16[A]	ILE	4.9
1	A	12[A]	LYS	3.6
1	A	79	THR	3.4
1	C	51	ASP	3.3
1	C	73	VAL	3.2
1	C	17[A]	VAL	3.2
1	A	60[A]	ILE	3.0
1	C	47	GLY	2.9
1	C	60[A]	ILE	2.8
1	C	84[A]	MET	2.7
1	C	95	ALA	2.7
1	A	76	VAL	2.6
1	A	90	VAL	2.6
1	A	46[B]	LYS	2.6
1	A	78	CYS	2.6
1	C	18	PHE	2.6
1	C	23[A]	LEU	2.6
1	C	31	VAL	2.5
1	A	119	LEU	2.5
1	A	87	VAL	2.5
1	A	115[A]	LEU	2.5
1	C	64	TYR	2.5
1	C	42	VAL	2.4
1	C	34	ILE	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	56	PHE	2.3
1	C	32	THR	2.3
1	A	64	TYR	2.3
1	C	8	LEU	2.3
1	A	3	PHE	2.3
1	A	57	LYS	2.2
1	A	25	VAL	2.2
1	C	87	VAL	2.2
1	A	84[A]	MET	2.2
1	C	24[A]	LYS	2.2
1	A	18	PHE	2.2
1	C	117	ALA	2.2
1	A	89	VAL	2.2
1	A	8	LEU	2.2
1	A	5	VAL	2.2
1	A	82	TYR	2.2
1	A	32	THR	2.2
1	A	123	GLY	2.1
1	C	78	CYS	2.1
1	C	90	VAL	2.1
1	C	111	ALA	2.1
1	A	30	THR	2.1
1	C	52	GLY	2.1
1	C	123	GLY	2.1
1	A	97	ALA	2.1
1	A	42	VAL	2.1
1	C	30	THR	2.1
1	C	3	PHE	2.1
1	C	121	ALA	2.0
1	C	69[A]	THR	2.0
1	A	10	LYS	2.0
1	A	23	LEU	2.0
1	A	33	PHE	2.0
1	A	68	PHE	2.0
1	C	92	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CU	A	201	1/1	1.00	0.10	9,9,9,9	0
2	CU	C	201	1/1	1.00	0.11	9,9,9,9	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.