



wwPDB X-ray Structure Validation Summary Report i

Nov 16, 2023 – 04:00 AM JST

PDB ID : 6KOP
Title : Mycobacterium tuberculosis initial transcription complex comprising sigma H and 5'-OH RNA of 9 nt
Authors : Li, L.; Zhang, Y.
Deposited on : 2019-08-12
Resolution : 3.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)
RNA backbone	3102	1117 (3.70-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...



2 Entry composition [\(i\)](#)

There are 11 unique types of molecules in this entry. The entry contains 24560 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	218	Total	C 1646	N 1038	O 281	S 325	2	0	0
1	B	227	Total	C 1606	N 1013	O 270	S 321	2	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP P9WGZ1
A	-19	GLY	-	expression tag	UNP P9WGZ1
A	-18	HIS	-	expression tag	UNP P9WGZ1
A	-17	HIS	-	expression tag	UNP P9WGZ1
A	-16	HIS	-	expression tag	UNP P9WGZ1
A	-15	HIS	-	expression tag	UNP P9WGZ1
A	-14	HIS	-	expression tag	UNP P9WGZ1
A	-13	HIS	-	expression tag	UNP P9WGZ1
A	-12	HIS	-	expression tag	UNP P9WGZ1
A	-11	HIS	-	expression tag	UNP P9WGZ1
A	-10	HIS	-	expression tag	UNP P9WGZ1
A	-9	HIS	-	expression tag	UNP P9WGZ1
A	-8	SER	-	expression tag	UNP P9WGZ1
A	-7	SER	-	expression tag	UNP P9WGZ1
A	-6	GLY	-	expression tag	UNP P9WGZ1
A	-5	HIS	-	expression tag	UNP P9WGZ1
A	-4	ILE	-	expression tag	UNP P9WGZ1
A	-3	GLU	-	expression tag	UNP P9WGZ1
A	-2	GLY	-	expression tag	UNP P9WGZ1
A	-1	ARG	-	expression tag	UNP P9WGZ1
A	0	HIS	-	expression tag	UNP P9WGZ1
B	-20	MET	-	initiating methionine	UNP P9WGZ1
B	-19	GLY	-	expression tag	UNP P9WGZ1
B	-18	HIS	-	expression tag	UNP P9WGZ1
B	-17	HIS	-	expression tag	UNP P9WGZ1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP P9WGZ1
B	-15	HIS	-	expression tag	UNP P9WGZ1
B	-14	HIS	-	expression tag	UNP P9WGZ1
B	-13	HIS	-	expression tag	UNP P9WGZ1
B	-12	HIS	-	expression tag	UNP P9WGZ1
B	-11	HIS	-	expression tag	UNP P9WGZ1
B	-10	HIS	-	expression tag	UNP P9WGZ1
B	-9	HIS	-	expression tag	UNP P9WGZ1
B	-8	SER	-	expression tag	UNP P9WGZ1
B	-7	SER	-	expression tag	UNP P9WGZ1
B	-6	GLY	-	expression tag	UNP P9WGZ1
B	-5	HIS	-	expression tag	UNP P9WGZ1
B	-4	ILE	-	expression tag	UNP P9WGZ1
B	-3	GLU	-	expression tag	UNP P9WGZ1
B	-2	GLY	-	expression tag	UNP P9WGZ1
B	-1	ARG	-	expression tag	UNP P9WGZ1
B	0	HIS	-	expression tag	UNP P9WGZ1

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1138	Total	C 8655	N 5413	O 1503	S 1701	38	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	MET	-	initiating methionine	UNP P9WGY9
C	0	VAL	-	expression tag	UNP P9WGY9

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1263	Total	C 9699	N 6086	O 1735	S 1837	41	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	MET	-	initiating methionine	UNP P9WGY7
D	1	VAL	-	expression tag	UNP P9WGY7

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	75	Total	C	N	O	0	0	0

589 375 99 115

- Molecule 5 is a protein called ECF RNA polymerase sigma factor SigH.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	159	Total	C	N	O	S	0	0

1252 788 220 240 4

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-1	GLY	-	expression tag	UNP P9WGH9
F	0	ALA	-	expression tag	UNP P9WGH9

- Molecule 6 is a DNA chain called DNA ($5'$ -D(*TP*TP*GP*TP*GP*GP*GP*AP*GP*CP*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*A)- $3'$).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	G	23	Total	C	N	O	P	0	0

475 226 89 138 22

- Molecule 7 is a DNA chain called DNA ($5'$ -D(*TP*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*GP*GP*TP*G)- $3'$).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	H	21	Total	C	N	O	P	0	0

434 206 82 126 20

- Molecule 8 is a RNA chain called RNA ($5'$ -R(*CP*AP*CP*CP*CP*UP*CP*GP*A)- $3'$).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	I	9	Total	C	N	O	P	0	0

184 84 32 60 8

- Molecule 9 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	2	Total	Zn	0	0

2 2

- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	D	1	Total Mg 1 1	0	0

- Molecule 11 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	1	Total O 1 1	0	0
11	C	7	Total O 7 7	0	0
11	D	8	Total O 8 8	0	0
11	H	1	Total O 1 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase subunit alpha




- Molecule 1: DNA-directed RNA polymerase subunit alpha





- Molecule 2: DNA-directed RNA polymerase subunit beta






- Molecule 3: DNA-directed RNA polymerase subunit beta'






- Molecule 4: DNA-directed RNA polymerase subunit omega



- Molecule 5: ECF RNA polymerase sigma factor SigH




- Molecule 6: DNA (5'-D(*TP*TP*GP*TP*GP*GP*GP*AP*GP*CP*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*A)-3')



- Molecule 7: DNA (5'-D(*TP*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*GP*GP*GP*TP*G)-3')



- Molecule 8: RNA (5'-R(*CP*AP*CP*CP*CP*UP*CP*GP*A)-3')

Chain I:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	127.37 Å 162.64 Å 133.42 Å 90.00° 117.95° 90.00°	Depositor
Resolution (Å)	49.25 – 3.30 49.25 – 3.30	Depositor EDS
% Data completeness (in resolution range)	98.3 (49.25-3.30) 98.4 (49.25-3.30)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.81 (at 3.33 Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R , R_{free}	0.217 , 0.255 0.217 , 0.256	Depositor DCC
R_{free} test set	2401 reflections (3.39%)	wwPDB-VP
Wilson B-factor (Å ²)	107.2	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 57.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	24560	wwPDB-VP
Average B, all atoms (Å ²)	121.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1671	0.43	0/2274
1	B	0.24	0/1629	0.45	0/2233
2	C	0.25	0/8813	0.43	0/11978
3	D	0.24	0/9860	0.42	0/13357
4	E	0.23	0/601	0.43	0/818
5	F	0.24	0/1274	0.40	0/1724
6	G	0.53	0/533	0.94	0/823
7	H	0.52	0/487	0.93	0/752
8	I	0.28	0/204	0.82	0/315
All	All	0.26	0/25072	0.47	0/34274

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1646	0	1676	27	0
1	B	1606	0	1543	22	0
2	C	8655	0	8429	122	0
3	D	9699	0	9624	141	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	589	0	578	16	0
5	F	1252	0	1214	17	0
6	G	475	0	261	10	0
7	H	434	0	238	11	0
8	I	184	0	99	0	0
9	D	2	0	0	0	0
10	D	1	0	0	0	0
11	A	1	0	0	0	0
11	C	7	0	0	0	0
11	D	8	0	0	0	0
11	H	1	0	0	0	0
All	All	24560	0	23662	326	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 326 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:122:THR:HG22	2:C:163:ASN:H	1.40	0.85
2:C:652:ILE:HD11	2:C:682:PRO:HB3	1.63	0.81
3:D:925:LEU:HA	3:D:962:VAL:HA	1.66	0.76
3:D:1050:THR:HG23	3:D:1107:VAL:HG23	1.67	0.76
4:E:31:PRO:HB2	4:E:36:ASN:HB3	1.67	0.75

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	214/368 (58%)	210 (98%)	4 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	223/368 (61%)	215 (96%)	8 (4%)	0	100	100
2	C	1136/1174 (97%)	1095 (96%)	41 (4%)	0	100	100
3	D	1257/1317 (95%)	1226 (98%)	31 (2%)	0	100	100
4	E	71/110 (64%)	70 (99%)	1 (1%)	0	100	100
5	F	155/218 (71%)	150 (97%)	5 (3%)	0	100	100
All	All	3056/3555 (86%)	2966 (97%)	90 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/315 (59%)	183 (99%)	2 (1%)	73	85
1	B	165/315 (52%)	162 (98%)	3 (2%)	59	78
2	C	925/995 (93%)	913 (99%)	12 (1%)	69	82
3	D	1007/1096 (92%)	991 (98%)	16 (2%)	62	79
4	E	63/90 (70%)	60 (95%)	3 (5%)	25	56
5	F	125/175 (71%)	120 (96%)	5 (4%)	31	61
All	All	2470/2986 (83%)	2429 (98%)	41 (2%)	60	78

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	938	VAL
4	E	55	TYR
3	D	1130	VAL
3	D	1246	ASN
5	F	143	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	124	HIS
3	D	767	HIS

5.3.3 RNA [\(i\)](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
8	I	8/9 (88%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	218/368 (59%)	-0.06	4 (1%) 68 67	75, 107, 150, 192	0
1	B	227/368 (61%)	0.18	12 (5%) 26 24	116, 158, 181, 191	0
2	C	1138/1174 (96%)	0.05	18 (1%) 72 70	68, 103, 172, 200	0
3	D	1263/1317 (95%)	0.11	45 (3%) 42 40	74, 115, 182, 215	0
4	E	75/110 (68%)	0.13	2 (2%) 54 52	120, 143, 166, 172	0
5	F	159/218 (72%)	0.60	14 (8%) 10 10	102, 151, 174, 197	2 (1%)
6	G	23/23 (100%)	-0.15	0 100 100	112, 154, 180, 190	3 (13%)
7	H	21/21 (100%)	-0.59	0 100 100	76, 91, 167, 177	0
8	I	9/9 (100%)	-0.10	0 100 100	71, 79, 130, 133	0
All	All	3133/3608 (86%)	0.10	95 (3%) 50 49	68, 117, 176, 215	5 (0%)

The worst 5 of 95 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	F	122	SER	7.9
3	D	1093	ASP	7.0
2	C	229	THR	4.9
2	C	1157	GLY	4.7
5	F	101	PRO	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	ZN	D	2002	1/1	0.94	0.13	150,150,150,150	0
9	ZN	D	2001	1/1	0.99	0.11	117,117,117,117	0
10	MG	D	2003	1/1	0.99	0.33	81,81,81,81	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.