



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 12:27 PM BST

PDB ID : 6N29  
Title : Crystal structure of monomeric von Willebrand Factor D'D3 assembly  
Authors : Dong, X.; Arndt, J.W.; Springer, T.A.  
Deposited on : 2018-11-12  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	
1	B	483	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called von Willebrand factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	478	3645	2246	633	708	58	0	1	0
1	B	475	3628	2236	630	704	58	0	1	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	852	ARG	GLN	conflict	UNP P04275
A	1099	ALA	CYS	conflict	UNP P04275
A	1142	ALA	CYS	conflict	UNP P04275
A	1245	PRO	-	expression tag	UNP P04275
A	1246	ARG	-	expression tag	UNP P04275
B	852	ARG	GLN	conflict	UNP P04275
B	1099	ALA	CYS	conflict	UNP P04275
B	1142	ALA	CYS	conflict	UNP P04275
B	1245	PRO	-	expression tag	UNP P04275
B	1246	ARG	-	expression tag	UNP P04275

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucofuranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	B	1	14	8	1	5	0	0
2	B	1	14	8	1	5	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	B	1	1	1	0	0
3	A	2	2	2	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	47	Total O 47 47	0	0
5	B	53	Total O 53 53	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: von Willebrand factor



- Molecule 1: von Willebrand factor



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	174.69Å 174.69Å 104.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.45 – 2.50 48.45 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.45-2.50) 98.7 (48.45-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.02 (at 2.51Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, $R_{free}$	0.202 , 0.241 0.205 , 0.245	Depositor DCC
$R_{free}$ test set	2102 reflections (3.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	78.3	Xtrriage
Anisotropy	0.054	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 66.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7460	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	101.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.86% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PEG, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/3724	0.50	0/5061
1	B	0.31	0/3706	0.50	0/5036
All	All	0.31	0/7430	0.50	0/10097

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3645	0	3444	25	0
1	B	3628	0	3430	30	0
2	A	42	0	39	0	0
2	B	28	0	26	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	7	0	10	0	0
4	B	7	0	10	0	0
5	A	47	0	0	1	0
5	B	53	0	0	4	0
All	All	7460	0	6959	54	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1122:THR:HG23	1:B:1125:LEU:H	1.48	0.77
1:A:791:THR:HG23	1:A:793:GLN:H	1.56	0.70
1:A:1122:THR:HG23	1:A:1125:LEU:H	1.57	0.69
1:B:791:THR:HG23	1:B:793:GLN:H	1.58	0.67
1:A:965:LEU:HD21	1:A:1166:PRO:HD3	1.80	0.63
1:B:1225:CYS:HB3	1:B:1232:LEU:HD11	1.83	0.60
1:A:945:ARG:NH1	1:B:946:PRO:O	2.27	0.60
1:B:921:CYS:SG	1:B:922:LYS:N	2.78	0.57
1:B:787:GLU:HB2	5:B:1415:HOH:O	2.04	0.57
1:A:1204:ARG:NH1	5:A:1403:HOH:O	2.37	0.56
1:B:939:GLY:O	1:B:976:ARG:NH1	2.38	0.56
1:B:952:HIS:CE1	1:B:968:LYS:HE3	2.41	0.56
1:A:1052:LYS:NZ	1:A:1091:CYS:O	2.38	0.54
1:B:1206:PHE:CD1	1:B:1212:VAL:HG21	2.43	0.53
1:A:1123:ALA:HA	1:A:1127:PRO:HB3	1.90	0.53
1:A:925:VAL:HB	1:A:936:LEU:HB2	1.92	0.52
1:A:777:PRO:HG2	1:A:780:ASN:HB2	1.92	0.52
1:B:985:LYS:NZ	5:B:1403:HOH:O	2.44	0.51
1:B:953:PHE:HA	1:B:965:LEU:O	2.10	0.51
1:B:796:ASP:N	1:B:796:ASP:OD1	2.37	0.51
1:B:965:LEU:HD21	1:B:1166:PRO:HD3	1.91	0.50
1:B:925:VAL:HB	1:B:936:LEU:HB2	1.94	0.50
1:B:969:ALA:HB1	1:B:985:LYS:HB2	1.93	0.50
1:A:1046:CYS:HB3	1:A:1053[B]:GLN:HG2	1.94	0.49
1:A:796:ASP:N	1:A:796:ASP:OD1	2.45	0.48
1:B:791:THR:HG22	1:B:794:ASN:OD1	2.14	0.47
1:A:965:LEU:CD2	1:A:1166:PRO:HD3	2.44	0.47
1:B:1052:LYS:NZ	1:B:1091:CYS:O	2.32	0.47
1:A:870:ILE:HD11	1:A:876:LEU:HD22	1.97	0.46
1:B:786:LEU:HD11	1:B:807:GLY:HA3	1.98	0.46
1:A:953:PHE:HA	1:A:965:LEU:O	2.16	0.46
1:B:953:PHE:O	1:B:1210:LYS:NZ	2.49	0.46
1:A:791:THR:HG22	1:A:794:ASN:OD1	2.17	0.45
1:B:881:LEU:HB2	1:B:1003:GLN:HG3	1.99	0.45
1:A:921:CYS:SG	1:A:922:LYS:N	2.89	0.44
1:B:894:VAL:HG11	1:B:982:VAL:HG11	1.98	0.44
1:B:1156:THR:HA	1:B:1182:ILE:HG23	2.00	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1150:ALA:HB3	1:A:1174:HIS:CE1	2.53	0.44
1:A:1156:THR:HA	1:A:1182:ILE:HG23	1.99	0.44
1:B:1018:PRO:HD2	5:B:1431:HOH:O	2.18	0.43
1:A:1112:ALA:HB1	1:A:1149:CYS:SG	2.59	0.43
1:A:881:LEU:HB2	1:A:1003:GLN:HG3	2.01	0.43
1:B:855:LYS:HG3	5:B:1410:HOH:O	2.18	0.42
1:B:1100:PHE:HE2	1:B:1125:LEU:HD11	1.84	0.42
1:A:782:ARG:HA	1:A:782:ARG:HD3	1.84	0.42
1:B:927:ILE:HG21	1:B:966:LEU:HD13	2.01	0.42
1:A:826:ARG:HD3	1:A:826:ARG:HA	1.81	0.42
1:B:881:LEU:HD21	1:B:1007:LEU:HG	2.01	0.41
1:B:794:ASN:HA	1:B:797:LEU:HD12	2.02	0.41
1:A:1133:ARG:NH2	1:A:1173:CYS:HB2	2.35	0.41
1:A:894:VAL:HG11	1:A:982:VAL:HG11	2.03	0.41
1:A:1208:SER:OG	1:A:1228:ASP:O	2.35	0.41
1:B:785:GLY:N	1:B:804:CYS:HB2	2.36	0.40
1:B:1199:CYS:HB3	1:B:1232:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	477/483 (99%)	455 (95%)	21 (4%)	1 (0%)	47	68
1	B	474/483 (98%)	453 (96%)	20 (4%)	1 (0%)	47	68
All	All	951/966 (98%)	908 (96%)	41 (4%)	2 (0%)	47	68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1044	ALA

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	1044	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	420/427 (98%)	417 (99%)	3 (1%)	84 94
1	B	418/427 (98%)	414 (99%)	4 (1%)	76 90
All	All	838/854 (98%)	831 (99%)	7 (1%)	81 93

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1031	CYS
1	A	1149	CYS
1	A	1165	CYS
1	B	1031	CYS
1	B	1149	CYS
1	B	1165	CYS
1	B	1181	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PEG	A	1306	-	6,6,6	0.49	0	5,5,5	0.29	0
2	NAG	A	1302	1	14,14,15	0.38	0	17,19,21	0.49	0
2	NAG	A	1301	1	14,14,15	0.30	0	17,19,21	0.39	0
2	NAG	B	1301	1	14,14,15	0.28	0	17,19,21	0.45	0
2	NAG	A	1303	1	14,14,15	0.30	0	17,19,21	0.47	0
2	NAG	B	1302	1	14,14,15	0.29	0	17,19,21	0.38	0
4	PEG	B	1304	-	6,6,6	0.49	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	1306	-	-	1/4/4/4	-
2	NAG	A	1302	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1301	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1301	1	-	2/6/23/26	0/1/1/1
2	NAG	A	1303	1	-	1/6/23/26	0/1/1/1
2	NAG	B	1302	1	-	0/6/23/26	0/1/1/1
4	PEG	B	1304	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1301	NAG	O5-C5-C6-O6
2	B	1301	NAG	C4-C5-C6-O6
2	A	1303	NAG	O5-C5-C6-O6
4	A	1306	PEG	C1-C2-O2-C3
4	B	1304	PEG	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	478/483 (98%)	0.66	45 (9%) <b>8</b> <b>8</b>	52, 93, 172, 293	0
1	B	475/483 (98%)	0.72	43 (9%) <b>9</b> <b>9</b>	52, 94, 163, 249	0
All	All	953/966 (98%)	0.69	88 (9%) <b>9</b> <b>9</b>	52, 94, 170, 293	0

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	781	LEU	11.0
1	B	781	LEU	10.6
1	B	782	ARG	7.6
1	A	783	ALA	7.1
1	B	800	MET	7.0
1	B	802	MET	6.9
1	B	779	ASP	6.7
1	A	779	ASP	6.4
1	A	802	MET	6.1
1	A	853	ASP	5.2
1	A	837	ALA	5.1
1	B	830	PHE	4.9
1	A	916	HIS	4.8
1	B	1239	GLU	4.7
1	A	850	VAL	4.6
1	A	838	PRO	4.5
1	B	842	VAL	4.3
1	A	830	PHE	4.3
1	A	836	TYR	4.2
1	A	795	TYR	4.2
1	B	918	SER	4.2
1	B	765	LEU	4.1
1	A	855	LYS	4.1
1	B	836	TYR	4.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	1212	VAL	4.0
1	A	800	MET	3.9
1	A	854	ARG	3.9
1	B	916	HIS	3.8
1	B	792	CYS	3.7
1	B	835	GLU	3.7
1	A	841	THR	3.7
1	B	902	PRO	3.7
1	A	852	ARG	3.7
1	A	1192	ASP	3.7
1	B	840	GLU	3.6
1	B	1206	PHE	3.5
1	A	902	PRO	3.4
1	B	822	VAL	3.2
1	A	840	GLU	3.2
1	A	851	CYS	3.1
1	B	1179	PRO	3.1
1	B	1201	VAL	3.1
1	B	778	ALA	3.1
1	B	809	LEU	3.1
1	A	1031	CYS	2.9
1	A	787	GLU	2.9
1	B	827	CYS	2.9
1	B	855	LYS	2.9
1	B	853	ASP	2.8
1	A	914	CYS	2.8
1	A	1143	GLU	2.8
1	A	809	LEU	2.8
1	B	795	TYR	2.8
1	B	1232	LEU	2.8
1	B	1181	LYS	2.8
1	A	915	SER	2.7
1	A	764	SER	2.7
1	B	837	ALA	2.6
1	B	834	LYS	2.6
1	A	789	THR	2.5
1	A	828	PRO	2.5
1	A	1015	GLU	2.5
1	A	782	ARG	2.5
1	A	799	CYS	2.4
1	B	1014	VAL	2.4
1	B	1145	ARG	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	1193	PRO	2.4
1	B	817	HIS	2.3
1	B	839	GLY	2.3
1	A	1179	PRO	2.3
1	B	919	VAL	2.3
1	A	831	HIS	2.3
1	A	833	GLY	2.2
1	A	834	LYS	2.2
1	A	786	LEU	2.2
1	B	1131	GLU	2.2
1	B	1144	TRP	2.2
1	A	1194	GLU	2.2
1	A	1145	ARG	2.2
1	A	1016	GLU	2.1
1	A	839	GLY	2.1
1	A	1144	TRP	2.1
1	B	1188	GLN	2.1
1	B	1182	ILE	2.1
1	B	799	CYS	2.1
1	B	1194	GLU	2.1
1	B	1191	VAL	2.0
1	A	1035	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PEG	A	1306	7/7	0.64	0.32	115,119,130,134	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	A	1303	14/15	0.69	0.31	137,163,181,184	0
4	PEG	B	1304	7/7	0.74	0.40	118,137,143,146	0
3	CA	A	1305	1/1	0.75	0.05	144,144,144,144	0
2	NAG	A	1301	14/15	0.80	0.44	137,157,164,167	0
2	NAG	B	1301	14/15	0.83	0.24	132,145,167,171	0
2	NAG	B	1302	14/15	0.91	0.30	106,128,140,148	0
2	NAG	A	1302	14/15	0.92	0.13	105,116,127,129	0
3	CA	A	1304	1/1	0.98	0.11	79,79,79,79	0
3	CA	B	1303	1/1	0.98	0.27	101,101,101,101	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.