

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2023 – 01:47 AM EDT

PDB ID : 6NH3

Title : Structure of human endothelial nitric oxide synthase heme domain in complex

with (S)-6-(3-fluoro-5-(2-(pyrrolidin-2-yl)ethyl)phenethyl)-4-methylpyridin-2-

amine

Authors : Chreifi, G.; Li, H.; Poulos, T.L.

Deposited on : 2018-12-21

Resolution : 2.01 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as 541 be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)


There are 9 unique types of molecules in this entry. The entry contains 13985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Endothelial nitric oxide synthase splice variant eNOS13A.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	404	Total	С	N	О	S	0	2	0
1 A	404	3237	2062	570	589	16	0	2		
1	В	402	Total	С	N	О	S	0	3	0
1	Б	402	3221	2051	566	587	17	0		
1	C	401	Total	С	N	О	S	0	1	0
1		401	3203	2040	563	584	16	0		
1	D	402	Total	С	N	О	S	0	2	0
	ט	402	3221	2051	566	587	17	0	3	U

• Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf				
9	A	Λ	1	Total	С	Fe	N	О	0	0			
		1	43	34	1	4	4		0				
2	В	D	B	R	D	1	Total	С	Fe	N	О	0	0
2		В		43	34	1	4	4	0				


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf							
2	С		1	Total	С	Fe	N	О	0	0				
2		1	43	34	1	4	4		0					
9	D	D 1	D	D	D	D	1	Total	С	Fe	N	О	0	0
2			43	34	1	4	4							

• Molecule 3 is 6-[2-(3-fluoro-5-{2-[(2S)-pyrrolidin-2-yl]ethyl}phenyl)ethyl]-4-methylpyridin-2 -amine (three-letter code: KL4) (formula: $C_{20}H_{26}FN_3$).




Mol	Chain	Residues	A	ton	ıs		ZeroOcc	AltConf	
3	A	1	Total	С	F	N	0	0	
3	A	1	24	20	1	3	U	0	
3	A	1	Total	С	F	N	0	0	
5		1	24	20	1	3		U	
3	В	1	Total	С	F	N	0	0	
3	D	1	24	20	1	3	U		
3	В	1	Total	\mathbf{C}	F	Ν	0	0	
3	D	D 1	24	20	1	3	O		
3	C	C	1	Total	\mathbf{C}	F	N	0	0
	C	1	24	20	1	3	U	U	
3	$^{\rm C}$	1	Total	\mathbf{C}	F	N	0	0	
		1	24	20	1	3	U	0	
3	$^{\rm C}$	1	Total	\mathbf{C}	F	N	0	0	
ა	C	1	24	20	1	3	U		
3	D	1	Total	С	F	Ν	0	0	
		1	24	20	1	3	0	0	

 \bullet Molecule 4 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN



E-1,3-DIOL (three-letter code: BTB) (formula: $\mathrm{C_8H_{19}NO_5}).$




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 14 8 1 5	0	0
4	A	1	Total C N O 14 8 1 5	0	0
4	В	1	Total C N O 14 8 1 5	0	0
4	В	1	Total C N O 14 8 1 5	0	0
4	В	1	Total C N O 14 8 1 5	0	0
4	С	1	Total C N O 14 8 1 5	0	0
4	С	1	Total C N O 14 8 1 5	0	0
4	D	1	Total C N O 14 8 1 5	0	0
4	D	1	Total C N O 14 8 1 5	0	0

• Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total Zn 3 3	0	0
5	С	3	Total Zn 3 3	0	0



• Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	С	1	Total C O 6 3 3	0	0

• Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Cl 1 1	0	0
7	В	1	Total Cl 1 1	0	0
7	С	1	Total Cl 1 1	0	0
7	D	1	Total Cl 1 1	0	0

• Molecule 8 is GADOLINIUM ATOM (three-letter code: GD) (formula: Gd).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Gd 1 1	0	0
8	В	1	Total Gd 1 1	0	0
8	С	1	Total Gd 1 1	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	D	1	Total Gd 1 1	0	0

• Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	93	Total O 93 93	0	0
9	В	164	Total O 164 164	0	0
9	С	126	Total O 126 126	0	0
9	D	204	Total O 204 204	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	59.47Å 153.09Å 109.15Å	Depositor
a, b, c, α , β , γ	90.00° 90.80° 90.00°	Depositor
Resolution (Å)	88.87 - 2.01	Depositor
% Data completeness	98.4 (88.87-2.01)	Depositor
(in resolution range)	,	
R_{merge}	0.17	Depositor
R_{sym}	0.17	Depositor
$< I/\sigma(I) > 1$	1.33 (at 2.02Å)	Xtriage
Refinement program	PHENIX 1.11.1-2575_1496	Depositor
R, R_{free}	0.206 , 0.257	Depositor
Wilson B-factor (\mathring{A}^2)	31.0	Xtriage
Anisotropy	0.747	Xtriage
L-test for twinning ²	$< L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	0.087 for h,-k,-l	Xtriage
Total number of atoms	13985	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 37 ligands modelled in this entry, 14 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	В	802	1	41,50,50	1.51	7 (17%)	45,82,82	1.83	12 (26%)
2	HEM	D	501	1	41,50,50	1.61	7 (17%)	45,82,82	1.75	13 (28%)
2	HEM	С	501	1	41,50,50	1.66	8 (19%)	45,82,82	1.76	11 (24%)
4	BTB	С	505	_	13,13,13	0.38	0	7,16,16	0.43	0
2	HEM	A	501	1	41,50,50	1.51	6 (14%)	45,82,82	1.61	11 (24%)
3	KL4	A	509	-	25,26,26	1.21	3 (12%)	32,35,35	1.27	4 (12%)
3	KL4	С	503	-	25,26,26	1.20	3 (12%)	32,35,35	1.36	4 (12%)
4	BTB	A	504	-	13,13,13	0.40	0	7,16,16	0.61	0
4	BTB	В	805	-	13,13,13	0.42	0	7,16,16	0.65	0
3	KL4	С	502	-	25,26,26	1.19	3 (12%)	32,35,35	1.55	6 (18%)
6	GOL	A	506	-	5,5,5	0.34	0	5,5,5	0.40	0
3	KL4	С	511	-	25,26,26	1.19	3 (12%)	32,35,35	1.39	4 (12%)
4	BTB	В	804	8	13,13,13	0.52	0	7,16,16	0.72	0
6	GOL	С	507	-	5,5,5	0.39	0	5,5,5	0.34	0
4	BTB	D	503	8	13,13,13	0.43	0	7,16,16	0.57	0
4	BTB	D	504	-	13,13,13	0.50	0	7,16,16	0.68	0
4	ВТВ	В	808	-	13,13,13	0.83	1 (7%)	7,16,16	0.98	1 (14%)
3	KL4	В	801	-	25,26,26	1.22	3 (12%)	32,35,35	1.32	4 (12%)
4	BTB	A	503	-	13,13,13	0.39	0	7,16,16	0.52	0
3	KL4	D	502	_	25,26,26	1.29	3 (12%)	32,35,35	1.63	8 (25%)
4	BTB	С	504	8	13,13,13	0.41	0	7,16,16	0.51	0
3	KL4	В	803	-	25,26,26	1.23	3 (12%)	32,35,35	1.85	8 (25%)
3	KL4	A	502	-	25,26,26	1.22	3 (12%)	32,35,35	1.52	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	В	802	1	-	4/12/54/54	-
2	HEM	D	501	1	-	2/12/54/54	-
2	HEM	С	501	1	-	6/12/54/54	-
4	BTB	С	505	-	-	6/21/21/21	-
2	HEM	A	501	1	-	4/12/54/54	-

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KL4	A	509	-	-	2/10/17/17	0/3/3/3
3	KL4	С	503	-	-	1/10/17/17	0/3/3/3
4	BTB	A	504	-	-	8/21/21/21	-
4	BTB	В	805	-	-	7/21/21/21	-
3	KL4	С	502	-	-	4/10/17/17	0/3/3/3
6	GOL	A	506	-	-	4/4/4/4	-
3	KL4	С	511	-	-	3/10/17/17	0/3/3/3
4	BTB	В	804	8	-	4/21/21/21	-
6	GOL	С	507	-	-	4/4/4/4	-
4	BTB	D	503	8	-	7/21/21/21	-
4	BTB	D	504	-	-	11/21/21/21	-
4	BTB	В	808	-	-	7/21/21/21	-
3	KL4	В	801	-	-	3/10/17/17	0/3/3/3
4	BTB	A	503	-	-	11/21/21/21	-
3	KL4	D	502	-	-	4/10/17/17	0/3/3/3
4	ВТВ	С	504	8	-	7/21/21/21	-
3	KL4	В	803	-	-	2/10/17/17	0/3/3/3
3	KL4	A	502	-	-	3/10/17/17	0/3/3/3

The worst 5 of 53 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
2	D	501	HEM	C3C-C2C	-4.13	1.34	1.40
2	В	802				1.56	1.47
2	С	501	HEM	C3C-CAC	3.99	1.56	1.47
3	D	502	KL4	C25-N21	-3.80	1.35	1.49
2	D	501	HEM	C3C-CAC	3.76	1.55	1.47

The worst 5 of 89 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	С	501	HEM	CMA-C3A-C4A	-5.33	120.27	128.46
3	D	502	KL4	C02-N01-C06	4.96	121.86	118.10
2	В	802	HEM	CMA-C3A-C4A	-4.91	120.92	128.46
3	A	502	KL4	C02-N01-C06	4.77	121.72	118.10
2	D	501	HEM	CMA-C3A-C4A	-4.67	121.29	128.46

There are no chirality outliers.



5 of 114 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	HEM	C1A-C2A-CAA-CBA
2	A	501	HEM	C3A-C2A-CAA-CBA
2	В	802	HEM	C1A-C2A-CAA-CBA
2	В	802	HEM	C3A-C2A-CAA-CBA
2	С	501	HEM	C1A-C2A-CAA-CBA

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















































4.7 Other polymers (i)

There are no such residues in this entry.



4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

