

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 3, 2023 – 05:15 AM EDT

PDB ID	:	6OCX
Title	:	Structure of human CIB1 in complex with peptide inhibitor UNC10245109
Authors	:	Puhl, A.C.; Godoy, A.S.; Pearce, K.
Deposited on		
Resolution	:	1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	FAILED
:	1.13
:	FAILED
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	Engh & Huber $(2001)$
:	Parkinson et al. (1996)
:	2.35.1
	: : : :

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\hbox{-}RAY\,DIFFRACTION$ 

The reported resolution of this entry is 1.90 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



#### 60CX

# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5802 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	150	Total	С	Ν	0	S	1	2	0
	А	159	1229	780	210	235	4	1		
1	В	159	Total	С	Ν	0	S	0	1	0
	I D	109	1227	778	210	235	4			
1	С	C 159	Total	С	Ν	0	S	0	1	0
		159	1214	772	207	231	4	0	1	0
1 D	D 159	Total	С	Ν	0	S	1	1	0	
		1231	781	211	235	4	L		0	

• Molecule 1 is a protein called Calcium and integrin-binding protein 1.

• Molecule 2 is a protein called Peptide inhibitor UNC10245109.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	2 F	15	Total	С	Ν	Ο	$\mathbf{S}$	0	1	0
	Г		129	85	23	20	1	0		
2	Н	15	Total	С	Ν	Ο	S	0	1	0
	2 П	10	129	85	23	20	1			
2	т	J 15	Total	С	Ν	0	S	0	1	0
	J		129	85	23	20	1			
2 L	15	Total	С	Ν	Ο	S	0	1	0	
	L	L	15	129	85	23	20	1	0	

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	2	Total Ca 2 2	0	0
3	В	2	Total Ca 2 2	0	0
3	С	2	Total Ca 2 2	0	0
3	D	2	Total Ca 2 2	0	0



• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	85	Total O 85 85	0	0
4	В	81	Total O 81 81	0	0
4	С	81	Total O 81 81	0	0
4	D	107	Total O 107 107	0	0
4	F	8	Total O 8 8	0	0
4	Н	5	$\begin{array}{cc} \text{Total} & \text{O} \\ 5 & 5 \end{array}$	0	0
4	J	6	Total O 6 6	0	0
4	L	4	Total O 4 4	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



# 3 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	75.87Å 33.03Å 162.40Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $99.85^{\circ}$ $90.00^{\circ}$	Depositor	
Resolution (Å)	27.64 - 1.90	Depositor	
% Data completeness	99.5 (27.64-1.90)	Depositor	
(in resolution range)		Depositor	
$R_{merge}$	(Not available)	Depositor	
R <sub>sym</sub>	0.07	Depositor	
$< I/\sigma(I) > 1$	$1.59 (at 1.91 \text{\AA})$	Xtriage	
Refinement program	BUSTER 2.10.3	Depositor	
$R, R_{free}$	0.212 , $0.246$	Depositor	
Wilson B-factor $(Å^2)$	29.7	Xtriage	
Anisotropy	0.338	Xtriage	
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	5802	wwPDB-VP	
Average B, all atoms $(Å^2)$	37.0	wwPDB-VP	

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 42.43 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.0422e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

### 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 4.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 4.7 Other polymers (i)

There are no such residues in this entry.

## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 5 Fit of model and data (i)

## 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

## 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

