



# wwPDB X-ray Structure Validation Summary Report i

Jun 24, 2025 – 05:11 pm BST

PDB ID : 6R5O / pdb\_00006r5o  
Title : The crystal structure the Glycoside Hydrolase BglX inactive mutant D286N from *P. aeruginosa* in complex with two glucose molecules  
Authors : Batuecas, M.T.; Hermoso, J.A.  
Deposited on : 2019-03-25  
Resolution : 2.40 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

### X-RAY DIFFRACTION


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 11721 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Periplasmic beta-glucosidase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	B	730	Total	C	N	O	S		
			5631	3538	1011	1061	21	0	5
1	A	730	Total	C	N	O	S		
			5626	3534	1011	1061	20	0	4

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	286	ASN	ASP	engineered mutation	UNP Q9I311
A	286	ASN	ASP	engineered mutation	UNP Q9I311

- Molecule 2 is beta-D-glucopyranose (CCD ID: BGC) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			12	6	6		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 12 6 6	0	0
2	A	1	Total C O 12 6 6	0	0
2	A	1	Total C O 12 6 6	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	205	Total O 205 205	0	0
4	A	209	Total O 209 209	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Periplasmic beta-glucosidase



- Molecule 1: Periplasmic beta-glucosidase





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.50 Å 74.91 Å 82.10 Å 65.64° 74.18° 68.90°	Depositor
Resolution (Å)	46.72 – 2.40 46.72 – 2.40	Depositor EDS
% Data completeness (in resolution range)	83.9 (46.72-2.40) 83.9 (46.72-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.11 (at 2.39 Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
$R$ , $R_{free}$	0.238, 0.276 0.242, 0.284	Depositor DCC
$R_{free}$ test set	2315 reflections (4.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.9	Xtriage
Anisotropy	0.470	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.25, 20.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	11721	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.08	0/5747	1.44	1/7785 (0.0%)
1	B	1.07	0/5755	1.44	3/7794 (0.0%)
All	All	1.08	0/11502	1.44	4/15579 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	172	GLY	CA-C-O	-6.38	115.90	122.28
1	A	172	GLY	CA-C-O	-6.02	116.26	122.28
1	B	657	LEU	CA-C-N	5.67	128.19	120.54
1	B	657	LEU	C-N-CA	5.67	128.19	120.54

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5626	0	5622	99	0
1	B	5631	0	5631	93	0
2	A	24	0	24	0	0
2	B	24	0	24	4	0
3	A	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
4	A	209	0	0	32	1
4	B	205	0	0	28	1
All	All	11721	0	11301	190	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 190 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:MET:HA	4:A:904:HOH:O	1.26	1.23
1:B:63:HIS:O	4:B:901:HOH:O	1.55	1.22
1:B:59:VAL:HA	4:B:901:HOH:O	1.56	1.00
1:B:150:LEU:O	4:B:902:HOH:O	1.80	0.97
1:A:479:GLU:OE1	1:A:512:GLN:NE2	2.02	0.92

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:1101:HOH:O	4:A:1072:HOH:O[1_546]	2.13	0.07

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	730/733 (100%)	678 (93%)	49 (7%)	3 (0%)	30 44
1	B	731/733 (100%)	671 (92%)	57 (8%)	3 (0%)	30 44
All	All	1461/1466 (100%)	1349 (92%)	106 (7%)	6 (0%)	30 44

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	373	PHE
1	B	461	ALA
1	B	754	ALA
1	A	154	PHE
1	A	461	ALA

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	587/586 (100%)	572 (97%)	15 (3%)	41 62
1	B	588/586 (100%)	570 (97%)	18 (3%)	35 56
All	All	1175/1172 (100%)	1142 (97%)	33 (3%)	38 59

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	486	SER
1	A	550	ASN
1	A	757	GLU
1	B	407	LYS
1	B	403	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	376	ASN
1	A	550	ASN
1	A	738	HIS
1	A	560	GLN
1	B	562	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	BGC	A	802	-	12,12,12	0.60	0	17,17,17	1.54	4 (23%)
2	BGC	B	802	-	12,12,12	0.86	1 (8%)	17,17,17	1.33	2 (11%)
2	BGC	B	801	-	12,12,12	0.86	0	17,17,17	1.87	5 (29%)
2	BGC	A	801	-	12,12,12	0.87	0	17,17,17	1.09	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	802	-	-	2/2/22/22	0/1/1/1
2	BGC	B	802	-	-	2/2/22/22	0/1/1/1
2	BGC	B	801	-	-	0/2/22/22	0/1/1/1
2	BGC	A	801	-	-	2/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	802	BGC	O1-C1	2.00	1.46	1.39

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	BGC	O2-C2-C1	4.31	119.16	109.16
2	B	801	BGC	C1-O5-C5	-3.12	107.78	113.66
2	A	802	BGC	C1-C2-C3	-2.81	104.49	110.31
2	A	802	BGC	C6-C5-C4	-2.68	106.72	113.00
2	B	802	BGC	O5-C5-C6	2.65	113.04	106.44

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	802	BGC	C4-C5-C6-O6
2	B	802	BGC	O5-C5-C6-O6
2	A	802	BGC	C4-C5-C6-O6
2	A	802	BGC	O5-C5-C6-O6
2	A	801	BGC	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	802	BGC	1	0
2	B	801	BGC	3	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	730/733 (99%)	0.56	27 (3%) 45 43	26, 51, 75, 96	4 (0%)
1	B	730/733 (99%)	0.57	29 (3%) 43 40	23, 51, 72, 120	5 (0%)
All	All	1460/1466 (99%)	0.56	56 (3%) 44 42	23, 51, 74, 120	9 (0%)

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	418	LEU	4.3
1	B	204	ALA	4.0
1	A	623	PRO	3.6
1	B	515	ALA	3.6
1	A	205	ALA	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i

There are no oligosaccharides in this entry.

### 6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BGC	A	802	12/12	0.87	0.09	41,50,55,56	0
2	BGC	B	801	12/12	0.88	0.09	32,36,39,44	0
2	BGC	B	802	12/12	0.89	0.08	34,41,44,44	0
2	BGC	A	801	12/12	0.92	0.07	32,36,40,42	0
3	MG	B	803	1/1	0.96	0.09	61,61,61,61	0
3	MG	A	803	1/1	0.98	0.09	40,40,40,40	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.