

wwPDB X-ray Structure Validation Summary Report (i)

Nov 10, 2020 – 11:19 AM GMT

PDB ID : 6SP8

Title : Structure of hyperstable haloalkane dehalogenase variant DhaA115 prepared

by the 'soak-and-freeze' method under 150 bar of krypton pressure

Authors: Chmelova, K.; Markova, K.; Damborsky, J.; Marek, M.

Deposited on : 2019-08-31

Resolution : 1.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.14.6 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.14.6

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	291	90%	10%
1	В	291	89%	11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	KR	A	301	-	-	X	-
2	KR	A	302	-	-	X	-
2	KR	A	304	-	-	X	-
2	KR	A	307	-	-	X	-
2	KR	A	308	-	-	X	-
2	KR	A	312	-	-	X	-
2	KR	В	301	-	-	X	-
2	KR	В	302	-	-	X	-
2	KR	В	305	-	-	X	-
2	KR	В	306	-	-	X	X
2	KR	В	307	-	_	X	-
2	KR	В	308	-	-	X	-
3	SCN	В	310	-	-	X	
5	GOL	В	317	_	_	X	_



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5574 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Haloalkane dehalogenase.

\mathbf{Mol}	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
1	Δ	201	Total C		N	V O S		0	5	0
1	Λ	291	2416	1584	410	416	6	0	9	
1	B	291	Total	С	N	О	S	0	5	0
T	D	$\Delta \vartheta 1$	2411	1580	405	419	7	0	9	

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	20	SER	GLU	engineered mutation	UNP P0A3G2
A	80	ARG	PHE	engineered mutation	UNP P0A3G2
A	128	PHE	CYS	engineered mutation	UNP P0A3G2
A	148	LEU	THR	engineered mutation	UNP P0A3G2
A	155	PRO	ALA	engineered mutation	UNP P0A3G2
A	172	ILE	ALA	engineered mutation	UNP P0A3G2
A	176	PHE	CYS	engineered mutation	UNP P0A3G2
A	198	TRP	ASP	engineered mutation	UNP P0A3G2
A	219	TRP	VAL	engineered mutation	UNP P0A3G2
A	262	LEU	CYS	engineered mutation	UNP P0A3G2
A	266	PHE	ASP	engineered mutation	UNP P0A3G2
В	20	SER	GLU	engineered mutation	UNP P0A3G2
В	80	ARG	PHE	engineered mutation	UNP P0A3G2
В	128	PHE	CYS	engineered mutation	UNP P0A3G2
В	148	LEU	THR	engineered mutation	UNP P0A3G2
В	155	PRO	ALA	engineered mutation	UNP P0A3G2
В	172	ILE	ALA	engineered mutation	UNP P0A3G2
В	176	PHE	CYS	engineered mutation	UNP P0A3G2
В	198	TRP	ASP	engineered mutation	UNP P0A3G2
В	219	TRP	VAL	engineered mutation	UNP P0A3G2
В	262	LEU	CYS	engineered mutation	UNP P0A3G2
В	266	PHE	ASP	engineered mutation	UNP P0A3G2

• Molecule 2 is KRYPTON (three-letter code: KR) (formula: Kr) (labeled as "Ligand of Interest" by author).



\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
2	В	9	Total Kr 9 9	0	0
2	A	12	Total Kr 12 12	0	0


• Molecule 3 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
3	A	1	Total	С	N	S	0	0
J 3	A	1	3	1	1	1	0	U
3	A	1	Total	С	N	S	0	0
)	Λ	1	3	1	1	1	U	0
3	Λ	1	Total	С	N	S	0	0
	A	T	3	1	1	1	U	U
3	В	1	1 Total C N S	0	0			
	Ъ	T	3	1	1	1	U	U
3	В	1	Total	С	N	S	0	0
	Ъ	T	3	1	1	1	U	U
3	В	1	Total	С	N	S	0	0
	ם	I.	3	1	1	1		U


• Molecule 4 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYL AMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: B3P) (formula: $C_{11}H_{26}N_2O_6$) (labeled as "Ligand of Interest" by author).





Mol	Chain	Residues	A	ton	ıs		ZeroOcc	AltConf
1	Λ.	1	Total	С	N	О	0	0
4	A	1	19	11	2	6		U
1	D	1	Total	С	N	О	0	0
4	Б	1	19	11	2	6	U	U

• Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	В	1	Total C O 6 3 3	0	0
5	В	1	Total C O 6 3 3	0	0
5	В	1	Total C O 6 3 3	0	0
5	В	1	Total C O 6 3 3	0	0
5	В	1	Total C O 6 3 3	0	0
5	В	1	Total C O 6 3 3	0	0

• Molecule 6 is water.


Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
6	A	319	Total O 323 323	0	4
6	В	298	Total O 299 299	0	1



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Haloalkane dehalogenase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	67.98Å 82.05Å 144.18Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.20 - 1.55	Depositor
Resolution (A)	49.20 - 1.55	EDS
% Data completeness	98.8 (49.20-1.55)	Depositor
(in resolution range)	98.8 (49.20-1.55)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.13 (at 1.55Å)	Xtriage
Refinement program	PHENIX	Depositor
D D.	0.160 , 0.181	Depositor
R, R_{free}	0.161 , 0.163	DCC
R_{free} test set	5763 reflections $(4.97%)$	wwPDB-VP
Wilson B-factor (Å ²)	18.3	Xtriage
Anisotropy	0.839	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.39, 48.5	EDS
L-test for twinning ²	$ < L > = 0.53, < L^2> = 0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5574	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 24.94 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.4808e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SCN, B3P, KR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.34	0/2504	0.53	0/3426	
1	В	0.35	0/2499	0.54	0/3421	
All	All	0.34	0/5003	0.54	0/6847	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2416	0	2359	27	0
1	В	2411	0	2344	36	0
2	A	12	0	0	19	0
2	В	9	0	0	25	0
3	A	9	0	0	1	0
3	В	9	0	0	2	0
4	A	19	0	26	0	0
4	В	19	0	26	0	0
5	A	12	0	16	0	0
5	В	36	0	48	8	0
6	A	323	0	0	4	0

Continued on next page...



6SP8

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	В	299	0	0	3	0
All	All	5574	0	4819	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 70 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$egin{array}{c} ext{Clash} \ ext{overlap } (ext{Å}) \end{array}$
2:A:303:KR:KR	2:A:304:KR:KR	2.94	1.08
2:B:302:KR:KR	2:B:303:KR:KR	3.03	0.98
1:B:221:LEU:HB3	2:B:307:KR:KR	2.40	0.83
1:A:208:GLU:HG3	2:A:308:KR:KR	2.42	0.80
1:B:208[A]:GLU:OE2	2:B:306:KR:KR	2.63	0.77

There are no symmetry-related clashes.

Torsion angles (i) 5.3

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	$294/291 \; (101\%)$	284 (97%)	10 (3%)	0	100	100
1	В	$294/291 \; (101\%)$	283 (96%)	11 (4%)	0	100	100
All	All	588/582 (101%)	567 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$256/251 \; (102\%)$	254 (99%)	2 (1%)	81 66
1	В	$256/251 \; (102\%)$	254 (99%)	2 (1%)	81 66
All	All	$512/502 \; (102\%)$	508 (99%)	4 (1%)	81 66

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	168	PHE
1	A	225	TYR
1	В	168	PHE
1	В	225	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 37 ligands modelled in this entry, 21 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
WIOI	туре	Chain	res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SCN	A	313	-	1,2,2	1.14	0	0,1,1	0.00	-
3	SCN	A	314	-	1,2,2	0.57	0	0,1,1	0.00	-
3	SCN	В	312	-	1,2,2	0.37	0	0,1,1	0.00	-
5	GOL	В	317	-	5,5,5	0.48	0	5, 5, 5	0.33	0
5	GOL	В	319	-	5,5,5	0.62	0	5,5,5	0.38	0
5	GOL	В	314	-	5,5,5	0.51	0	5, 5, 5	0.19	0
3	SCN	A	315	ı	1,2,2	0.35	0	0,1,1	0.00	-
4	ВЗР	В	313	_	18,18,18	0.69	0	21,23,23	1.07	2 (9%)
4	ВЗР	A	316	-	18,18,18	0.63	0	21,23,23	1.31	1 (4%)
5	GOL	A	317	-	5,5,5	0.54	0	5, 5, 5	0.34	0
5	GOL	A	318	ı	5,5,5	0.53	0	5, 5, 5	0.28	0
5	GOL	В	315	_	5,5,5	0.54	0	5, 5, 5	0.38	0
3	SCN	В	310	_	1,2,2	1.50	0	0,1,1	0.00	_
5	GOL	В	316		5,5,5	0.47	0	5, 5, 5	0.73	0
5	GOL	В	318	_	5,5,5	0.52	0	5,5,5	0.31	0
3	SCN	В	311	-	1,2,2	0.48	0	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	В	317	_	_	2/4/4/4	_
5	GOL	В	319	-	-	3/4/4/4	-
5	GOL	В	314	-	-	0/4/4/4	-
4	ВЗР	В	313	_	-	0/28/28/28	-
5	GOL	A	317	_	_	4/4/4/4	_
5	GOL	A	318	-	-	0/4/4/4	-
5	GOL	В	315	_	-	0/4/4/4	-
4	ВЗР	A	316	_	-	2/28/28/28	_
5	GOL	В	316	-	-	0/4/4/4	-
5	GOL	В	318	_	-	2/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
4	A	316	B3P	C2-N2-C8	-3.85	110.61	116.08
4	В	313	B3P	C2-N2-C8	-2.26	112.88	116.08
4	В	313	B3P	C3-N1-C4	-2.18	112.99	116.08

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	В	317	GOL	C1-C2-C3-O3
5	A	317	GOL	C1-C2-C3-O3
5	В	318	GOL	O1-C1-C2-C3
5	В	319	GOL	C1-C2-C3-O3
5	В	319	GOL	O2-C2-C3-O3


There are no ring outliers.


6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	313	SCN	1	0
5	В	317	GOL	4	0
5	В	319	GOL	2	0
3	В	310	SCN	2	0
5	В	316	GOL	1	0
5	В	318	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	$291/291 \; (100\%)$	-0.21	4 (1%) 75 80	12, 17, 30, 51	0
1	В	291/291 (100%)	-0.23	4 (1%) 75 80	12, 18, 31, 52	0
All	All	$582/582 \; (100\%)$	-0.22	8 (1%) 75 80	12, 18, 31, 52	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	${f Res}$	Type	RSRZ
1	В	3	GLU	4.6
1	В	292	ALA	3.4
1	A	3	GLU	3.0
1	В	144	PHE	2.9
1	В	143	GLU	2.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	KR	В	306	1/1	0.49	0.49	19,19,19,19	1
2	KR	В	307	1/1	0.71	0.38	25,25,25,25	1
5	GOL	A	317	6/6	0.72	0.20	36,43,46,48	0
4	B3P	A	316	19/19	0.78	0.15	19,24,37,45	19
5	GOL	В	317	6/6	0.78	0.22	33,35,40,45	0
2	KR	A	308	1/1	0.81	0.36	25,25,25,25	1
4	B3P	В	313	19/19	0.81	0.14	19,25,41,45	19
5	GOL	В	318	6/6	0.83	0.26	23,35,41,49	0
5	GOL	A	318	6/6	0.83	0.26	24,36,47,54	0
5	GOL	В	319	6/6	0.84	0.32	30,34,38,41	0
2	KR	A	312	1/1	0.85	0.15	36,36,36,36	1
2	KR	В	308	1/1	0.85	0.17	27,27,27,27	1
2	KR	A	307	1/1	0.86	0.19	30,30,30,30	1
5	GOL	В	314	6/6	0.87	0.12	25,26,35,35	0
5	GOL	В	316	6/6	0.87	0.33	22,37,42,42	0
5	GOL	В	315	6/6	0.90	0.23	31,34,38,44	0
2	KR	В	302	1/1	0.91	0.17	24,24,24,24	1
2	KR	A	304	1/1	0.91	0.21	23,23,23,23	1
3	SCN	В	312	3/3	0.95	0.25	39,39,47,57	0
2	KR	В	309	1/1	0.95	0.13	30,30,30,30	1
2	KR	В	304	1/1	0.96	0.10	22,22,22,22	1
2	KR	A	311	1/1	0.97	0.07	28,28,28,28	1
2	KR	A	310	1/1	0.98	0.08	21,21,21,21	1
2	KR	A	305	1/1	0.98	0.07	35,35,35,35	1
2	KR	В	301	1/1	0.98	0.08	15,15,15,15	1
3	SCN	В	311	3/3	0.98	0.10	22,22,33,37	3
2	KR	В	303	1/1	0.98	0.07	23,23,23,23	1
3	SCN	A	315	3/3	0.98	0.21	28,28,31,40	0
2	KR	A	302	1/1	0.98	0.14	16,16,16,16	1
2	KR	В	305	1/1	0.98	0.10	18,18,18,18	1
2	KR	A	303	1/1	0.98	0.08	22,22,22,22	1
2	KR	A	301	1/1	0.99	0.14	18,18,18,18	1
2	KR	A	309	1/1	0.99	0.07	19,19,19,19	1
3	SCN	В	310	3/3	0.99	0.09	18,18,20,24	0
2	KR	A	306	1/1	0.99	0.13	19,19,19,19	1
3	SCN	A	313	3/3	0.99	0.07	18,18,20,25	0
3	SCN	A	314	3/3	0.99	0.12	21,21,34,37	3


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





Electron density around KR B 306: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around KR B 307:













Electron density around GOL A 317: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)














Electron density around KR A 308:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around B3P B 313:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around GOL B 318:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





















Electron density around KR A 312:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around KR B 308: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o ext{-}{ m DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



















Electron density around GOL B 315: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around KR B 302:











Electron density around KR A 304: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o ext{-}{ m DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SCN B 312:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













Electron density around KR B 309:
















Electron density around KR A 311: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c \ (\mathrm{at}\ 0.7\ \mathrm{rmsd}) \ \mathrm{in}\ \mathrm{gray}$

 ${
m mF}_o{
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around KR A 310: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)









Electron density around KR B 301:















Electron density around SCN B 311:
















Electron density around SCN A 315:













Electron density around KR A 302:






















Electron density around KR A 301: $\,$

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
























Electron density around SCN A 313:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around SCN A 314:









6.5 Other polymers (i)

There are no such residues in this entry.

