



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 12:57 PM BST

PDB ID : 6TBL
Title : Crystal structure of MMS19(CTD)-CIAO1-CIAO2B CIA targeting complex
Authors : Kassube, S.A.; Thoma, N.H.
Deposited on : 2019-11-01
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	125	
1	B	125	
2	C	159	
2	D	159	
3	E	338	
3	F	338	

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 9476 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MMS19 nucleotide excision repair protein homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	119	917	589	157	165	6	0	1	0
1	B	120	925	595	158	166	6	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	907	GLY	-	expression tag	UNP Q9D071
A	908	GLY	-	expression tag	UNP Q9D071
A	909	GLY	-	expression tag	UNP Q9D071
A	910	ARG	-	expression tag	UNP Q9D071
B	907	GLY	-	expression tag	UNP Q9D071
B	908	GLY	-	expression tag	UNP Q9D071
B	909	GLY	-	expression tag	UNP Q9D071
B	910	ARG	-	expression tag	UNP Q9D071

- Molecule 2 is a protein called MIP18 family protein galla-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	150	1195	750	215	227	3	0	1	0
2	D	151	1199	752	216	228	3	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP Q9VTC4
C	-1	GLY	-	expression tag	UNP Q9VTC4
C	0	GLY	-	expression tag	UNP Q9VTC4
C	1	ARG	-	expression tag	UNP Q9VTC4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	GLY	-	expression tag	UNP Q9VTC4
D	-1	GLY	-	expression tag	UNP Q9VTC4
D	0	GLY	-	expression tag	UNP Q9VTC4
D	1	ARG	-	expression tag	UNP Q9VTC4

- Molecule 3 is a protein called Probable cytosolic iron-sulfur protein assembly protein Ciao1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	334	Total	C	N	O	S	0	0	0
			2598	1621	450	514	13			
3	F	336	Total	C	N	O	S	0	1	0
			2618	1632	456	517	13			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-2	GLY	-	expression tag	UNP Q7K1Y4
E	-1	GLY	-	expression tag	UNP Q7K1Y4
E	0	GLY	-	expression tag	UNP Q7K1Y4
E	1	ARG	-	expression tag	UNP Q7K1Y4
F	-2	GLY	-	expression tag	UNP Q7K1Y4
F	-1	GLY	-	expression tag	UNP Q7K1Y4
F	0	GLY	-	expression tag	UNP Q7K1Y4
F	1	ARG	-	expression tag	UNP Q7K1Y4

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	K	0	0
			1	1		
4	A	1	Total	K	0	0
			1	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total O 1 1	0	0
6	C	2	Total O 2 2	0	0
6	D	6	Total O 6 6	0	0
6	F	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MMS19 nucleotide excision repair protein homolog




- Molecule 1: MMS19 nucleotide excision repair protein homolog




- Molecule 2: MIP18 family protein galla-2




- Molecule 2: MIP18 family protein galla-2





- Molecule 3: Probable cytosolic iron-sulfur protein assembly protein Ciao1





- Molecule 3: Probable cytosolic iron-sulfur protein assembly protein Ciao1



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	144.07Å 144.07Å 166.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.99 – 2.65 19.99 – 2.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.99-2.65) 100.0 (19.99-2.65)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 2.67Å)	Xtrriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.198 , 0.242 0.198 , 0.242	Depositor DCC
R_{free} test set	2567 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	81.7	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 55.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9476	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/935	0.38	0/1268
1	B	0.23	0/943	0.39	0/1280
2	C	0.24	0/1218	0.39	0/1656
2	D	0.23	0/1222	0.39	0/1661
3	E	0.24	0/2664	0.46	0/3630
3	F	0.25	0/2687	0.46	0/3660
All	All	0.24	0/9669	0.43	0/13155

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	917	0	988	14	0
1	B	925	0	998	11	0
2	C	1195	0	1218	11	0
2	D	1199	0	1221	12	0
3	E	2598	0	2461	28	0
3	F	2618	0	2486	28	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
5	C	8	0	12	2	0
5	D	4	0	6	1	0
6	B	1	0	0	0	0
6	C	2	0	0	0	0
6	D	6	0	0	0	0
6	F	1	0	0	0	0
All	All	9476	0	9390	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:225:HIS:HA	3:F:239:THR:HG22	1.67	0.77
2:C:74:SER:HB3	5:C:202:EDO:H11	1.70	0.72
3:E:228:ASN:ND2	3:E:232:VAL:O	2.23	0.71
2:C:76:HIS:HB2	5:C:202:EDO:H21	1.74	0.70
2:C:73:ASN:OD1	2:C:107:ARG:NH2	2.28	0.67
3:F:77:ALA:HB2	3:F:109:VAL:HG22	1.77	0.66
2:C:109:LYS:NZ	3:E:305:ASP:OD1	2.30	0.65
1:B:1013:ARG:NH2	2:D:32:ASP:O	2.32	0.62
1:A:1009:LYS:HB2	1:A:1012:VAL:HG12	1.82	0.61
3:F:210:VAL:HG22	3:F:260:TRP:CE3	2.35	0.61
3:E:210:VAL:HG22	3:E:260:TRP:CE3	2.35	0.61
1:B:983:THR:OG1	1:B:984:ARG:NH1	2.34	0.60
2:D:140:GLU:HB2	3:F:60:ARG:HH21	1.68	0.59
3:E:187:THR:HG23	3:E:238:GLN:HB3	1.83	0.58
1:A:1013:ARG:NH2	2:C:32:ASP:O	2.37	0.58
3:E:77:ALA:HB2	3:E:109:VAL:HG22	1.86	0.57
1:A:958:THR:O	1:A:962:ASN:ND2	2.37	0.57
3:F:222:ARG:HG2	3:F:224:TYR:CE1	2.40	0.56
2:D:99:LYS:HB2	2:D:152:ILE:HD12	1.87	0.56
3:E:87:SER:OG	3:E:96:ASN:OD1	2.20	0.55
2:D:70:ASP:OD2	2:D:107:ARG:NH1	2.40	0.54
1:A:1028:SER:O	3:F:46[B]:ASN:ND2	2.40	0.54
3:E:16:ARG:HG2	3:E:34:GLU:HB2	1.88	0.54
3:E:204:ALA:HA	3:E:262:LYS:HG3	1.89	0.54
3:E:113:ARG:HD3	3:E:158:PRO:O	2.07	0.54
3:F:82:THR:HG22	3:F:100:GLU:HG2	1.90	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:212:CYS:HB2	3:F:255:ILE:HB	1.90	0.53
3:F:237:GLN:CD	3:F:237:GLN:H	2.12	0.53
3:E:30:ALA:HB3	3:E:65:ILE:HD13	1.91	0.53
2:C:99:LYS:HB2	2:C:152:ILE:HD12	1.91	0.52
3:E:157:HIS:HB2	3:E:162:ILE:HB	1.90	0.52
2:D:68:ILE:HG13	2:D:75:VAL:HG22	1.92	0.52
1:A:1029:PRO:O	3:F:46[A]:ASN:ND2	2.43	0.52
3:E:185:ASP:OD1	3:E:186:CYS:N	2.42	0.51
1:A:960:PHE:CE2	1:A:978:CYS:HB3	2.46	0.51
1:B:1006:ASP:OD1	2:D:107:ARG:NH2	2.43	0.51
3:E:212:CYS:HB2	3:E:255:ILE:HB	1.92	0.51
3:F:190:LEU:HD23	3:F:221:TRP:CE3	2.46	0.51
1:A:1005:LEU:HD11	1:A:1020:ARG:HD3	1.93	0.51
3:F:109:VAL:HG13	3:F:120:THR:HG22	1.93	0.50
3:F:16:ARG:HG2	3:F:34:GLU:HB2	1.94	0.50
1:A:943:LEU:HA	1:A:984:ARG:HD2	1.94	0.50
3:F:83:THR:HB	3:F:99:LEU:HB2	1.94	0.49
3:E:4:LEU:HD23	3:E:298:GLU:HB2	1.94	0.49
2:C:68:ILE:HG13	2:C:75:VAL:HG22	1.93	0.49
3:F:203:ASP:OD2	3:F:204:ALA:N	2.43	0.49
1:B:972:ARG:NH1	1:B:1012:VAL:HG21	2.28	0.49
3:F:190:LEU:HD23	3:F:221:TRP:CD2	2.48	0.49
3:F:210:VAL:HG22	3:F:260:TRP:HE3	1.76	0.49
3:F:203:ASP:HB2	3:F:208:ARG:HB2	1.95	0.49
3:F:4:LEU:HD23	3:F:298:GLU:HB2	1.95	0.49
2:C:140:GLU:HB2	3:E:60:ARG:HH21	1.77	0.48
3:F:157:HIS:HB2	3:F:162:ILE:HB	1.95	0.48
1:B:912:LEU:HB3	1:B:913:PRO:HD3	1.95	0.48
3:E:82:THR:HG22	3:E:100:GLU:HG2	1.95	0.48
3:E:220:ILE:HB	3:E:245:CYS:HB3	1.96	0.47
1:A:994:LYS:NZ	1:A:1027:GLY:H	2.12	0.47
1:B:997:VAL:HG11	1:B:1023:TRP:CD2	2.50	0.46
2:D:12:VAL:O	2:D:16:ILE:HG12	2.15	0.46
3:E:225:HIS:HA	3:E:239:THR:HG22	1.97	0.46
3:E:190:LEU:HD23	3:E:221:TRP:CD2	2.51	0.46
3:E:66:ARG:HA	3:E:66:ARG:HD3	1.80	0.45
1:B:985:LEU:HB2	1:B:990:LEU:HD21	1.98	0.45
1:B:1010[B]:ARG:NH2	2:D:33:PRO:O	2.50	0.45
1:A:997:VAL:HG11	1:A:1023:TRP:CD2	2.52	0.45
3:F:163:LEU:HB3	3:F:175:PHE:HB2	1.98	0.45
3:E:190:LEU:HD23	3:E:221:TRP:CE3	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:79:PHE:HA	3:F:105:GLU:HB3	2.00	0.44
1:B:973:ILE:HG23	1:B:1015:GLU:HG3	2.00	0.44
2:C:13:TYR:CZ	2:C:17:LYS:HD2	2.53	0.44
3:F:73:TYR:CD1	3:F:87:SER:HB3	2.53	0.43
1:B:960:PHE:CE2	1:B:978:CYS:HB3	2.53	0.43
3:F:79:PHE:CD1	3:F:105:GLU:HB3	2.53	0.43
1:A:1025:LEU:O	1:A:1028:SER:HB2	2.18	0.43
3:E:109:VAL:HG13	3:E:120:THR:HG22	1.99	0.43
3:E:169:ASP:OD1	3:E:171:THR:OG1	2.30	0.43
3:F:58:HIS:HE2	3:F:76:SER:HG	1.61	0.43
2:C:42:LEU:O	2:C:99:LYS:NZ	2.40	0.43
3:E:285:SER:HB2	3:E:291:THR:H	1.84	0.43
1:A:1010[B]:ARG:NH1	2:C:35:ASP:OD1	2.46	0.43
2:D:141:ASN:HB3	2:D:144:LEU:HB3	2.00	0.43
2:D:17:LYS:O	2:D:20:VAL:HG22	2.19	0.43
3:E:73:TYR:CD1	3:E:87:SER:HA	2.54	0.43
3:F:135:ASP:OD2	3:F:136:ASP:N	2.52	0.42
3:E:157:HIS:CG	3:E:158:PRO:HD2	2.55	0.42
2:D:74:SER:HB3	5:D:201:EDO:H12	2.02	0.42
3:E:23:HIS:CG	3:E:24:PRO:HD2	2.55	0.41
3:F:51:LYS:HG3	3:F:52:THR:HG23	2.02	0.41
2:D:99:LYS:HD3	2:D:152:ILE:HG23	2.02	0.41
1:A:959:LYS:HA	1:A:959:LYS:HD3	1.84	0.41
3:F:213:SER:OG	3:F:214:ASP:N	2.53	0.41
3:E:222:ARG:NE	3:E:287:PRO:O	2.54	0.41
1:B:1005:LEU:HD11	1:B:1020:ARG:HD3	2.02	0.40
1:A:994:LYS:HZ2	1:A:1027:GLY:H	1.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	118/125 (94%)	118 (100%)	0	0	100	100
1	B	119/125 (95%)	119 (100%)	0	0	100	100
2	C	147/159 (92%)	147 (100%)	0	0	100	100
2	D	148/159 (93%)	148 (100%)	0	0	100	100
3	E	332/338 (98%)	321 (97%)	11 (3%)	0	100	100
3	F	335/338 (99%)	322 (96%)	13 (4%)	0	100	100
All	All	1199/1244 (96%)	1175 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/109 (98%)	107 (100%)	0	100	100
1	B	108/109 (99%)	107 (99%)	1 (1%)	78	87
2	C	138/142 (97%)	138 (100%)	0	100	100
2	D	138/142 (97%)	138 (100%)	0	100	100
3	E	283/284 (100%)	277 (98%)	6 (2%)	53	72
3	F	285/284 (100%)	281 (99%)	4 (1%)	67	81
All	All	1059/1070 (99%)	1048 (99%)	11 (1%)	76	86

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	962	ASN
3	E	109	VAL
3	E	170	ASN
3	E	178	GLU
3	E	210	VAL
3	E	229	THR
3	E	274	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	F	87	SER
3	F	109	VAL
3	F	113	ARG
3	F	210	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	C	202	-	3,3,3	0.44	0	2,2,2	0.16	0
5	EDO	D	201	-	3,3,3	0.44	0	2,2,2	0.35	0
5	EDO	C	201	-	3,3,3	0.46	0	2,2,2	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	C	202	-	-	1/1/1/1	-
5	EDO	D	201	-	-	1/1/1/1	-
5	EDO	C	201	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	202	EDO	O1-C1-C2-O2
5	D	201	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	202	EDO	2	0
5	D	201	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	119/125 (95%)	-0.15	3 (2%) 57 53	64, 88, 123, 175	0
1	B	120/125 (96%)	-0.03	5 (4%) 36 33	66, 85, 127, 198	0
2	C	150/159 (94%)	-0.12	8 (5%) 26 23	60, 74, 145, 176	0
2	D	151/159 (94%)	-0.10	9 (5%) 21 18	61, 72, 137, 172	0
3	E	334/338 (98%)	0.69	49 (14%) 2 1	77, 125, 192, 266	0
3	F	336/338 (99%)	0.23	22 (6%) 18 16	66, 97, 158, 197	0
All	All	1210/1244 (97%)	0.21	96 (7%) 12 10	60, 95, 165, 266	0

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	231	GLY	7.7
2	C	1	ARG	5.5
1	A	1031	SER	5.5
3	E	91	GLY	5.2
3	E	92	GLU	4.9
2	D	21	LEU	4.9
1	B	912	LEU	4.9
3	F	1	ARG	4.7
3	F	180	ILE	4.5
3	E	90	SER	4.5
3	F	321	SER	4.4
3	E	121	CYS	4.3
3	F	3	ARG	3.9
2	D	28	GLU	3.9
2	D	1	ARG	3.9
3	E	136	ASP	3.9
1	B	1031	SER	3.9
3	E	222	ARG	3.8
2	C	155	LYS	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	E	133	ALA	3.7
3	E	179	PRO	3.6
2	C	21	LEU	3.6
2	D	155	LYS	3.6
3	F	179	PRO	3.5
3	E	3	ARG	3.5
2	C	30	VAL	3.5
3	E	232	VAL	3.5
2	D	30	VAL	3.4
3	E	120	THR	3.4
3	E	180	ILE	3.4
3	E	30	ALA	3.4
2	D	20	VAL	3.3
3	E	60	ARG	3.3
3	F	121	CYS	3.3
3	E	59	LYS	3.3
3	F	133	ALA	3.3
3	E	237	GLN	3.2
3	E	321	SER	3.2
3	E	73	TYR	3.2
3	F	0	GLY	3.2
3	E	119	ALA	3.2
3	E	25	LYS	3.1
3	F	235	PRO	3.1
3	E	283	SER	3.1
1	B	967	TYR	3.0
2	D	-1	GLY	3.0
3	E	94	GLU	3.0
3	E	288	ASP	3.0
3	F	182	ASN	2.9
3	E	322	CYS	2.9
1	B	913	PRO	2.9
3	E	229	THR	2.8
2	D	18	GLU	2.8
3	E	235	PRO	2.8
3	F	90	SER	2.7
3	E	284	ASP	2.7
3	F	92	GLU	2.7
3	E	181	ASP	2.6
3	F	286	LYS	2.6
3	E	57	GLY	2.6
3	F	119	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	103	GLU	2.5
3	E	75	ALA	2.5
3	E	31	SER	2.5
3	F	91	GLY	2.5
3	E	221	TRP	2.5
3	E	159	THR	2.5
3	E	138	PHE	2.5
3	E	289	GLU	2.4
3	E	287	PRO	2.4
3	E	137	GLU	2.4
3	F	120	THR	2.4
2	C	18	GLU	2.4
3	E	269	THR	2.4
3	F	289	GLU	2.4
3	F	59	LYS	2.4
3	E	142	ALA	2.4
3	E	113	ARG	2.3
3	E	83	THR	2.3
1	A	967	TYR	2.3
3	F	181	ASP	2.3
2	D	15	ARG	2.2
1	B	1029	PRO	2.2
3	E	89	SER	2.1
3	E	56	ASP	2.1
2	C	11	ASN	2.1
2	C	29	ASN	2.1
3	E	319	LEU	2.1
3	E	320	ILE	2.1
2	C	0	GLY	2.1
1	A	913	PRO	2.1
3	E	286	LYS	2.1
3	F	159	THR	2.1
3	E	228	ASN	2.0
3	E	106	VAL	2.0
3	F	56	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	EDO	D	201	4/4	0.81	0.48	107,115,122,126	0
4	K	B	1101	1/1	0.85	0.09	104,104,104,104	0
5	EDO	C	202	4/4	0.87	0.33	102,102,102,103	0
4	K	A	1101	1/1	0.91	0.14	106,106,106,106	0
5	EDO	C	201	4/4	0.96	0.42	103,106,106,109	0

6.5 Other polymers [i](#)

There are no such residues in this entry.